

# New 6-in-1 vaccine study to guide UK vaccine policy

<b>Submission date</b> 02/06/2023	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 04/09/2023	<b>Overall study status</b> Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 20/05/2025	<b>Condition category</b> Other	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

A vaccine protecting against Haemophilus influenzae type b (Hib) and group C meningococcus (MenC), currently given in the UK routine immunisation programme at age 12 months, branded Menitorix, will be unavailable from 2025. The Joint Committee on Vaccination and Immunisation (JCVI) has considered options for adapting the current schedule. This study will evaluate a proposed schedule before its implementation.

The JCVI suggests that MenC vaccination is no longer required in the UK infant schedule, as a successful vaccination programme in adolescents has led to effective “herd immunity” against MenC. The JCVI also suggests a new vaccination timepoint in the routine schedule at age 18 months, to give a second dose of measles, mumps, rubella (MMR) vaccine and a booster dose of the “6-in-1” vaccine (which protects against diphtheria, tetanus, poliovirus, pertussis, hepatitis B and Hib). Moving forward the second dose of MMR vaccine (currently given at pre-school age) may improve vaccination uptake. An initial course of the 6-in-1 vaccine is currently routinely given at 2, 3 and 4 months of age. The boost to Hib immunity currently achieved with Menitorix may instead be achievable by giving the 6-in-1 vaccine at 18 months.

There are two licensed 6-in-1 vaccines, Infanrix hexa and Vaxelis, whose components differ. Currently, the same vaccine is recommended for each of the initial three doses. This study will investigate whether the vaccines may be used interchangeably for the booster, which would facilitate the delivery of the vaccine programme. In addition, the researchers will study the difference in immune response to routine vaccinations between infants born at less than 37 weeks gestational age versus those born at greater than or equal to 37 weeks gestational age receiving this vaccine schedule.

### Who can participate?

Children aged 12 months who have completed the infant UK immunisation schedule before 6 months of age

### What does the study involve?

Participants will be given the vaccinations routinely scheduled at 12 months (except Menitorix). At age 18 months, they will be given either Infanrix hexa or Vaxelis (determined by

randomisation) and MMR vaccine. Participants will also be offered two doses of chickenpox vaccine (which is not routinely given in UK) as an optional benefit. Blood samples will be taken at ages 18 and 19 months to assess immune responses to the vaccines.

What are the possible benefits and risks of participating?

The following benefits are possible when participating in this study:

Participants will be given routine vaccines in their own homes or in a suitable and convenient location.

Participants will be offered two doses of a vaccine which helps to prevent chicken pox. This vaccine is not routinely offered in the UK.

Participants will have 24-hour telephone access to an on-call study team member for urgent clinical queries related to the study.

Participants in the study will receive their booster dose of an MMR vaccine at the age of 18 months old instead of at 40 months; this may result in increased protection against measles, mumps and rubella from an earlier age. Similarly, by receiving the 6-in-1 vaccine at the age of 18 months old, participants may gain increased immunity against tetanus, diphtheria, pertussis, polio and hepatitis B.

Participants in this trial will be receiving licensed vaccines which are already included in the childhood routine UK immunisation schedule or licensed for use in the UK. The risks to participants include the common side effects of each vaccination, listed within the summary of product characteristics. These include local injection site reactions such as pain, erythema, swelling, pruritis and induration and systemic side effects including fever, irritability, rash, malaise, drowsiness, vomiting and loss of appetite. Both MMR and varicella vaccines are associated with an increased risk of febrile seizures, typically 7 to 10 days following vaccination. A delay in administration of the Hib booster from 12 to 18 months may result in reduced immunity against Hib, however, this risk is minimal due to low numbers of Haemophilus influenzae Type B cases reported in the UK.

Removal of the MenC booster from the vaccinations given at 12 months may result in reduced immunity against MenC. The risk of disease resulting from this is extremely low, given the low rates of carriage in the UK since the introduction of routine adolescent MenACWY vaccination in 2015.

As with all vaccines, there is a small chance of an allergic reaction including severe reactions such as anaphylaxis (the risk is less than one in a million doses for existing vaccines).

The risks associated with blood sampling include bleeding, pain at the injection site, bruising and infection. Anaesthetic cream is offered to minimise any pain during blood sampling. The volume of blood taken at each visit does not exceed the recommended European limits for infants.

Where is the study run from?

University of Oxford (UK)

When is the study starting and how long is it expected to run for?

May 2023 to June 2026

Who is funding the study?

MCM Vaccine (Netherlands)

Who is the main contact?

Prof. Dominic Kelly, info@ovg.ox.ac.uk (UK)

## Contact information

**Type(s)**

Public, Scientific, Principal investigator

**Contact name**

Prof Dominic Kelly

**Contact details**

Oxford Vaccine Group

Oxford

United Kingdom

OX3 7LE

+44 (0)1865 611400 ext 0000

info@ovg.ox.ac.uk

## Additional identifiers

**Clinical Trials Information System (CTIS)**

2022-003425-22

**Integrated Research Application System (IRAS)**

1006942

**Central Portfolio Management System (CPMS)**

57091

**Protocol serial number**

OVG2022/04

## Study information

**Scientific Title**

Heterologous boosting for hexavalent paediatric vaccines in the UK schedule

**Acronym**

The 6-in-1 Part 2 Vaccine Study

**Study objectives**

Primary objective:

To determine if the anti-PRP seroprotection rate in the heterologous hexavalent schedule is non-inferior to the homologous schedule at 1 month following boost with either Hex-I or Hex-V

Secondary objectives:

1. To determine if the anti-PRP seroprotection rate in the heterologous schedule is superior to the homologous schedule at 1 month following boost
2. To characterize the immunogenicity of selected antigens in Infanrix-hexa and Vaxelis at 18 months in study participants after receiving homologous Infanrix-hexa or Vaxelis as prime vaccination at 2, 3 and 4 months
3. To characterize the immunogenicity of selected antigens in Infanrix-hexa and Vaxelis at 18 and 19 months in study participants receiving either homologous or heterologous Infanrix-Hexa or Vaxelis as boost vaccination at 18 months

4. To characterize the immunogenicity of a schedule of MMR prime at 12 months and MMR boost at 18 months
5. To characterize rates of solicited and unsolicited adverse events for both heterologous and homologous schedules

Added 22/03/2024:

6. To determine if there is a difference in anti-PRP seroprotection rates after the primary vaccination series of 6-in-1 at 18 months of age in infants born at  $\geq 32$  weeks gestational age compared to infants born at  $< 32$  weeks gestational age
7. To determine if there is a difference in anti-PRP seroprotection rates at one month following the 6-in-1 booster vaccination in infants born at  $\geq 32$  weeks gestational age compared to infants born at  $< 32$  weeks gestational age
8. To characterize the immunogenicity of selected antigens at 18 months in infants born at  $\geq 32$  weeks gestational age and in infants born at  $< 32$  weeks gestational age, after receiving 6-in-1 prime vaccination at 2, 3 and 4 months
9. To characterize the immunogenicity of selected antigens at 19 months in infants born at  $\geq 32$  weeks gestational age and in infants born at  $< 32$  weeks gestational age, after receiving 6-in-1 boost vaccination at 18 months

Updated 29/01/2025:

6. To determine if there is a difference in anti-PRP seroprotection rates after the primary vaccination series of 6-in-1 at 18 months of age in infants born at  $\geq 37$  weeks gestational age compared to infants born at  $< 37$  weeks gestational age
7. To determine if there is a difference in anti-PRP seroprotection rates at one month following the 6-in-1 booster vaccination in infants born at  $\geq 37$  weeks gestational age compared to infants born at  $< 37$  weeks gestational age
8. To characterize the immunogenicity of selected antigens at 18 months in infants born at  $\geq 37$  weeks gestational age and in infants born at  $< 37$  weeks gestational age, after receiving 6-in-1 prime vaccination at 2, 3 and 4 months
9. To characterize the immunogenicity of selected antigens at 19 months in infants born at  $\geq 37$  weeks gestational age and in infants born at  $< 37$  weeks gestational age, after receiving 6-in-1 boost vaccination at 18 months

### **Ethics approval required**

Ethics approval required

### **Ethics approval(s)**

approved 30/08/2023, East of England - Cambridge South Research Ethics Committee (Equinox House, City Link, Nottingham, NG2 4LA, United Kingdom; None available; cambridgesouth.rec@hra.nhs.uk), ref: 23/EE/0121

### **Study design**

Single-blind randomized controlled trial

### **Primary study design**

Interventional

### **Study type(s)**

Prevention

### **Health condition(s) or problem(s) studied**

This study is to evaluate a proposed new vaccine schedule before its potential implementation

## **Interventions**

All participants receive the following vaccines at 12 months of age:

- Meningococcal Group B vaccine (Bexsero), 0.5 ml by intramuscular injection
- Pneumococcal polysaccharide conjugate vaccine (Prevenar 13), 0.5 ml by intramuscular injection
- Measles, mumps and rubella vaccine (Priorix or M-M-RvaxPro), 0.5 ml by intramuscular injection

Participants do not receive the Hib/MenC vaccine (Menitorix), which is currently routinely given in the UK immunisation schedule at 12 months.

All participants receive the following vaccine at 18 months of age:

- Measles, mumps and rubella vaccine (Priorix or M-M-RvaxPro), 0.5 ml by intramuscular injection

All participants are randomised to receive at 18 months of age:

EITHER hexavalent DTaP/IPV/Hib/HepB vaccine (Infanrix hexa), 0.5 ml by intramuscular injection  
OR hexavalent DTaP/IPV/Hib/HepB vaccine (Infanrix hexa), 0.5 ml by intramuscular injection

Randomisation (1:1) will be achieved by computer-generated randomisation lists, prepared by the study statistician using stratified block randomization

There will be four study groups, determined by the hexavalent vaccine received for priming in infancy (Infanrix hexa or Vaxelis) and by the hexavalent vaccine received at 18 months of age (Infanrix hexa or Vaxelis). In addition, all study participants will be offered varicella vaccine (either Varivax or Varilrix) at 12 and 18 months of age (each dose 0.5 ml, by intramuscular injection). This is optional and decided by the participant's parent/legal guardian, who may opt for 0, 1 or 2 doses of the varicella vaccine to be given.

## **Intervention Type**

Biological/Vaccine

## **Phase**

Phase IV

## **Drug/device/biological/vaccine name(s)**

Infanrix hexa, Powder and suspension for suspension for injection [Diphtheria toxoid, Tetanus toxoid, Bordetella pertussis toxoid, Bordetella pertussis Filamentous Haemagglutinin, Bordetella pertussis pertactin, Hepatitis B surface antigen, Poliovirus (inactivated) type 1 (Mahoney strain), Poliovirus (inactivated) type 2 (MEF-1 strain), Poliovirus (inactivated) type 3 (Saukett strain), Haemophilus influenzae type b polysaccharide, Tetanus toxoid (as carrier protein for Haemophilus influenzae type b polysaccharide), Aluminium hydroxide (hydrated), Aluminium phosphate] , Vaxelis [Diphtheria Toxoid, Tetanus Toxoid, Bordetella pertussis toxoid, Bordetella pertussis Filamentous Haemagglutinin, Bordetella pertussis Pertactin, Bordetella pertussis Fimbriae Types 2 and 3, Hepatitis B surface antigen, Poliovirus (Inactivated) Type 1 (Mahoney), Poliovirus (Inactivated) Type 2 (MEF-1), Poliovirus (Inactivated) Type 3 (Saukett), Haemophilus influenzae type b polysaccharide (Polyribosylribitol Phosphate), Meningococcal protein (as carrier for Haemophilus influenzae type b polysaccharide), Aluminium phosphate, Amorphous aluminium hydroxyphosphate sulfate] , Priorix [Measles virus Schwarz strain (live, attenuated), Mumps virus RIT 4385 strain, derived from Jeryl Lynn strain (live, attenuated),

Rubella virus Wistar RA 27/3 strain (live, attenuated)] , M-M-RvaxPro [Measles virus Enders' Edmonston strain (live, attenuated), Mumps virus Jeryl Lynn™ (Level B) strain (live, attenuated), Rubella virus Wistar RA 27/3 strain (live, attenuated)]

### **Primary outcome(s)**

Anti-PRP (Hib) IgG concentrations  $\geq 1.0$  mcg/ml, measured by ELISA at 19 months of age

### **Key secondary outcome(s)**

1. IgG concentrations against PRP (Hib), diphtheria toxoid, tetanus toxoid, hepatitis B surface antigen and pertussis toxoid (PT, FHA, fimbriae, pertactin), measured by ELISA at 18 and 19 months of age
2. Antibody against polio types 1, 2 and 3, measured by poliovirus binding inhibition multiplex immunoassay at 18 and 19 months of age
3. IgG concentrations against measles, mumps and rubella, measured by ELISA at 18 and 19 months of age
4. Solicited events recorded in participant diaries, from the vaccination at age 18 months until 7 days post-vaccination
5. Daily temperature recorded in participant diaries, from the vaccination at age 18 months until 21 days post-vaccination
6. Unsolicited medically attended adverse events reported by participants from the vaccination at age 18 months until 28 days post-vaccination
7. Serious adverse events (SAEs) reported by participants from the vaccination at age 12 months until the final study visit

Added 22/03/2024:

8. Anti-PRP (Hib) IgG concentrations  $\geq 1.0$   $\mu\text{g/ml}$  and anti-PRP (Hib) IgG concentrations  $\geq 0.15$   $\mu\text{g/ml}$  at 18 months of age, as measured by ELISA
9. Anti-PRP (Hib) IgG concentrations  $\geq 1.0$   $\mu\text{g/ml}$  and anti-PRP (Hib) IgG concentrations  $\geq 0.15$   $\mu\text{g/ml}$  at 19 months of age, as measured by ELISA
10. IgG concentrations at 18 months against PRP (Hib), diphtheria toxoid, tetanus toxoid, hepatitis B surface antigen, pertussis toxoid (PT, FHA, fimbriae, pertactin), measles, mumps, rubella, measured by ELISA
11. Antibody against polio types 1, 2, 3 determined by MIA\* at 18 months
12. IgG concentrations at 19 months against PRP (Hib), diphtheria toxoid, tetanus toxoid, hepatitis B surface antigen, pertussis toxoid (PT, FHA, fimbriae, pertactin), measured by ELISA
13. Antibody against polio types 1, 2, 3 determined by MIA\* at 19 months

### **Completion date**

30/06/2026

## **Eligibility**

### **Key inclusion criteria**

1. Children aged 12 months (up to 12 months +42 days), who have completed the infant UK immunisation schedule before 6 months of age [including 3 (homologous) doses of Hex-I or Hex-V]
2. Parents/legal guardians are willing and able to comply with study procedures
3. Parents/legal guardians who are over 18 years of age and willing to provide written informed consent for their child's participation in the study

Added 22/03/2024:

4. For cohort B only: children born at <32 weeks gestational age

Updated 29/01/2025:

4. For cohort B only: children born at <37 weeks gestational age

### **Participant type(s)**

Healthy volunteer

### **Healthy volunteers allowed**

No

### **Age group**

Child

### **Lower age limit**

12 months

### **Sex**

All

### **Key exclusion criteria**

Children may not participate in the study if they have:

1. Received any of the vaccines/vaccine doses scheduled for the UK '12-month' immunisation episode
2. Received any additional doses (outside the routine schedule) of any vaccines containing antigens for diphtheria, tetanus, pertussis, polio, hepatitis B, Hib, measles, mumps, rubella or varicella
3. Received influenza vaccine
4. Confirmed or suspected immunodeficiency
5. Household contacts with severe immunodeficiency
6. Any of the contraindications to the study vaccinations as specified in The Green Book
7. Confirmed anaphylactic reaction to any constituent or excipient of the study vaccines
8. Confirmed anaphylactic reaction to neomycin, streptomycin or polymyxin B (which may be present in trace amounts in the tetanus vaccine), kanamycin, histidine, sodium chloride or sucrose (which may be present in trace amounts in the Meningococcal B vaccine) or to gelatin (which may be present in trace amounts in the MMR vaccine)
9. Latex hypersensitivity (the syringe cap of the Meningococcal B vaccine Bexsero may contain natural rubber latex)
10. Any other significant disease or disorder which, in the opinion of the Investigator, may either put the participants at risk through participation in the study, or may influence the result of the study or the participant's ability to participate in the study
11. Parents/legal guardians who are on the delegation log of this study

Added 22/03/2024:

Temporary exclusion criteria

Children are temporarily excluded from participating if they:

1. Within the 28 days prior to study vaccines have received live yellow fever, varicella or any other live vaccine that requires a 28-day interval before receiving MMR or varicella vaccine
2. Have received any other vaccine within 14 days prior to study vaccines
3. Have scheduled elective surgery, planned admission or other procedures requiring general

anaesthesia within 7 days of receiving a vaccine

4. Have a febrile illness (axillary temperature  $\geq 38.0^{\circ}\text{C}$ ) within the previous 24 hours of a scheduled vaccination

5. Are currently participating in another interventional clinical trial

6. Have had a tuberculin skin test (Mantoux) which has not been read (MMR vaccine should be delayed until after the test has been read unless protection against measles, mumps and rubella is required urgently)

**Date of first enrolment**

12/09/2023

**Date of final enrolment**

31/07/2025

## **Locations**

**Countries of recruitment**

United Kingdom

England

**Study participating centre**

**Oxford Vaccine Group**

Centre for Clinical Vaccinology & Tropical Medicine (CCVTM)

Churchill Hospital

Oxford

United Kingdom

OX3 7LE

**Study participating centre**

**Bristol Royal Hospital for Children**

Bristol Vaccine Centre

Upper Maudlin St,

Bristol

United Kingdom

BS2 8BJ

**Study participating centre**

**Southampton General Hospital**

Tremona Road

Southampton

United Kingdom

SO16 6YD

**Study participating centre**

**St George's University Hospitals NHS Foundation Trust**  
Blackshaw Road  
London  
United Kingdom  
SW17 0QT

**Study participating centre****The Adam Practice**

Upton Surgery  
Upton  
Poole  
Dorset  
United Kingdom  
BH16 5PW

## Sponsor information

**Organisation**

University of Oxford

**ROR**

<https://ror.org/052gg0110>

## Funder(s)

**Funder type**

Industry

**Funder Name**

MCM Vaccine

## Results and Publications

**Individual participant data (IPD) sharing plan**

The data-sharing plans for the current study are unknown and will be made available at a later date

**IPD sharing plan summary**

Data sharing statement to be made available at a later date

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Study website</a>	Study website	11/11/2025	11/11/2025	No	Yes