

Investigating conditions causing breathlessness in athletes

Submission date 27/07/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 24/08/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 03/09/2025	Condition category Respiratory	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Exercise-induced dyspnea (shortness of breath) in athletes is common and is often attributed to asthma. However, all wheeze is not due to asthma and diagnosis and treatment should be based on careful clinical evaluation. Exercise-induced laryngeal obstruction (EILO) is an important differential diagnosis to asthma in athletes. In EILO, the larynx (voice box) is transiently obstructed during exercise resulting in decreased airflow and airway symptoms, typically with a sudden onset followed by rapid cessation when exercise is stopped. EILO is diagnosed with a continuous laryngoscopy during exercise (CLE) test, in which larynx is continuously visualized during peak exercise. This study aims to estimate the prevalence of EILO among athletes and to describe the natural history of EILO. Secondary aims of the study are to describe the prevalence of asthma and to what extent the conditions coexist.

Who can participate?

Healthy athletes at competitive level, 15-35 years of age

What does the study involve?

Participation in the study involves one or two visits at the Clinical Research Centre, where testing is done. Each visit includes a clinical examination by a physician, spirometry (lung function tests), asthma provocation test with a eucapnic voluntary hyperventilation test (EVH), skin prick test (only first visit), questionnaire, and continuous laryngoscopy during exercise (CLE) test.

What are the possible benefits and risks of participating?

The participants receive financial compensation of about £90 (SEK 1000). The pre-test requires exercise to exhaustion which may be uncomfortable for participants who do not frequently undertake high-intensity exercise. Oxygen uptake is measured using a mouthpiece and nose clip that might cause a little discomfort during exercise. Exposure to sub-zero temperatures is expected to cause short-lived cold-related symptoms. Blood sampling causes short-lived pain at the site of vein puncture. The lung function tests involve several large exhalations which may feel exhaustive in the moment.

Where is the study run from?
Clinical Research Centre, Östersund Hospital (Sweden)

When is the study starting and how long is it expected to run for?
April 2014 to November 2017

Who is funding the study?
1. VISARE NORR Fund, Northern Country Councils Regional Federation (Sweden)
2. Unit of Research, Education and Development, Region Jämtland Härjedalen (Sweden)

Who is the main contact?
Dr Nikolai Stenfors (scientific)
nikolai.stenfors@umu.se

Contact information

Type(s)
Scientific

Contact name
Dr Nikolai Stenfors

ORCID ID
<https://orcid.org/0000-0002-1684-1301>

Contact details
Östersund Hospital
Kyrkgatan 16
Östersund
Sweden
83183
+46 (0)63 153000
nikolai.stenfors@umu.se

Additional identifiers

Protocol serial number
EILO hos konditionsidrottare.

Study information

Scientific Title
Prevalence of exercise-induced laryngeal obstruction and asthma among athletes

Study objectives
Athletes have a high prevalence of exercise-induced laryngeal obstruction (EILO).

Ethics approval required
Old ethics approval format

Ethics approval(s)

Approved 26/03/2015, Regional Ethical Review Board Umea (Samverkanshuset Universitetsområdet, 901 87 Umeå, Sweden; +46 (0)90 786 7252; epn@adm.umu.se), ref: Dnr 2015-43-31M

Study design

Single-center longitudinal cohort study

Primary study design

Observational

Study type(s)

Screening

Health condition(s) or problem(s) studied

Prevalence of exercise-induced laryngeal obstruction (EILO) in athletes

Interventions

Participants undergo a clinical examination by the study physician, dynamic spirometry, eucapnic voluntary hyperventilation test, skin prick test, and continuous laryngoscopy during exercise test (CLE). CLE test includes running on a treadmill at incrementing speed to >90% maximum heart rate, while video recording larynx with a nasal fiberscope.

Intervention Type

Other

Primary outcome(s)

Prevalence and natural history of exercise-induced laryngeal obstruction (EILO), defined as laryngeal obstruction at glottic or supraglottic level using a visual scoring system, assessed with continuous laryngoscopy during exercise test (CLE), at the first and second visit (summer or winter season)

Key secondary outcome(s)

1. Current asthma, defined as a) physician-diagnosed asthma, and b) use of asthma medication in the last 12 months, evaluated by questionnaire at first visit
2. Bronchial hyperreactivity, defined as 10% decrease in FEV1 after EVH test, measured at the first and second visit (summer or winter season)
3. Allergy, defined as a) reaction with >2 mm wheal to skin prick test, and b) self-reported allergic symptoms, measured at first visit

Completion date

28/11/2017

Eligibility

Key inclusion criteria

1. Competitive athlete (i.e. not at recreational level)
2. Age 15-35 years

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

15 years

Upper age limit

35 years

Sex

All

Total final enrolment

111

Key exclusion criteria

1. Pregnancy
2. FEV1 <50% of expected value at spirometry
3. Blood pressure (mmHg) > 200 systolic or >100 diastolic at rest
4. Aortic aneurysm
5. Previous heart attack or stroke

Date of first enrolment

07/04/2015

Date of final enrolment

28/11/2017

Locations

Countries of recruitment

Sweden

Study participating centre

Clinical Research Centre, Östersund Hospital

Kyrkgatan 16

Östersund

Sweden

831 83

Sponsor information

Organisation
Umeå University

Funder(s)

Funder type
Government

Funder Name
VISARE NORR Fund, Northern Country Councils Regional Federation

Funder Name
Unit of Research, Education and Development, Region Jämtland Härjedalen

Results and Publications

Individual participant data (IPD) sharing plan

The data generated during and/or analysed during the current study are not expected to be made available due to a requirement in the ethical approval, where there is a statement that the data will not be shared with unauthorized persons. The data will be held in a server at Umeå University.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/06/2021	15/04/2021	Yes	No
Other publications	Longitudinal follow-up	15/07/2023	17/07/2023	Yes	No
Protocol file	In Swedish version 2	29/06/2020	30/08/2022	No	No