

# Markers of Aggressive Local Therapy In Newly diagnosed Glioblastomas

|                                        |                                                   |                                                                                                              |
|----------------------------------------|---------------------------------------------------|--------------------------------------------------------------------------------------------------------------|
| <b>Submission date</b><br>02/01/2010   | <b>Recruitment status</b><br>No longer recruiting | <input type="checkbox"/> Prospectively registered<br><input type="checkbox"/> Protocol                       |
| <b>Registration date</b><br>04/03/2011 | <b>Overall study status</b><br>Completed          | <input type="checkbox"/> Statistical analysis plan<br><input type="checkbox"/> Results                       |
| <b>Last Edited</b><br>20/02/2019       | <b>Condition category</b><br>Cancer               | <input type="checkbox"/> Individual participant data<br><input type="checkbox"/> Record updated in last year |

## Plain English summary of protocol

<http://cancerhelp.cancerresearchuk.org/trials/a-study-looking-how-glioma-brain-tumours-behave>

## Contact information

### Type(s)

Scientific

### Contact name

Mr Stephen Price

### Contact details

Neurosurgery Division  
Department of Clinical Neurosciences  
Box 167, Addenbrooke's Hospital  
Hills Road  
Cambridge  
United Kingdom  
CB2 0QQ  
+44 (0)12 2327 4295  
sjp58@cam.ac.uk

## Additional identifiers

### Protocol serial number

NIHR/CS/009/011

## Study information

### Scientific Title

# Magnetic resonance imaging to characterise invasive phenotypes in cerebral gliomas: an observational prospective cohort study

## Acronym

MALTING

## Study objectives

1. The spectrum of high grade gliomas invasion can be identified using diffusion tensor imaging (DTI). Combining these findings with further information relating to angiogenesis and metabolic activity provided by perfusion magnetic resonance imaging (MRI) and magnetic resonance spectroscopy (MRS), will predict patterns of recurrence and the time-to-progression.
2. Tumours with a less invasive imaging pattern will have a better response to aggressive local therapy
3. Regions of extensive invasion will exhibit a biological phenotype characterised by increased extracellular expression of matrix metalloproteinase 2 gene (MMP-2). Vascular endothelial cell growth factor (VEGF) and chemokine stromal cell derived factor-1 (CXCL12) measured using microdialysis and an increased expression of membrane type 1 metalloprotease (MT1-MMP) measured using tumour biopsies.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Not provided at time of registration

## Primary study design

Observational

## Study design

Prospective observational cohort study

## Study type(s)

Screening

## Health condition(s) or problem(s) studied

Glioblastomas

## Interventions

This study will include three patient cohorts:

1. Markers of Aggressive Local Therapies in Newly diagnosed Glioblastomas (MALTING):  
This project will involve recruiting 95 patients with glioblastomas that are felt by the surgical team to be fully resectable. Patients will be imaged pre-operatively using the standard protocol. Patients will undergo an image-guided craniotomy and tumour debulking, assisted using 5-aminolevulinic acid (5-ALA) fluorescence to improve and standardise tumour resection. Carmustine wafers (Gliadel®) will be inserted according to our local protocol. Patients will be imaged post-operatively to assess the extent of resection. Post-operatively patients will be treated with radiotherapy plus concomitant and adjuvant Temozolomide according to the Stupp protocol. All adverse events (both expected and unexpected) will be recorded in the post-operative period.

## 2. Correlation of Imaging Parameters with Biological Markers of Invasion:

This cohort will involve 50 patients who are undergoing craniotomy and tumour resection. These numbers should provide 10 - 14 limited invasion and 10 diffusely invasive patients. These studies will be performed in collaboration and will study:

2.1. Microdialysis: This will be performed in conjunction with Mr. Peter Hutchinson and Dr. Keri Carpenter, Department of Clinical Neurosciences. In 50 patients who are undergoing craniotomy and tumour resection, two CMA-71 microdialysis catheters (membrane cut-off of 100 kD) will be inserted at the time of surgery: one catheter in the most invasive region (as defined by DTI) and another in normal brain. They will be perfused at a rate of 0.3  $\mu\text{L min}^{-1}$ . After 6 hours equilibration time, samples will be taken at 8 hourly intervals over a 48 hour period. A post-op imaging will confirm catheter location. Small molecule analysis will be performed using a CMA 600 analyser to measure concentrations of glucose, pyruvate, lactate and glutamate. Urea will also be measured and used as an endogenous reference. Macromolecules will be analysed using a multiplex immunoassay and chemiluminescence detection to measure concentration of VEGF, MMP-2, TIMP-1, CXCL12). The concentration of these proteins will be correlated with imaging measures of invasion.

2.2. Cellular Measures: Using multimodal imaging we will identify surgically accessible regions of potential biological interest and these regions will be biopsied. Cell populations will be derived under serum-free conditions and assayed for tumour competency in the Watts lab using established protocols. Cell populations from multiple patient samples will be evaluated for the expression of therapeutically targetable ligands associated with tumour invasion. In particular we will focus on the expression of MT1-MMP.

2.3. High-Resolution Magic Angle Spinning NMR Spectroscopy (HR MAS): This technique images tumour samples and provides a biochemical profile of the tumour that does not destroy the tumour sample. This work will be performed in conjunction with Prof. John Griffiths, CRUK Cambridge Research Institute.

2.4. Genomic Profiling: Gene expression profiles will compare invasive with minimally invasive tumours. Samples will be stored as part of our Brain Tumour Bank and will involve storing both brain tumour samples (image-guided) with 20mls of blood.

## Intervention Type

Other

## Phase

Not Applicable

## Primary outcome(s)

Pattern of contrast enhancement at first recurrence. This will be assessed by co-registering anatomical MR's at recurrence with pre-RT and highlighting areas of new contrast enhancement. Invasive GBM's will be defined as radiological evidence of > 80% of the recurrent tumour occurring outside the radiotherapy 95% isodose. For the MALTING study, the percentage of patients surviving 2 years will be the main outcome measure.

## Key secondary outcome(s)

1. Overall survival

2. Time to radiological progression: Radiological progression is defined as per MacDonald criteria i.e. a 25% or greater increase in the size of the tumour (as defined by the product of two perpendiculars of the enhancing component) or the appearances of new contrast-enhancing lesions.

3. Time to clinical progression: This will be defined as the presence of any of the following:

3.1. Neurological deterioration with or without the need for increased steroid use

- 3.2. Increased steroid requirements for more than 2 weeks including for increasing neurological deficit and/or features of increased intracranial pressure suggestive of tumour progression when other causes have been excluded.
- 3.3. Deterioration of  $\geq 1$  point in WHO performance status, compared with previous assessment
- 3.4. Increased symptoms of raised intracranial pressure (headache, nausea/vomiting etc.)
4. The extent to which conventionally planned RT volumes encompassed the abnormalities identified using advanced imaging
5. For the MALTING study, patients outcome will be compared to predicted outcome from the prognostic model proposed by Gorlia et al (Lancet Oncology, 2008). This uses a nomogram that involves MGMT promoter methylation status, age, performance status, extent of resection, and Mini-Mental State Examination (MMSE) to predict outcome. Using a modified Sliding Dichotomy design, the outcome for each patient is compared to their predicted outcome.

**Completion date**

31/12/2018

## Eligibility

**Key inclusion criteria**

1. Imaging appearances of a high grade glioma
2. Likely to be suitable for radiotherapy (60 Gy) with concomitant and adjuvant temozolomide
3. World Health Organization (WHO) performance status (PS) grade 0 or 2
4. Aged 18 - 75 years, either sex
5. Resection or biopsy (although only those suitable for maximal resection will be considered for the MALTING Trial)

Patients for the MALTING Trial will be felt by their consultant neurosurgeon to be suitable for 5-aminolevulinic acid (5-ALA) fluorescence-guided resection with insertion of carmustine wafers.

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

All

**Key exclusion criteria**

1. Unsuitable for a contrast-enhanced MRI (MR unsafe metallic implants, claustrophobia, allergy to gadolinium contrast agent or severe renal impairment)
2. Pregnant

3. Allergic to aminolevulinic acid
4. Suffering from porphyria. Care will be taken if the patient is taking other photosensitising drugs.

**Date of first enrolment**

01/03/2010

**Date of final enrolment**

01/03/2015

## Locations

**Countries of recruitment**

United Kingdom

England

**Study participating centre****Neurosurgery Division**

Cambridge

United Kingdom

CB2 0QQ

## Sponsor information

**Organisation**

Cambridge University Hospitals NHS Foundation Trust (UK)

**ROR**

<https://ror.org/04v54gj93>

## Funder(s)

**Funder type**

Government

**Funder Name**

National Institute for Health Research (NIHR) (UK) - Clinician Scientist Award (ref: NIHR/CS/009/011)

# Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration