ISRCTN62699180 https://doi.org/10.1186/ISRCTN62699180

Efficiency of 7.2% hypertonic saline hydroxyethyl starch 200/0.5 versus mannitol 15% in the treatment of increased intracranial pressure in neurosurgical patients

Submission date
23/06/2005

Recruitment status
No longer recruiting

Registration date
24/06/2005

Overall study status
Completed

Last Edited
15/04/2008

Condition category
Nervous System Diseases

[ ] Prospectively registered
[ ] Protocol
[ ] Statistical analysis plan
[X] Results
[ ] Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Jens Soukup

Contact details
Martin-Luther-University Halle
Department of Anaesthesia and Critical Care
Klinikum Kröllwitz
Ernst-Grube-Str. 40
Halle
Germany
06097
+49 345 5575992
jens.soukup@medizin.uni-halle.de

Additional identifiers

Study information

Scientific Title

Study objectives
Severe brain injury can lead to a brain edema with increased intracranial pressure. This fact leads to a reduced cerebral blood flow and cerebral oxygenation. These situations can extend the brain edema with a possible poor patient outcome.

7.2% hypertonic saline hydroxyethyl starch 200/0.5 is more effective compared to mannitol 15% in the treatment of increased intracranial pressure in neurosurgical patients.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Not provided at time of registration

Primary study design
Interventional

Study design
Randomised controlled trial

Study type(s)
Treatment

Health condition(s) or problem(s) studied
Increased intracranial pressure

Interventions
Cerebral perfusion pressure (CPP) directed therapy with CPP >70 mmHg, sedation, normoventilation.

If ICP >20 mmHg patients receive either 7.2% hydroxyethyl starch 200/0.5 or mannitol 15%.

Intervention Type
Drug

Phase
Not Specified

Drug/device/biological/vaccine name(s)
7.2% hypertonic saline hydroxyethyl starch 200/0.5, mannitol 15%

Primary outcome(s)
ICP <15 mmHg

Key secondary outcome(s)
Survival, discharge status

Completion date

31/08/2004

Eligibility

Key inclusion criteria
1. Neurosurgical patients >18 years with severe neuronal damage being at risk of increased intracranial pressure (ICP)
2. Cerebral edema - visualized by computed tomography (CT) scan, continuous monitoring of ICP

Participant type(s)
Patient

Healthy volunteers allowed
No

Age group
Adult

Lower age limit
18 Years

Sex
All

Key exclusion criteria
Exclusion criteria were elevated ICP due to space occupying lesions with indication for neurosurgical intervention, severe renal failure, metabolic disorders, initial serum sodium >150 mmol/l and initial serum osmolarity >320 mosm/kg

Date of first enrolment
01/02/2003

Date of final enrolment
31/08/2004

Locations

Countries of recruitment
Germany

Study participating centre
Martin-Luther-University Halle
Halle
Germany
06097

Sponsor information

Organisation
Martin-Luther-University Halle - Department of Anesthesia and Critical Care (Germany)

ROR
https://ror.org/05gqaka33

Funder(s)

Funder type
University/education

Funder Name
Self-funded trial, Department of Anesthesia and Critical Care, Martin-Luther University Halle (Germany)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Results:	05/10/2005		Yes	No