

The effectiveness of intrauterine insemination (IUI) in subfertile couples with an isolated cervical factor: a randomised controlled trial

Submission date 08/02/2006	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 08/02/2006	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 19/10/2007	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Mrs P Steures

Contact details
Academic Medical Center
Center of Reproductive Medicine
OFO-project
H4-213
P.O. Box 22660
Amsterdam
Netherlands
1100 DD
+31 (0)20 5663857
ofoproject@amc.uva.nl

Additional identifiers

Protocol serial number
2

Study information

Scientific Title

Study objectives

1. We hypothesised a beneficial effect of IUI in couples with an isolated cervical factor
2. Furthermore we hypothesised that the post-coital test can identify those couples who would benefit from IUI without ovarian hyperstimulation

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from the local medical ethics committee

Primary study design

Interventional

Study design

Multicentre, randomised, active controlled, parallel group trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Subfertility

Interventions

Couples were randomly allocated to IUI for six months or expectant management for six months. In the first three IUI cycles no controlled ovarian hyperstimulation (COH) was given. If these attempts failed subsequent IUI cycles were performed with COH.

Couples allocated to expectant management were followed until an ongoing pregnancy occurred within six months. If no pregnancy occurred, follow-up ended after this period. If a pregnancy miscarried, follow-up continued until the next pregnancy or the end of the six months period.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

The primary endpoint was ongoing pregnancy within six months. Ongoing pregnancy was defined as the presence of foetal cardiac activity at transvaginal sonography at a gestational age of at least 12 weeks.

Key secondary outcome(s)

Secondary endpoints were total number of clinical pregnancies, miscarriages and multiple pregnancies.

Completion date

01/07/2005

Eligibility

Key inclusion criteria

Couples with a cervical factor and otherwise no factors that reduced their fertility i.e. a prognosis for a treatment independent ongoing pregnancy in the next year higher than 30%. A cervical factor was diagnosed by a well-timed, non-progressive post-coital test (PCT) with normal semen parameters.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Key exclusion criteria

All other subfertile couples

Date of first enrolment

01/06/2002

Date of final enrolment

01/07/2005

Locations

Countries of recruitment

Netherlands

Study participating centre

Academic Medical Center

Amsterdam

Netherlands

1100 DD

Sponsor information

Organisation

Academic Medical Center Amsterdam (The Netherlands)

Funder(s)

Funder type

Research organisation

Funder Name

The Netherlands Organisation for Health Research and Development (ZonMw) (The Netherlands)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Results	01/12/2007		Yes	No