

A comparison of intravenous salbutamol and aminophylline in the management of acute severe asthma in children

Submission date 23/01/2004	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 23/01/2004	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 22/02/2008	Condition category Respiratory	☐ Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Alex Habel

Contact details
West Middlesex Hospital
Isleworth
United Kingdom
TW7 6AF

Additional identifiers

Protocol serial number
RDC01664

Study information

Scientific Title

Study objectives

To determine whether a bolus of intravenous salbutamol or an aminophylline infusion is the most effective therapy for children with acute severe asthma.

Ethics approval required

Old ethics approval format

Ethics approval(s)

The study was approved by the Thames Multicentre Research ethics committee and the local ethics committees.

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Asthma

Interventions

1. Bolus of intravenous salbutamol
2. Aminophylline infusion

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Salbutamol, aminophylline

Primary outcome(s)

Duration of the need for supplementary oxygen to maintain saturations of above 91%.

Key secondary outcome(s))

Other end points:

1. Asthma severity score two hours after starting intravenous therapy
2. Time to stopping intravenous therapy
3. Time to decision to discharge subject

Completion date

10/01/2002

Eligibility

Key inclusion criteria

1. Age 1 to 15 years
2. Acute severe asthma with an asthma severity score of greater than 7
3. Poor response to 3 "back to back" salbutamol/ipratropium nebulisers with an improvement in asthma severity score of less than 2

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Lower age limit

1 years

Upper age limit

15 years

Sex

All

Key exclusion criteria

1. A life threatening exacerbation
2. An underlying respiratory disease other than asthma
3. Cardiac disease
4. Treatment with a medication that alters the metabolism of aminophylline

Date of first enrolment

10/01/2000

Date of final enrolment

10/01/2002

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

West Middlesex Hospital

Isleworth

United Kingdom

TW7 6AF

Sponsor information

Organisation

NHS R&D Regional Programme Register - Department of Health (UK)

Funder(s)

Funder type

Government

Funder Name

NHS Executive London (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Results	01/04/2003		Yes	No