

Treating IBS with an Intestinal Microbiota Product for Health - TriuMPH

Submission date 05/12/2023	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 10/04/2025	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 14/05/2026	Condition category Digestive System	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

The study aims to assess the safety and efficacy of EBX-102-02 in patients with constipation-predominant Irritable bowel syndrome (IBS-C) and diarrhea-predominant irritable bowel syndrome (IBS-D). IBS is a common condition that affects the digestive system. It causes symptoms like stomach cramps, bloating, diarrhoea and constipation. These tend to come and go over time and can last for days, weeks or months at a time. It's usually a lifelong problem. It can be very frustrating to live with and can have a big impact on everyday life. The exact cause is unknown. Current treatments focus on relieving the symptoms. There is increased interest in the microbiota of IBS and EBX-102 has the potential to improve microbial diversity. EBX-102 capsules contain communities of dried, intestinal microorganisms taken from carefully screened pooled human stool samples.

Who can participate?

Patients aged ≥ 18 years up to 70 years old with a clinical diagnosis of IBS-C

What does the study involve?

The study will be conducted in 2 centres - The functional Gut Clinics, based in Manchester and London. 120 participants randomised in a 2:1 ratio (60 participants for the IBS-C and 60 for the IBS-D cohort with approximately 40 in the active and 20 in the placebo group for each cohort, respectively). Once eligibility criteria are confirmed and written informed consent has been obtained, participants will undergo at least a 2-week screening period to confirm the diagnosis of IBS-C. Once randomisation occurs, 8 capsules of the study drug will be administered on day 1 and day 8. Participants will then be followed up for 6 weeks (after the last dose of the study drug). The total period of study for an individual participant will be approximately 9 weeks (2 weeks screening, 1 week treatment, 6 weeks follow-up). The following assessment information will be performed/collected - medical history, physical exam, vital signs, ECG, blood sampling, stool collection, completion of questionnaires, patient diary, concomitant medication, demographics and breath tests.

What are the possible benefits and risks of participating?

There are risks, discomforts, and inconveniences associated with any research study. It is possible that these general risks could be increased by the addition of test medications. Some of the general risks may be potentially life-threatening and may not have been previously reported.

Study Assessment Risks:

Some of these procedures take place more often than they would if patients were not taking part in this study.

Blood Collection:

Taking blood samples may cause bruising and discomfort and a risk of infection or blood clots at the site of the blood collection.

Study Treatment:

Will be given in a clinic with emergency equipment and staff who are trained to monitor for and respond to any potential medical emergencies.

Risks Associated with EBX-102-02 :

Stomach pain and other gastro-type symptoms after taking the capsules. The known side effects of this type of treatment typically include bloating, diarrhoea and stomach cramps. These symptoms do not occur in everybody who takes them. Participants are encouraged to inform their study team about any side effects that they experience. The study doctor may provide additional medications to ease the experience of side effects; however, typically, symptoms resolve within a few days.

Risk of exposure to pathogens contained in the IMP due to the nature of intestinal microbial ecosystems. The sponsor operates a robust biosafety programme focussed on ensuring that starting material is appropriately and extensively screened, rendering it safe for its intended use. The sponsor works with a small number of accredited and audited laboratory partners as well as utilising in-house pathogen screening capabilities that include state-of-the-art technology. Regular monitoring of safety data for indications that pathogens have been transmitted.

Unknown Risks:

Side effects of EBX-102-02, which are unknown at this time, may occur during the study. Any new information that may affect participants' health or which may make the participants want to stop taking part in the study will be shared with them as soon as it becomes available.

Pregnancy Prevention:

There may be a risk in exposing an unborn child to study drugs, and all risks are not known at this time. Women must take precautions to avoid exposing an unborn child to study drugs, as described in the PIS-ICF.

Patients will be informed of all of the above risks in the Patient Information Sheet and will be asked to notify their study doctor or study staff should they experience any side effects during the study. Patients will be monitored throughout the study to minimise risks.

Where is the study run from?

EnteroBiotix Ltd

When is the study starting and how long is it expected to run for?

December 2023 to December 2024

Who is funding the study?
EnteroBiotix Ltd

Who is the main contact?
Shinofa Rizan, clinops@enterobiotix.com

Contact information

Type(s)
Scientific

Contact name
Dr James McIlroy

ORCID ID
<https://orcid.org/0000-0002-8265-1418>

Contact details
EnteroBiotix Limited, Phoenix House, Phoenix Crescent, Strathclyde Business Park
Bellshill, Lanarkshire
United Kingdom
ML4 3NJ
+44 (0)1698 501370
j.mcilroy@enterobiotix.com

Type(s)
Principal investigator

Contact name
Dr Paul Goldsmith

Contact details
262 Deansgate
Manchester
United Kingdom
M3 4BG
+44 (0)7787 113 150
paul.goldsmith@mft.nhs.uk

Type(s)
Public

Contact name
Ms Shinofa Rizan

Contact details
EnteroBiotix Limited
Phoenix House, Phoenix Crescent,
Strathclyde Business Park
Bellshill, Lanarkshire

United Kingdom
ML4 3NJ
-
clinops@enterobiotix.com

Additional identifiers

Integrated Research Application System (IRAS)
1009182

Protocol serial number
EBX-102-02-201

Study information

Scientific Title

A randomised, double-blind, placebo-controlled, phase II trial assessing the safety and efficacy of EBX-102-02 in patients with Irritable Bowel Syndrome (IBS)

Acronym

TrluMPH

Study objectives

Current study objectives as of 14/05/2026:

The primary objective of the study is to assess the safety and tolerability of EBX 102-02 in irritable bowel syndrome participants

The main secondary objective is to assess the effects of EBX-102-02 on IBS symptomology including pain, stool consistency and frequency and constipation.

Previous study objectives:

To assess the safety and tolerability of EBX 102-02 in irritable bowel syndrome participants

To assess the effects of EBX-102-02 on:

IBS-C symptomatology including:

Pain

Bloating

Stool consistency and frequency

Biomarker breath test measurements

Intestinal microbiome composition

Quality of life/global well-being

Anxiety and depression

Stool based short chain fatty acids (SCFAs) and bile acids (BAs)

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 23/01/2024, West Midlands - Edgbaston Research Ethics Committee (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 (0)20 7104 8155; edgbaston.rec@hra.nhs.uk), ref: 23/WM/0274

Study design

Randomized double-blind placebo-controlled phase II trial

Primary study design

Interventional

Study type(s)

Efficacy, Safety

Health condition(s) or problem(s) studied

Irritable bowel syndrome (IBS)

Interventions

EBX-102-02 is a gastro-resistant hard hydroxypropyl methylcellulose (HPMC) capsule containing communities of dried, intestinal microorganisms derived from rigorously screened pooled human stool samples. Visual and weight-matched placebo capsules are available to maintain the blind. Participants will be randomly assigned in a 2:1 ratio to receive active or placebo IMP via an Interactive Web-based Randomisation System.

Participants will receive 8 capsules of IMP to be taken at each of 2 separate administrations 1 week apart. Administrations will be directly observed in the clinic. Participants will then be followed up for 6 weeks (after the last dose of IMP).

Intervention Type

Drug

Phase

Phase II

Drug/device/biological/vaccine name(s)

EBX-102-02 [EBX-102]

Primary outcome(s)

Current primary outcomes as of 14/05/2026:

Incidence and type of adverse events (AEs) measured using data collected during safety monitoring assessments at baseline (dosing 1) and weeks 1 (dosing 2), 3 (follow-up 1) and 7 (final follow-up), clinically significant abnormal vital signs at baseline and weeks 1, 3 and 7, and clinically significant abnormal ECG and clinical laboratory assessments at week -1 (pre-treatment) and week 7

Previous primary outcomes:

Incidence of adverse events (AEs) and safety data (including vital signs, physical examinations and laboratory test results) up to and including 6 weeks post-treatment

Key secondary outcome(s)

Current secondary outcomes as of 14/05/2026:

1. Irritable bowel symptom severity measured using the Irritable Bowel Syndrome Symptom Severity Scale (IBS-SSS) questionnaire at baseline and weeks 1, 3 and 7
2. Change in symptoms of constipation measured using Patient Assessment of Constipation Symptoms (PAC-SYM) questionnaire at baseline and weeks 1, 3 and 7
3. Change in stool consistency measured using the Bristol Stool Scale at baseline and weeks 1, 3 and 7
4. Change in stool frequency and other symptomatology (i.e. severity of pain) at baseline and weeks 1, 3 and 7

Previous secondary outcomes:

The following secondary outcome measures will be assessed up to and including 6 weeks post-treatment:

1. Change in IBS-C symptomatology measured on the IBS Symptom Severity Score (IBS-SSS).
2. Change in responses to Patient Assessment of Constipation Symptoms (PAC-SYM) questionnaire
3. Change in stool consistency measured on the Bristol Stool Scale
4. Change in stool frequency and other symptomatology measured by a bowel habit diary
5. Change in fermentation profiles, total gas production, and small intestinal bacterial overgrowth in biomarker breath test measurements
6. Taxonomic microbiome analysis
7. Change in Quality of Life scores measured on the IBS Quality of Life questionnaire (IBS-QOL)
8. Change in Hospital Anxiety and Depression score
9. Requirement for rescue medication
10. Metabolite analysis: SCFAs and BAs in stool

Completion date

23/12/2024

Eligibility

Key inclusion criteria

1. Willing and able to provide informed consent
 2. Male or female aged ≥ 18 years up to 70 years
 3. A clinical diagnosis of IBS-C or IBS-D, as confirmed by Rome IV grading criteria, excluding patients with mild disease by using an IBS-SSS inclusion of ≥ 175
 4. Willing to discontinue all medications for bowel habit abnormalities after providing consent
 5. Willing to abstain from consuming regular 'over-the-counter' pre- or probiotics from pharmacies or other retailers from screening through to end of follow-up
 6. If women of childbearing potential (WOCBP), subjects must have a negative serum pregnancy test at screening, a negative urine pregnancy test at randomisation and must be willing to use a highly effective method of birth control for the duration of the study. Acceptable methods of contraception:
 - 6.1. Hormonal contraception associated with inhibition of ovulation
 - 6.2. Intrauterine device (IUD)
 - 6.3. Intrauterine hormone-releasing system (IUS)
 - 6.4. Bilateral tubal occlusion
 - 6.5. Vasectomised partner
 - 6.6. Sexual abstinence, in line with the preferred and usual lifestyle of the subject
- Note: Periodic abstinence (such as calendar, ovulation, symptothermal, post-ovulation methods) and withdrawal, and condom with spermicide, are not considered a highly effective method of

contraception for female subjects of childbearing potential.

7. If male, subjects must be prepared to use reliable barrier method contraception and a second method such as spermicide for the duration of the study unless surgically sterile

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

18 years

Upper age limit

70 years

Sex

All

Total final enrolment

122

Key exclusion criteria

1. Women who are pregnant or breastfeeding
2. Planned surgery requiring general anaesthetic during the course of the study
3. Participants who are planning to significantly change their diet (e.g. weight loss programme, becoming vegetarian) during the study period. Patients established on a low fermentable oligosaccharides, disaccharides, monosaccharides and polyols (FODMAP) diet can continue without changes to it
4. Confirmed diagnosis of mixed type IBS (IBS-M), or unclassified IBS (IBS-U)
5. In IBS-C subjects: diarrhoea within 7 days prior to screening
 - 5.1. If possible infective diarrhoea, then wait 3 weeks before starting bowel habit diary
6. In IBS-D subjects
 - 6.1. Nocturnal diarrhoea
 - 6.2. BSS type 7 on more than 5 days per week
 - 6.3. Diarrhoea associated with foreign travel in the 4 weeks prior to screening
7. Other chronic gastrointestinal (GI) disease including:
 - 7.1. Inflammatory bowel disease
 - 7.2. Diverticulitis
 - 7.3. Malabsorption syndromes e.g. lactose intolerance (with proven lactase deficiency)
8. Any history of malignant tumours (primary or secondary) affecting any part of the GI tract
9. History of colectomy/ileostomy at any time
10. History of colonic perforation or fistula
11. History of any malignancy within the 5 years prior to screening, excluding non-melanoma skin cancers
12. Conditions associated with increased risk of GI cancer, e.g. familial adenomatous polyposis coli
13. Abdominal surgical intervention except for appendectomy, hernia repair, and gynaecological

and urological procedures

14. Known lactulose intolerance

15. Ongoing requirement for medications known to cause constipation e.g. Iron supplementation, opiates or diarrhoea e.g. antacids, non-steroidal anti-inflammatory drugs (NSAIDs)

16. Use of any prohibited medications for which a participant cannot complete the appropriate washout period

17. Aspartate aminotransferase (AST) or alanine aminotransferase (ALT) $\geq 2.5x$ upper limit of normal (ULN)

18. History of human immunodeficiency virus (HIV), hepatitis B virus (HBV) or hepatitis C virus (HCV) infection, regardless of current viral status and test results

19. Use of systemic antibiotics within 7 days prior to screening, or intended use during the study

20. Faecal microbiota transplantation (FMT) within the past 12 months

21. History of sensitivity to any of the study drug components, or a history of drug allergy that in the opinion of the Investigator contraindicates study participation

22. Participants with dysphagia, or inability to ingest capsules (e.g. severe nausea, vomiting, delayed gastric emptying) or history of 'choking' on capsules

23. Have taken an IMP within the last 3 months

24. Planned or active participation in any other study with an IMP

25. Any autoimmune or oncologic disease requiring, or that may require, systemic treatment with steroids and/or other immunosuppressants/immunomodulators

26. Significant bleeding disorder

27. Anaphylactic food allergy

28. Requirement for vasopressors

29. Valvular heart disease or known structural defects of the heart

30. Clinically significant medical or surgical history or any condition that could interfere with study participation or confound the assessments in the opinion of the study Investigator

Date of first enrolment

11/03/2024

Date of final enrolment

11/11/2024

Locations

Countries of recruitment

United Kingdom

Study participating centre

The Functional Gut Clinic (Manchester)

262 Deansgate

Manchester

England

M3 4BG

Study participating centre

The Functional Gut Clinic
8 Dorset Square
London
England
NW1 6PU

Sponsor information

Organisation
EnteroBiotix Limited

Funder(s)

Funder type
Industry

Funder Name
EnteroBiotix Ltd

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not expected to be made available