

A multicentre randomised controlled trial of amnioinfusion

Submission date 16/11/2005	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 16/11/2005	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 15/11/2013	Condition category Neonatal Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr William Donald Fraser

Contact details
Hôpital Sainte-Justine
Département d'Obstétrique-Gynécologie
3175, Chemin de la Côte-Ste-Catherine
4e Étage, Bloc 9, Local 4986-B
Montréal
Canada
H3T 1C5
+1 (0) 514 345 4931 (4155)
william.fraser@umontreal.ca

Additional identifiers

Protocol serial number
MCT-15221

Study information

Scientific Title

A multicentre randomised controlled trial of amnioinfusion for thickly meconium stained amniotic fluid to reduce the risk of moderate to severe meconium aspiration syndrome or perinatal death

Study objectives

1. To assess when compared to standard care, a policy of amnioinfusion for thickly meconium stained amniotic fluid reduces the risk of either moderate to severe meconium aspiration syndrome or perinatal death
2. To assess the effects of amnioinfusion on the risk of occurrence of other indicators of neonatal morbidity, cesarean section and indicators of severe maternal morbidity
3. Will evaluate the acceptability of amnioinfusion to women and determine the cost-effectiveness of the procedure

Ethics approval required

Old ethics approval format

Ethics approval(s)

University Hospital of Québec Research Ethics Committee approved on the 26th June 2002.

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Meconium aspiration syndrome

Interventions

Experimental: Intrapartum transcervical infusion of 800 ml of sterile normal saline over a period of 40 minutes at 20 ml/min followed by continuous infusion up to total volume of 1500 ml
Control: usual care

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Occurrence of perinatal death or moderate/severe meconium aspiration syndrome.

Key secondary outcome(s)

Maternal:

1. Haemorrhage requiring blood transfusion
2. Hysterectomy
3. Uterine rupture

4. Febrile morbidity
5. Caesarean section
6. Acceptability of amnioinfusion to women
7. Determine the cost-effectiveness of the procedure

Foetal:

1. Foetal heart rate tracing abnormalities
2. Positive blood or cerebrospinal fluid culture
3. Convulsions
4. Need for tube feeding
5. Fractures and palsies
6. Duration of oxygen supplementation
7. Duration of ventilation
8. Abnormal chest x-ray

Completion date

30/04/2002

Eligibility

Key inclusion criteria

1. Pregnant women of childbearing age, with a single baby in cephalic presentation
2. Ruptured membranes (spontaneous or artificial)
3. Gestational age greater than or equal to 36 weeks
4. Established labour defined as the presence of regular contractions occurring at 5 minute intervals
5. Cervical dilation between 2 and 7 cm inclusively at randomisation
6. Thick meconium
7. Foetal status considered as acceptable after a 30-minute period of electronic foetal heart rate monitoring

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Key exclusion criteria

1. Cord prolapse
2. Known or suspected major foetal anomaly
3. Suspicion of chorioamnionitis on the basis of maternal fever or abnormal vaginal discharge
4. Known placenta praevia or vaginal bleeding due to other causes
5. Known intravenous drug user
6. Patient known or suspected to be at high risk of human immunodeficiency virus (HIV)
7. Hepatitis B or C

8. Active genital herpetic lesions
9. Uterine over distension
10. Previous uterine incision other than low transverse
11. Recurrent late decelerations
12. Prolonged decelerations
13. Scalp blood pH less than 7.15
14. Any contradictions to labour
15. Unable to comprehend the consent form
16. Use of narcotic analgesics prior to consent

Date of first enrolment

01/04/1999

Date of final enrolment

30/04/2002

Locations

Countries of recruitment

United Kingdom

Argentina

Belgium

Brazil

Canada

France

Mexico

Portugal

South Africa

Switzerland

Tunisia

United States of America

Uruguay

Study participating centre

Hôpital Sainte-Justine

Montréal

Canada

H3T 1C5

Sponsor information

Organisation

Hospital Sainte-Justine, Montréal (Canada)

ROR

<https://ror.org/01gv74p78>

Funder(s)

Funder type

Research organisation

Funder Name

Canadian Institutes of Health Research (CIHR) (Canada) - <http://www.cihr-irsc.gc.ca> (ref: MCT-15221)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/09/2005		Yes	No