

Randomised phase II trial comparing vinorelbine and capecitabine with docetaxel and capecitabine chemotherapy for metastatic breast cancer

Submission date 12/09/2003	Recruitment status Stopped	☐ Prospectively registered ☐ Protocol
Registration date 12/09/2003	Overall study status Stopped	☐ Statistical analysis plan ☐ Results
Last Edited 29/10/2012	Condition category Cancer	☐ Individual participant data ☐ Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Chris Gallagher

Contact details
Consultant
Medical Oncology Department
St Bartholomew's Hospital
West Smithfield
London
United Kingdom
EC1A 7BE
+44 (0)20 7601 8521
chris.gallagher@bartsandthelondon.nhs.uk

Additional identifiers

Protocol serial number
N0205108870

Study information

Scientific Title

Study objectives

Toxicity, response and survival.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Breast cancer

Interventions

Randomisation will be stratified by previous anthracycline exposure.

1. Vinorelbine 25mg/m² intravenous (IV) 30 min DI + 8q 21 days. Capecitabine 1 g/m² orally, twice daily for 14 in every 21 days.
2. Docetaxel 75 mg/m² IV 60 min every 21 days with dexamethasone 8 mg twice daily for three days starting 24 hours before each docetaxel treatment. Capecitabine 1 g/m² orally, twice daily for 14 in every 21 days.

Added 17 July 2008: the trial closed in 2006 due to poor recruitment.

Intervention Type

Drug

Phase

Phase II

Drug/device/biological/vaccine name(s)

Vinorelbine and capecitabine

Primary outcome(s)

Toxicity, response and survival.

Key secondary outcome(s)

Not provided at time of registration

Completion date

25/11/2005

Reason abandoned (if study stopped)

Poor recruitment

Eligibility

Key inclusion criteria

Metastatic breast cancer following initially histologically proven adenocarcinoma of the breast.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Female

Key exclusion criteria

Exclusions include previous treatment with vinorelbine, docetaxel or capecitabine and presence of other primary cancers.

Date of first enrolment

01/04/2002

Date of final enrolment

25/11/2005

Locations

Countries of recruitment

United Kingdom

England

Study participating centre**Consultant**

London

United Kingdom

EC1A 7BE

Sponsor information

Organisation

Department of Health (UK)

Funder(s)

Funder type

Government

Funder Name

Barts and The London NHS Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration