

# Supervised pelvic floor muscle training improves sexual function and diminishes sexual distress in women with multiple sclerosis

<b>Submission date</b> 03/07/2023	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 06/07/2023	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 03/09/2024	<b>Condition category</b> Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Pelvic floor muscle training (PFMT) contributes to better sexual function, as muscle strength and ability to properly contract to improve vaginal receptivity and responsiveness, orgasm and sexual pleasure. Multiple sclerosis (MS) is the most common chronic nervous system disorder leading to sexual dysfunction, with a prevalence of 40-70%. More than a third of MS patients experience signs of pelvic floor weakness. The primary outcome of this study is to evaluate the effect of PFMT on improving sexual function and sexual distress in women suffering from MS.

### Who can participate?

Adult women aged 18-45 years old with MS

### What does the study involve?

In this study, women suffering from MS will be included in an intervention Group A (12 weeks of PFMT), and in a control Group B (observation group; negative control group). All women will be assessed with the Female Sexual Function Index (FSFI) and Female Sexual Distress Scale-Revised (FSDS-R) at the study's beginning and end.

### What are the possible benefits and risks of participating?

Possible benefits include better sexual function. No risks are expected from the pelvic floor exercises.

### Where is the study run from?

Ey Prattein Rehabilitation Centre (Greece)

### When is the study starting and how long is it expected to run for?

May 2021 to May 2023

### Who is funding the study?

Investigator initiated and funded

Who is the main contact?

Athanasios Zachariou (Assistant Prof Urology), zahariou@otenet.gr (Greece)

## Contact information

### Type(s)

Principal investigator

### Contact name

Prof Athanasios Zachariou

### ORCID ID

<https://orcid.org/0000-0002-5287-4450>

### Contact details

3 Spyridi Street

Volos

Greece

38221

+306937227778

zahariou@otenet.gr

## Additional identifiers

## Study information

### Scientific Title

Supervised pelvic floor muscle training improves sexual function and diminishes sexual distress in women with multiple sclerosis: a randomized controlled study

### Study objectives

Pelvic floor muscle training (PFMT) contributes to better sexual function, as muscle strength and ability to properly contract improve vaginal receptivity and responsiveness, orgasm and sexual pleasure. Multiple sclerosis (MS) is the most common chronic nervous system disorder leading to sexual dysfunction, with a prevalence of 40-70%. More than a third of MS patients experience signs of pelvic floor weakness. The primary outcome of this study is to evaluate the effect of PFMT on improving sexual function and sexual distress in women suffering from MS.

### Ethics approval required

Ethics approval required

### Ethics approval(s)

approved 01/09/2021, KENTAVROS Center for Physical Medicine and Rehabilitation (Apollonos 94, Volos, 38222, Greece; +30 2421043000; info@kentavros.com.gr), ref: 12/2021

### Study design

Interventional randomized controlled trial

### Primary study design

Interventional

## Study type(s)

Quality of life

## Health condition(s) or problem(s) studied

Supervised pelvic floor muscle training improves sexual function and diminishes sexual distress in women with multiple sclerosis

## Interventions

The intervention included women suffering from MS with an EDSS score <4 and divided them into two groups.

In Group A, women followed a program of supervised pelvic floor muscle training (PFMT) for 12 weeks, and women in Group B comprised the control group. The total duration of the intervention was 12 weeks.

At the beginning and end of the study, all women completed the FSFI and FSDS-R questionnaires to assess their sexual function/distress and evaluate if there were any differences between the two groups.

1. Method of randomization. Women were consecutively randomized into the two groups.

Consecutive sampling is a non-probability sampling technique where samples are picked at the ease of the investigator, more like convenient sampling. In that way, we produced two groups with an equal number of women.

2. The principal investigator, doctor, and physiotherapist evaluated the pelvic floor to provide evidence that the woman could perform pelvic floor muscle exercises. All work in the rehabilitation center and are members of the pelvic floor unit. They already have the necessary specific training and are all considered specialists in their topic.

3. The instruction of the PFMT was face-to-face, but during the study period, there were a lot of telephone communications to strengthen our patients and increase adherence to the program.

4. All the necessary interventions occurred on the premises of Kentavros Rehabilitation Center.

## Intervention Type

Behavioural

## Primary outcome(s)

Effect of pelvic floor muscle training (PFMT) on improving sexual function and sexual distress in women suffering from MS measured using the Female Sexual Function Index (FSFI) and the Female Sexual Distress Scale-Revised (FSDS-R) questionnaires scores at the beginning and end of the twelve-week study

## Key secondary outcome(s)

There are no secondary outcome measures

## Completion date

01/05/2023

## Eligibility

### Key inclusion criteria

1. 18 years of age or older
2. Diagnosis of relapsing-remitting MS
3. Stable condition for a minimum period of six months
4. Kurtzke's Expanded Disability Status Scale (EDSS) score < 4
5. Women sexually active for at least four weeks
6. Cognitive ability to complete the questionnaires and study protocol
7. Ability to contract PFM evaluated by the primary investigator

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Upper age limit**

45 years

**Sex**

Female

**Total final enrolment**

84

**Key exclusion criteria**

1. Previous pelvic floor muscle training program
2. Ongoing pregnancy
3. Child delivery within the previous six months
4. Urinary or faecal incontinence
5. Pelvic organ prolapses greater than stage I
6. Perimenopause or menopause period

**Date of first enrolment**

01/09/2021

**Date of final enrolment**

01/02/2023

**Locations****Countries of recruitment**

Greece

**Study participating centre**  
**Ey Prattein Rehabilitation Centre**  
94 Apollonos Street  
Volos  
Greece  
38222

## Sponsor information

**Organisation**  
KENTAVROS

## Funder(s)

**Funder type**  
Hospital/treatment centre

**Funder Name**  
KENTAVROS Center for Physical Medicine and Rehabilitation

## Results and Publications

### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Athanasios Zachariou (Assistant Prof Urology), zahariou@otenet.gr.

### IPD sharing plan summary

Available on request

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		12/01/2024	03/09/2024	Yes	No
<a href="#">Participant information sheet</a>	Patient consent form		06/07/2023	No	Yes