

A comparison of once- and twice-weekly eccentric training on muscular health of older adults

Submission date 19/01/2024	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 24/01/2024	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 02/05/2024	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input checked="" type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Currently, the Chief Medical Officers in the United Kingdom recommend that older adults partake in strength or balance training twice per week to improve muscular function. However, adherence to these recommendations is low with time and physical capacity being reported as two barriers to meeting these guidelines. Eccentric resistance training (lengthening of the muscle whilst contracting) results in greater muscular adaptations than traditional resistance and requires a lower metabolic demand, therefore fewer weekly sessions may be required and they should be easier to tolerate, even for those with pre-existing comorbidities. Therefore, this study aimed to compare the effects of once-weekly eccentric resistance training to twice-weekly resistance training on muscular structure and function in healthy older adults.

Who can participate?

Community-dwelling older adults (aged 60 years and over) who can ambulate independently, do not suffer from any musculoskeletal or neuromuscular diseases, and are not taking any medication that may affect muscular function or balance

What does the study involve?

Participants are randomly allocated to one of three groups. The non-active control group maintained normal living conditions whereas the two training groups performed multi-joint isokinetic eccentric exercise for 12 weeks at 50% of their maximum eccentric strength. The once-weekly training group trained once per week and the twice-weekly training group trained twice per week; training volume was not matched between groups. The training lasted for 7 minutes in week 1 and progressed to 12 minutes in week 4, which remained the same hereafter until the training programme was completed in week 12. Participants were assessed at baseline, mid-training (week 7) and post-training (week 13).

What are the possible benefits and risks of participating?

Possible benefits are improvements in muscle strength, power, and size, whereas possible risks are muscle fatigue and temporary exercise-induced muscle damage.

Where is the study run from?
University of Northampton Health and Performance Laboratory (UK)

When is the study starting and how long is it expected to run for?
November 2018 to March 2021

Who is funding the study?
Wellcome Trust (UK)

Who is the main contact?
Mr Brett Baxter, brett.baxter@northampton.ac.uk

Contact information

Type(s)

Public, Scientific, Principal investigator

Contact name

Mr Brett Baxter

ORCID ID

<https://orcid.org/0000-0002-7479-3985>

Contact details

Waterside Campus
University Drive
Northampton
United Kingdom
NN1 5PH
+44 (0)1604 892293
brett.baxter@northampton.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Protocol serial number

IRAS 339894

Study information

Scientific Title

Effects of once- versus twice-weekly eccentric resistance training on muscle structure and function in older adults

Acronym

G1XG2XECC

Study objectives

1. Both eccentric resistance training groups (once- and twice-weekly) would alter muscle structure and function
2. The twice-weekly training group would alter muscle structure and function to a greater extent than the once-weekly group

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 11/12/2018, University of Northampton Research Ethics Committee (Faculty of Art, Science, and Technology, Waterside Campus, University Drive, Northampton, NN1 5PH, United Kingdom; +44 (0)1604892523; Merryn.Ekberg@Northampton.ac.uk), ref: ETH1819-0053

Study design

Parallel randomized controlled trial

Primary study design

Interventional

Study type(s)

Efficacy

Health condition(s) or problem(s) studied

Improving neuromuscular structure and function in healthy older adults

Interventions

Parallel randomisation was performed using a random number generator online. The non-active control group maintained normal living conditions whereas the two training groups performed multi-joint isokinetic eccentric exercise for 12 weeks at 50% of their maximum eccentric strength. The once-weekly training group trained once per week and the twice-weekly training group trained twice per week; training volume was not matched between groups. The training lasted for 7 minutes in week 1 and progressed to 12 minutes in week 4, which remained the same hereafter until the training programme was completed in week 12. Participants were assessed at baseline, mid-training (week 7) and post-training (week 13).

Intervention Type

Other

Primary outcome(s)

Lower-limb muscular strength measured via dynamometry at baseline, mid-training (week 7) and post-training (week 13)

Key secondary outcome(s)

1. Lower-limb muscular power measured using the 10-repetition sit-to-stand test at baseline, mid-training (week 7) and post-training (week 13)
2. Vastus lateralis muscle thickness measured using B-mode ultrasonography at baseline, mid-training (week 7) and post-training (week 13)

Completion date

26/03/2021

Eligibility

Key inclusion criteria

1. ≥ 60 years of age
2. Able to independently ambulate without walking aids
3. Free from any illnesses and/or medication that affected the neuromuscular system or balance
4. Not currently involved in a structured exercise programme

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Senior

Lower age limit

60 years

Upper age limit

100 years

Sex

All

Total final enrolment

42

Key exclusion criteria

1. < 60 years of age
2. Enrolled in a resistance training programme
3. Diagnosed with a neuromuscular or musculoskeletal disease that affects balance and/or strength
4. On medication that affects neuromuscular or musculoskeletal health

Date of first enrolment

29/04/2019

Date of final enrolment

31/05/2019

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
University of Northampton
University Drive
Northampton
United Kingdom
NN1 5PH

Sponsor information

Organisation
University of Northampton

ROR
<https://ror.org/04jp2hx10>

Funder(s)

Funder type
Charity

Funder Name
Wellcome Trust

Alternative Name(s)
Wellcome, WT

Funding Body Type
Private sector organisation

Funding Body Subtype
Trusts, charities, foundations (both public and private)

Location
United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are stored in a publicly available repository

The name of the repository: PURE

A persistent weblink to the dataset: <https://doi.org/10.24339/3373b688-e811-4847-9f38-ccf09a9c843a>

The type of data stored: objective discrete data

The process for requesting access (if non-publicly available): N/A

Dates of availability: N/A

Whether consent from participants was required and obtained: yes

Comments on data anonymization: data are anonymised using participant ID numbers

Any ethical or legal restrictions: no

IPD sharing plan summary

Stored in publicly available repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		26/04/2024	02/05/2024	Yes	No
Dataset		16/10/2023	19/01/2024	No	No
Participant information sheet			19/01/2024	No	Yes