

# Dose-intensified rechallenge with temozolomide, one week on one week off versus three weeks on one week off in patients with progressive or recurrent glioblastoma

<b>Submission date</b> 23/02/2009	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 23/04/2009	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 07/02/2019	<b>Condition category</b> Cancer	<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Prof Wolfgang Wick

**Contact details**  
University Hospital Heidelberg  
Karl-Ruprechts-University Heidelberg  
Vossstr. 2/ Geb. 4410  
Heidelberg  
Germany  
69120

## Additional identifiers

**ClinicalTrials.gov (NCT)**  
NCT00941460

**Clinical Trials Information System (CTIS)**  
2008-006871-60

# Study information

## Scientific Title

Dose-intensified rechallenge with temozolomide, one week on one week off versus three weeks on one week off in patients with progressive or recurrent glioblastoma: a prospective, multicentre, multinational, randomised, parallel-group, open, phase II trial

## Acronym

Director

## Study objectives

The primary objective of this study is to show the superiority of arm A (one week on temozolomide one week off) versus arm B (three weeks on temozolomide one week off) in terms of time to treatment failure.

On 05/01/2010 this record was updated to include further information on the site locations; this information can be found in the interventions section of this record under the above update date. At this time, the anticipated start date of this trial was also updated to the date of first participant recruitment. The initial anticipated start date of this trial was 16/03/2009.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

The Ethics Committee of the Medical Faculty of Heidelberg (Ethikkommission der Medizinischen Fakultät Heidelberg), 18/06/2009, ref: AFmu-050/2009

## Study design

Prospective multicentre multinational randomised parallel-group open phase II trial

## Primary study design

Interventional

## Study type(s)

Treatment

## Health condition(s) or problem(s) studied

Progressive or recurrent glioblastoma

## Interventions

In treatment arm A, patients will be treated with an initial dose of 120 mg/m<sup>2</sup> unless there was grade III or IV myelotoxicity with conventional temozolomide (5/28) previously. These patients will be started at 90 mg/m<sup>2</sup>. Temozolomide will be given orally on days 1 - 7 and 15 - 21. The dose will be modified if necessary.

In treatment arm B, patients will start with an initial dose of 80 mg/m<sup>2</sup> unless there was significant myelotoxicity with conventional temozolomide (5/28) previously. These patients will be started at 60 mg/m<sup>2</sup>. Temozolomide will be given orally on days 1 - 21. The dose will be modified if necessary.

1 year of treatment; follow-up until death or for one year.

Site location and principal investigator information added as of 05/01/2010:

University Hospital Zurich (Switzerland)

Department of Neurology

Zurich, Switzerland

Site PI: Prof Dr M Weller

Email: michael.weller@usz.ch

University Hospital Complex of Vaud (Centre Hospitalier Universitaire Vaudois [CHUV])  
(Switzerland)

University of Lausanne

Lausanne, Switzerland

Site PI: Prof Dr R Stupp

Email: roger.stupp@chuv.hospvd.ch

Charité - University Medicine Berlin (Charité - Universitätsmedizin Berlin) (Germany)

Department of Neurosurgery

Berlin, Germany

Site PI: Prof Dr P Vajkoczy

Email: peter.vajkoczy@charite.de

University Hospital Bonn (Universitätsklinikum Bonn) (Germany)

Department of Neurology

Bonn, Germany

Site PI: Prof Dr U Herrlinger

Email: ulrich.herrlinger@ukb.uni-bonn.de

The Miners' Hospital Bochum-Langendreer (Knappschaftskrankenhaus Bochum-Langendreer)  
(Germany)

Department of Neurology

Bochum, Germany

Site PI: Prof Dr U Schlegel

Email: uwe.schlegel@kk-bochum.de

Johann Wolfgang Goethe University Hospital (Klinikum der Johann-Wolfgang von Goethe-  
Universität) (Germany)

Dr Senckenbergisches Institut für Neuroonkologie

Zentrum für Neurologie und Neurochirurgie

Frankfurt am Main, Germany

Site PI: Prof Dr J Steinbach

Email: joachim.steinbach@med.uni-frankfurt.de

Saarland University Hospital (Universitätsklinikum des Saarlandes) (Germany)

Department of Neurosurgery

Homburg/Saar, Germany

Site PI: PD Dr R Ketter Homburg

Email: ncrket@uks.eu

University Hospital Heidelberg (Universitätsklinikum Heidelberg) (Germany)

Department of Neuro-oncology

Heidelberg, Germany

Site PI: Prof Dr med W Wick  
Email: wolfgang.wick@med.uni-heidelberg.de

University of Leipzig (Germany)  
Klinik und Poliklinik für Neurochirurgie  
Leipzig, Germany  
Site PI: Prof Dr J Meixensberger  
Email: meix@medizin.uni-leipzig.de

Ludwig Maximilians University of Munich (Germany)  
Grosshadern Hospital  
Department of Neurosurgery  
München, Germany  
Site PI: Prof Dr J Tonn  
Email: joerg.christian.tonn@med.uni-muenchen.de

University of Regensburg (Germany)  
Department of Neurology  
Regensburg, Germany  
Site PI: PD Dr P Hau  
Email: peter.hau@medbo.de

Medical University Vienna (Austria)  
Department of Internal Medicine I  
Vienna, Austria  
Site PI: Prof Dr C Marosi  
Email: christine.marosi@meduniwien.ac.at

State Mental Hospital of Wagner-Jauregg (Landesnervenklinik Wagner-Jauregg) (Austria)  
Linz, Austria  
Site PI: Dr J Pichler  
Email: josef.pichler@gespag.at

Site location and principal investigator information added as of 05/10/2011:

University Hospital Düsseldorf (Universitätsklinikum Düsseldorf, Neurochirurgische Klinik, Abt. Hirntumorchirurgie)  
Düsseldorf, Germany  
Site PI: Prof. Dr. med. Michael Sabel  
Email: sabel@uni-duesseldorf.de

University Hospital Freiburg (Universitätsklinikum Freiburg, Stereotaktische Neurochirurgie)  
Freiburg, Germany  
Site PI: Prof. Dr. med. Guido Nikkhah  
Email: stereo@uniklinik-freiburg.de

University Hospital Cologne (Universitätsklinikum Köln, Klinik für allgemeine Neurochirurgie)  
Köln, Germany  
Site PI: Prof. Dr. med. Roland Goldbrunner  
Email: roland.goldbrunner@uk-koeln.de

Updated 05/07/2012: the trial was stopped due to a slower than anticipated recruitment rate.

Updated 14/08/2014: the study is completed now and the publication is currently in preparation.

## **Intervention Type**

Drug

## **Phase**

Phase II

## **Drug/device/biological/vaccine name(s)**

Temozolomide

## **Primary outcome(s)**

Median time to treatment failure. Treatment failure is reached:

1. Upon tumour progression, measured every 8 weeks during treatment, every 3 months during follow-up
2. If treatment has to be terminated due to toxicity, measured at every contact between investigator and patient (according to protocol: weekly during first cycle, then every 4 weeks; during follow-up every 3 months)
3. If the patient dies for any reason

## **Key secondary outcome(s)**

1. Progression-free survival (PFS), measured every 8 weeks during treatment, every 3 months during follow-up
2. Overall survival, measured every 8 weeks during treatment, every 3 months during follow-up
3. Objective responses (complete response [CR] and partial response [PR]), measured every 8 weeks during treatment, every 3 months during follow-up
4. Outcome (PFS-6, PFS, survival, best response) relative to MGMT promoter methylation in recurrent tumour, measured every 8 weeks during treatment, every 3 months during follow-up
5. Outcome relative to duration of prior treatment (e.g. number of completed 5/28 cycles after radiation therapy)
6. Outcome relative to interval from completion of prior TMZ chemotherapy treatment (less than 3 months versus greater than 3 months)
7. Toxicity including lymphocytes, CD4 T cell and regulatory T cell counts, measured every 8 weeks during treatment, every 3 months during follow-up
8. Expression of the mismatch repair genes MLH-1, MSH-2, MSH-6 and PMS2 in tumour tissue determined by immunohistochemistry
9. Changes in MGMT status in recurrent disease relative to initial tumour tissue if applicable, measured weekly during first cycle, then every 8 weeks; during follow-up every 3 months
10. Changes in MGMT activity in peripheral blood during ongoing therapy will be assessed during the first cycle at days 1, 8, 15, 22, then MGMT activity will be investigated every 8 weeks; during follow-up every 3 months
11. Quality of life determined by EORTC QoL-Brain 20 Neurotoxicity determined by MRI, measured every 8 weeks during treatment, every 3 months during follow-up
12. Neurotoxicity determined by Mini-Mental State Examination (MMSE), MRI and NeuroCogFx neuropsychological examination, measured every 4 weeks during treatment, every 3 months during follow-up
13. Outcome relative to extent of resection (gross total resection versus resection with residual contrast-enhancing tumour versus biopsy)
14. Screening for aberrant MGMT promoter methylation in peripheral blood, measured weekly during first cycle, then every 8 weeks; during follow-up every 3 months

**Completion date**

16/03/2013

## Eligibility

**Key inclusion criteria**

Current inclusion criteria as of 05/10/2011:

1. Progressive or recurrent glioblastoma documented by magnetic resonance imaging (MRI) no earlier than 180 days after first surgery for glioblastoma and no earlier than 90 days after completion of radiotherapy
2. Histological diagnosis of glioblastoma
3. Tissue available for the determination of MGMT promoter methylation in the primary tumor or from the recurrent tumor of a patient undergoes a surgical procedure at recurrence prior to study entry
4. Prior treatment with temozolomide administered concomitantly with radiotherapy and at least for two cycles (5/28) as an adjuvant treatment
5. Informed consent
6. Aged 18 - 80 years, either sex
7. Karnofsky performance score greater than 50%
8. Neutrophil counts greater than 1,500/ $\mu$ l
9. Platelet counts greater than 100,000/ $\mu$ l
10. Haemoglobin greater than 10 g/dl
11. Serum creatinine less than 1.5-fold upper normal range
12. Aspartate aminotransferase (AST) or alanine aminotransferase (ALT) less than three-fold upper normal range unless attributed to anticonvulsants
13. Alkaline phosphatase less than three-fold upper normal range
14. Women with childbearing potential must have a negative serum pregnancy test less than or equal to 14 days prior to study enrolment

Added 05/10/2011:

15. Willingness to apply contraception according to local requirements (as stated in patient information)

Previous inclusion criteria:

3. Tissue available for the determination of MGMT gene promoter methylation in the recurrent tumour

All other points remained unchanged.

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

All

### **Key exclusion criteria**

1. Progressive or recurrent glioblastoma documented by MRI earlier than 180 days after first surgery for glioblastoma and earlier than 90 days after completion of radiotherapy
2. Treatment with any chemotherapy other than temozolomide according to the schedule of the EORTC NCIC trial except that an adjuvant starting dose of 200 mg/m<sup>2</sup> and more than six cycles of adjuvant temozolomide are allowed
3. Prior systemic or local treatment with deoxyribonucleic acid (DNA)-damaging agents, tyrosine kinase inhibitors or anti-angiogenic agents for any cancer
4. Allergy to or other intolerability of temozolomide
5. Unable to undergo MRI
6. Past medical history of diseases with poor prognosis, e.g. severe coronary heart disease, severe diabetes, immune deficiency, residual deficits after stroke, severe mental retardation
7. Human immunodeficiency virus (HIV) infection
8. Pregnancy
9. Breast feeding
10. Treatment within in any other clinical trial parallel to the treatment phase of the current study

### **Date of first enrolment**

23/09/2009

### **Date of final enrolment**

16/03/2013

## **Locations**

### **Countries of recruitment**

Austria

Germany

Switzerland

### **Study participating centre**

**University Hospital Heidelberg**

Heidelberg

Germany

69120

## **Sponsor information**

### **Organisation**

University Hospital Heidelberg (Universitätsklinikum Heidelberg) (Germany)

ROR

<https://ror.org/013czdx64>

## Funder(s)

### Funder type

Industry

### Funder Name

Essex Pharma GmbH (Germany)

## Results and Publications

### Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/05/2015		Yes	No