

Reduction of severe hypoglycaemia with continuous intraperitoneal insulin infusion in type one diabetic patients unsuccessfully treated by continuous subcutaneous insulin infusion

Submission date

17/09/2006

Recruitment status

No longer recruiting

☐ Prospectively registered

☐ Protocol

Registration date

15/11/2006

Overall study status

Completed

☐ Statistical analysis plan

☐ Results

Last Edited

08/09/2008

Condition category

Nutritional, Metabolic, Endocrine

☐ Individual participant data

☐ Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Andreas Liebl

Contact details

Woernerweg 30

Bad Heilbrunn

Germany

83670

dr.liebl@t-online.de

Additional identifiers

Study information

Scientific Title

Acronym

DiaPort Study

Study objectives

Objective:

Continuous intraperitoneal insulin infusion (CIPII) with the DiaPort system using regular insulin was compared to continuous subcutaneous insulin infusion (CSII) using insulin Lispro, to investigate the frequency of hypoglycaemias, blood glucose control, quality of life, and safety.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved by the Ethical Committee of the Bayerische Landesärztekammer in Munich, September 2000.

Primary study design

Interventional

Study design

Open, randomised, controlled, cross-over, multinational, 12-month study

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Diabetes mellitus type one

Interventions

60 type one diabetic patients with frequent hypoglycaemias and/or HbA1c greater than 7.0% with CSII were randomised to CIPII or CSII. The aim was to obtain the best possible blood glucose while avoiding hypoglycaemias.

1. Patients in the CSII group continued their CSII using insulin Lispro
2. Patients in the CIPII group had an implantation of a percutaneous DiaPort system under general anesthesia. The catheter was placed into the peritoneal cavity. The exact localisation of the DiaPort was chosen individually according to the regular habits and clothing of the patients. Mostly the port was implanted into the lower right or left quadrant of the abdomen. CSII was terminated, and the insulin pump was connected to the DiaPort. The insulin dosage was optimised during the stay in hospital and at each visit.

In both treatment groups, the target was to obtain fasting and pre-prandial blood glucose values between 80 - 120 mg/dl, and average blood glucose values below 150 mg/dl, while avoiding hypoglycemias at the same time. Also, in both groups H-TRONplus insulin pumps from Roche Diagnostics were used. For intraperitoneal infusion only regular insulin for pumps (Insuman Infusat® or H-tronin®, Aventis®) was administered.

In both groups, there were regular evaluations of diabetes complications, vital parameters, HbA1c, safety laboratory data (one central laboratory for all study sites), abdominal ultrasound examination, quality of life (using the Diabetes Quality of Life measure [DQoL]), port-related complications, and photographic documentation.

The patients performed and documented at least four blood glucose self measurements daily (prior to each main meal and just before bedtime). Before visits, additional measurements (two hours after each meal and during night time between 2:00 and 3:00 am) were performed.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Insulin Infusion

Primary outcome(s)

The primary endpoint of the study was the frequency of hypoglycaemias (defined as blood glucose below 54 mg/dl [3 mmol/l]) per patient year with CIPII using DiaPort in comparison to CSII with insulin Lispro.

Key secondary outcome(s)

1. Frequency of severe hypoglycaemias (defined by hospitalisation, unconsciousness, seizures or intravenous glucose administration)
2. Metabolic control (HbA1c, blood glucose, blood glucose fluctuations)
3. Quality of life (DQoL)
4. Safety of CIPII with DiaPort in comparison to CSII

Completion date

31/12/2001

Eligibility

Key inclusion criteria

1. Male or female patients at least 18 years of age
2. Type one diabetes
3. Unsuccessfully treated with CSII (i.e. frequent hypoglycaemias according to the assessment of the investigator and/or HbA1c above 7.0%)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

Not Specified

Key exclusion criteria

1. Lack of cooperation or of mental capacity
2. Pregnancy or wish for pregnancy
3. Abuse of alcohol or drugs
4. Lack of personal hygiene
5. Frequent change of treating physicians
6. Severe liver disease
7. Current malignant disease
8. Human immunodeficiency virus (HIV) infection
9. Continuous ambulatory peritoneal dialysis
10. Contraindications for anaesthesia or surgical operations
11. Severe eating disorders
12. Severe psychological or psychiatric disorders
13. Lack of willingness to perform at least four blood glucose self-measurements per day

Date of first enrolment

01/10/2000

Date of final enrolment

31/12/2001

Locations**Countries of recruitment**

Austria

France

Germany

Netherlands

Switzerland

Study participating centre

Woernerweg 30

Bad Heilbrunn

Germany

83670

Sponsor information**Organisation**

Disetronic Medical Systems AG (Switzerland)

ROR

<https://ror.org/00by1q217>

Funder(s)

Funder type

Industry

Funder Name

Disetronic Medical Systems AG (Switzerland)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration