

# Reiki practice and the quality of life of diabetic pregnant women

<b>Submission date</b> 01/08/2021	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 02/08/2021	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 12/04/2022	<b>Condition category</b> Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Women may get a type of diabetes that only happens in pregnancy. This is called gestational diabetes. Pregnancy can change how a woman's body uses glucose. This can make diabetes worse, or lead to gestational diabetes.

Reiki, a non-drug treatment used on people suffering from anxiety, depression, or both, appears to be an intriguing approach to treating the psychological impact of a woman's response to a diagnosis of gestational diabetes. However, randomized controlled trials comparing various Reiki strategies are required. The purpose of this trial is to analyse if the therapeutic modality of laying on of hands (Reiki) delivered by a trained Reiki practitioner, either face-to-face or remotely, was more effective than Sham-Reiki in controlling anxiety and depression in pregnant women with gestational diabetes.

### Who can participate?

Pregnant women with hyperglycaemia.

### What does the study involve?

The participants were randomized to Face-to-Face Reiki, Remote Reiki, or Sham-Reiki. Changes in the World Health Organization Quality of Life questionnaire, Beck depression inventory and Beck State-Trait anxiety surveys were the primary outcomes after 7 full Reiki sessions by applying standardized and validated surveys in Portuguese.

### What are the possible benefits and risks of participating?

Benefits are related to the improvement of quality of life as well as lowering the levels of depression and anxiety, in addition to have a holistic approach throughout prenatal care. There are no risks involved, since the Sham group continued to receive the medical and nursing care as all the groups. The only difference is that two groups received genuine Reiki while the other not.

### Where is the study run from?

Botucatu Medical School (Brazil)

### When is the study starting and how long is it expected to run for?

March 2016 to March 2019

Who is funding the study?  
Investigator initiated and funded

Who is the main contact?  
Guilherme Ferraz, guilhermerago@hotmail.com

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Mr Guilherme Ferraz

**ORCID ID**  
<https://orcid.org/0000-0001-6596-8782>

**Contact details**  
2902 Universitária Avenue  
Altos do Paraíso  
Botucatu  
Brazil  
18610034  
+55 14 981746951  
guilherme.rago@unesp.br

## Additional identifiers

**Protocol serial number**  
CAAE 52734216.1.0000.5411

## Study information

**Scientific Title**  
Reiki practice and the quality of life of diabetic pregnant women: a randomized clinical trial

**Study objectives**  
The aim of this study was to determine whether the therapeutic modality of laying on of hands (Reiki) may enhance the quality of life, as well as reducing depression and anxiety levels in hyperglycaemic pregnant women

**Ethics approval required**  
Old ethics approval format

**Ethics approval(s)**  
Approved 07/03/2016, Institutional Research Bureau of Botucatu Medical School, UNESP (Chacára Butignoli s/n, Rubião Júnior. Botucatu, State of São Paulo, 18618-970, Brazil; +55 14 3880-1609; cep.fmb@unesp.br), ref: CAAE 52734216.1.0000.5411

**Study design**

Double-blinded randomized controlled trial

**Primary study design**

Interventional

**Study type(s)**

Quality of life

**Health condition(s) or problem(s) studied**

Depression and anxiety levels in hyperglycaemic pregnant women

**Interventions**

Patients were allocated at random by the chief nurse into one of the three groups Sham, Face-to-Face or Remote Reiki at PDRC/HC FMB. Randomization was carried out by the Research Support Office (EAP) of FMB/UNESP using verified software, which is available on the website [www.randomization.com](http://www.randomization.com).

The participants and those evaluating the outcomes were all blinded. The control or intervention groups were only known to the Reiki therapist and Reiki simulation researchers.

The Face-to-Face intervention group had 7 sessions throughout prenatal care. The lead researcher is a licensed Reiki Master and has expertise with Reiki Therapy and has personally carried out the sessions with the approval of the pregnant person by a free explanation of permission and signed informed consent form. Reiki treatments were arranged on the same day as the pregnant woman's outpatient appointment, lasting 30 minutes each.

The Remote intervention group also had 7 sessions, which were also carried by the lead author. The remote treatments were planned on the same day that the pregnant woman returned for her outpatient visit, but without her presence, and lasted 30 minutes each session.

In the control group, participants received a simulation of laying on of hands, the hands' position used in the simulation was the same as the intervention group. This placebo was carried out by another research member who had been trained to imitate the session using the identical touch motions as the intervention group but had neither previous knowledge of Reiki nor competence. In the end, Sham Reiki replicated the identical technique as Face-to-Face Reiki therapy; however, only the researchers who used genuine and sham Reiki knew, and the patients had no idea who was in the control or intervention groups.

**Intervention Type**

Behavioural

**Primary outcome(s)**

Measured at baseline and after 7 sessions:

1. Quality of life measured using the World Health Organization Quality of Life questionnaire
2. Depression measured using the Beck depression inventory
3. Anxiety measured using the Beck State-Trait anxiety survey questionnaire

**Key secondary outcome(s))**

Glycaemic profile measured by blood tests taken throughout pregnancy

**Completion date**

20/03/2019

## Eligibility

**Key inclusion criteria**

1. Pregnant women
2. Fasting glucose 92 mg/dl and/or a risk factor for diabetes
3. Positive screening for Gestational Diabetes Mellitus (GDM), with the diagnosis made by the 75g oral glucose tolerance test (GTT) and glycaemic profile (GP).

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

Female

**Total final enrolment**

134

**Key exclusion criteria**

Does not meet inclusion criteria

**Date of first enrolment**

08/03/2016

**Date of final enrolment**

20/12/2018

## Locations

**Countries of recruitment**

Brazil

**Study participating centre****Perinatal Diabetes Research Centre**

Av. Prof. Montenegro, s/n

Distrito de Rubião Júnior

Botucatu

Brazil

18618-687

# Sponsor information

## Organisation

São Paulo State University

## ROR

<https://ror.org/00987cb86>

# Funder(s)

## Funder type

Other

## Funder Name

Investigator initiated and funded

# Results and Publications

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request ([guilhermerago@hotmail.com](mailto:guilhermerago@hotmail.com)).

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Protocol article</a>		02/06/2021	02/08/2021	Yes	No
<a href="#">Thesis results</a>		27/08/2021	12/04/2022	No	No