

Reduced frequency pembrolizumab immunotherapy: can the frequency of pembrolizumab treatment for non-small cell lung cancer be reduced without reducing its effectiveness?

Submission date 21/01/2022	Recruitment status Recruiting	☒ Prospectively registered ☒ Protocol
Registration date 10/03/2022	Overall study status Ongoing	☐ Statistical analysis plan ☐ Results
Last Edited 22/10/2025	Condition category Cancer	☐ Individual participant data ☒ Record updated in last year

Plain English summary of protocol

<https://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/a-trial-looking-at-how-often-to-give-pembrolizumab-for-non-small-lung-cancer-refine-lung>

Background and study aims

Lung cancer is the most common cause of cancer death. Things have improved with the use of drugs like pembrolizumab, which is often used following a diagnosis of advanced non-small cell lung cancer (NSCLC). Pembrolizumab uses the immune system, the body's natural defence. The immune system sends cells called T cells to fight infections and diseases. Cancer cells hide in the 'PD-1/PD-L1 pathway', allowing them to grow and spread. Pembrolizumab blocks the pathway. This prevents cancer cells from hiding, meaning they can be killed by T cells. The cancer may shrink or disappear as a result.

Treatment is usually every 6 weeks, sometimes with chemotherapy. Treatment can last 2 years, and this may be too much. Research has not shown that length increases benefit, and many people who stop before 2 years continue to benefit from pembrolizumab after it has finished. There is a possibility that researchers can reduce treatment frequency without effectiveness being reduced. This is the main aim of REFINE-Lung.

Who can participate?

Eligible participants will be about to or already receiving pembrolizumab as first treatment for NSCLC.

What does the study involve?

Participants will be randomly allocated to one of the following with or without chemotherapy until the cancer grows significantly:

- Pembrolizumab 6 weekly – the 'control' group, or standard treatment;
- Pembrolizumab 12 weekly.

After 150 participants are recruited, the researchers will see if 12 weekly is similar to 6 weekly, and if so 3 further groups will be opened:

- Pembrolizumab 9 weekly;
- Pembrolizumab 15 weekly;
- Pembrolizumab 18 weekly.

Up to 350 participants will be recruited into each group from up to 45 participating UK hospitals, 1750 in total. Each participant will be followed up for 18 months.

What are the possible benefits and risks of participating?

Benefits:

Similar effectiveness but fewer side effects

Improved quality of life

Fewer visits to and savings by hospitals

Risks:

CT Scans: The risk of causing cancer from one scan is around 1 in 1,000, in a healthy person. The dose from a scan is similar to 9 years of background radiation. Subjects would receive the scans as part of their normal care and the risks are considered low because of their existing condition. The patient may find the CT scanner claustrophobic. A contrast dye may be used which may cause discomfort, bruising, swelling and sometimes an allergic reaction. Severe reactions are very rare.

Blood collection: There is a possibility of redness, swelling and bruising after collection and participants may feel lightheaded or faint. The blood samples collected are not above those that would be performed routinely.

Pembrolizumab: Pembrolizumab is a common treatment for NSCLC. The common side effects are related to the immune system and include itching/rash, diarrhoea, cough, muscle/joint pain, fever, abdominal pain, sickness, headache and tiredness. The research arms reduce the treatment frequency which may reduce the side effects and reduce the burden on patients. Participants may be hesitant about this causing disease progression, hence we have allowed re-escalation to 6-weekly treatment.

Pregnancy: The risks are unknown, however, patients are aware of this having been on this treatment for the previous 6 months. Women who become pregnant will be withdrawn and female and male subjects must agree to take appropriate precautions to avoid pregnancy or fathering a child.

Data Protection: Agreements will be in place prior to the transfer of data externally and pseudo-anonymised will be sent to avoid participant identification.

COVID-19: Patients are included in the at-risk categories but the study aims to reduce the required visits. It is recommended that the vaccine be given as per current guidance for immunotherapy.

Additional visits: The patients will not have to attend hospital more than normal, except for an additional two visits for the screening and end of treatment.

Where is the study run from?

Imperial College London (UK)

When is the study starting and how long is it expected to run for?

January 2022 to August 2028

Who is funding the study?

The National Institute for Health Research (NIHR) Health Technology Assessment (HTA) Programme (UK).

Who is the main contact?
refine-lung@imperial.ac.uk

Contact information

Type(s)

Principal investigator

Contact name

Dr Michael Seckl

Contact details

Du Cane Road
London
United Kingdom
W12 0NN
+44 20 3311 1421
m.seckl@imperial.ac.uk

Type(s)

Scientific

Contact name

Dr Keith Boland

Contact details

RGIT, Room 221
Medical School Buidling
St Marys Campus
Norfolk Place
London
United Kingdom
W2 1PG
+44 20 7594 9480
refine-lung@imperial.ac.uk

Additional identifiers

ClinicalTrials.gov (NCT)

NCT05085028

Clinical Trials Information System (CTIS)

2021-004908-18

Integrated Research Application System (IRAS)

1004165

Central Portfolio Management System (CPMS)

52203

Protocol serial number

C/41/2021

Study information

Scientific Title

A randomised open-label Phase III trial of REduced Frequency pembrolizumab immuNothErapy for first-line treatment of patients with advanced non-small cell lung cancer (NSCLC) utilising a novel multi-arm frequency-response optimisation design

Acronym

REFINE-Lung

Study objectives

To determine the optimal dose frequency of pembrolizumab amongst patients with NSCLC who have benefited from and completed 6 months of standard therapy.

To determine the longest frequency of pembrolizumab treatment that has similar effectiveness in terms of the following when compared to standard 6 weekly pembrolizumab in NSCLC patients who have completed and benefitted from 6 months of standard treatment:

- Overall survival (OS)
- Progression-free survival (PFS), or the length of time that a participant is alive and the cancer has not come back or grown significantly
- Overall response rate (ORR), or the proportion of participants for whom the cancer either disappears or shrinks significantly
- Duration of response (DoR), or, in the proportion of patients for whom the cancer either disappears or shrinks significantly, the length of time between the disappearance or shrinking significantly and the shrinking stopping or the cancer coming back or growing significantly
- Safety and tolerability, or the side effects that are related to pembrolizumab
- Quality of life (QoL)
- Cost-effectiveness

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 09/03/2022, North West - Haydock Research Ethics Committee (Barlow House, 3rd Floor, 4 Minshull Street, Manchester, M1 3DZ, UK; +44 2071048248; haydock.rec@hra.nhs.uk), ref: 22/NW/0037

Primary study design

Interventional

Study design

Interventional randomized controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Lung cancer

Interventions

1. Pembrolizumab (400mg) every 6 weeks for up to 18 months from randomisation. Patients will be followed up for 18 months from randomisation if they stop treatment prior to this date.
 2. Pembrolizumab (400mg) every 12 weeks for up to 18 months from randomisation. Patients will be followed up for 18 months from randomisation if they stop treatment prior to this date.
 3. Pembrolizumab(400mg) every 9 weeks for up to 18 months from randomisation. Patients will be followed up for 18 months from randomisation if they stop treatment prior to this date.
 4. Pembrolizumab (400mg) every 15 weeks for up to 18 months from randomisation. Patients will be followed up for 18 months from randomisation if they stop treatment prior to this date.
 5. Pembrolizumab (400mg) every 18 weeks for up to 18 months from randomisation. Patients will be followed up for 18 months from randomisation if they stop treatment prior to this date.
- Randomisation will occur electronically through the eCRF system, OpenClinica.

Intervention Type

Drug

Phase

Phase III

Drug/device/biological/vaccine name(s)

pembrolizumab

Primary outcome(s)

Overall survival at 2 years, or whether each participant is alive or deceased due to any cause 2 years from initiation of pembrolizumab treatment measured using patient records at 2 years from initiation of pembrolizumab treatment i.e. 18 months from randomisation

Key secondary outcome(s)

1. Median overall survival, or whether each participant is alive or deceased due to any cause measured using patient records throughout the study
2. Median progression free survival, or the length of time a participant is alive without the cancer getting significantly worse as defined by RECIST v1.1 or deceased due to any cause measured using patient records at baseline and then every 12 weeks for a maximum of 2 years from initiation of pembrolizumab
3. Overall response rate, or the proportion of participants with cancer that shows a complete or partial response as defined by RECIST v1.1 measured using patient records at baseline and then every 12 weeks for a maximum of 2 years from initiation of pembrolizumab
4. Median duration of response, or the length of time between a participant with cancer showing a complete or partial response and the cancer getting significantly worse as defined by RECIST v1.1 or death due to any cause measured using patient records at baseline and then every 12 weeks for a maximum of 2 years from initiation of pembrolizumab
5. Safety and tolerability, as defined by adverse events in the Common Terminology Criteria for Adverse Events version 5.0 measured from baseline until the end of treatment visit
6. Quality of life, as defined by the validated EORTC questionnaires QLQ C30 and QLQ LC13 measured from baseline until 2 years from initiation of pembrolizumab
7. Cost effectiveness, as defined by the validated EuroQol EQ-5D measured from baseline until 2 years from initiation of pembrolizumab

Completion date

31/08/2028

Eligibility

Key inclusion criteria

1. Written informed consent prior to initiation of any study procedures and willingness and ability to comply with the study schedule
2. Any patient ≥ 18 years who has received 6 months of pembrolizumab treatment, with or without chemotherapy, for advanced NSCLC who is planned to continue immunotherapy every 6 weeks, or because of continued benefit.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

1. Disease progression or not tolerating treatment at 6 months into therapy
2. Clinician does not intend to continue immunotherapy
3. Is currently receiving an investigational agent or has participated in a study of an investigational agent and or used an investigational device within 28 days of randomisation

Added 04/07/2023:

4. Any patient with a synchronous primary cancer. This includes any new cancer diagnoses or relapse of previously treated cancer since starting pembrolizumab treatment.

Date of first enrolment

20/06/2022

Date of final enrolment

28/02/2027

Locations

Countries of recruitment

United Kingdom

England

Northern Ireland

Scotland

Wales

Study participating centre

Charing Cross Hospital

Fulham Palace Road

London

United Kingdom

W6 8RF

Study participating centre

Weston Park Hospital

Whitham Road

Sheffield

United Kingdom

S10 2SJ

Study participating centre

Royal Surrey County Hospital

Egerton Road

Guildford

United Kingdom

GU2 7XX

Study participating centre

The Royal Marsden Hospital (surrey)

Downs Road

Sutton

United Kingdom

SM2 5PT

Study participating centre

Beatson West of Scotland Cancer Centre

1053 Great Western Road

Glasgow

United Kingdom

G12 0YN

Study participating centre

New Victoria Hospital

52 Grange Rd
Glasgow
United Kingdom
G42 9LF

Study participating centre

Guys and St Thomas' NHS Foundation Trust

249 Westminster Bridge Road
London
United Kingdom
SE1 7EH

Study participating centre

Velindre Cancer Centre

Velindre Rd
Cardiff
United Kingdom
CF14 2TL

Study participating centre

Colchester General Hospital

Colchester District General Hosp.
Charter Way
Turner Road
Colchester
United Kingdom
CO4 5JL

Study participating centre

Ipswich Hospital

Heath Road
Ipswich
United Kingdom
IP4 5PD

Study participating centre

Clatterbridge Cancer Centre - Liverpool

Royal Liverpool University Hospital
Prescot Street

Liverpool
United Kingdom
L7 8XP

Study participating centre
Bristol Haematology & Oncology Centre
Horfield Road
Bristol
United Kingdom
BS2 8ED

Study participating centre
St James's University Hospital
Leeds Teaching Hospitals NHS Trust
Beckett Street
Leeds
United Kingdom
LS9 7TF

Study participating centre
Western General Hospital
Crewe Road South
Edinburgh
Lothian
United Kingdom
EH4 2XU

Study participating centre
Forth Valley Royal Hospital
Stirling Road
Larbert
United Kingdom
FK5 4WR

Study participating centre
Christie Hospital
Wilmslow Road
Manchester
United Kingdom
M20 4BX

Study participating centre
Nottingham City Hospital
Hucknall Road
Nottingham
United Kingdom
NG5 1PB

Study participating centre
Kent and Canterbury Hospital
Ethelbert Road
Canterbury
United Kingdom
CT1 3NG

Study participating centre
Peterborough City Hospital
Edith Cavell Campus
Bretton Gate
Bretton
Peterborough
United Kingdom
PE3 9GZ

Study participating centre
Queens Hospital
Rom Valley Way
Romford
United Kingdom
RM7 0AG

Study participating centre
Leicester Royal Infirmary
Infirmary Square
Leicester
United Kingdom
LE1 5WW

Study participating centre

Royal Devon and Exeter Hospital

Royal Devon & Exeter Hospital
Barrack Road
Exeter
United Kingdom
EX2 5DW

Study participating centre**Royal Cornwall Hospital (treiske)**

Treliske
Truro
United Kingdom
TR1 3LJ

Study participating centre**Royal Derby Hospital**

Uttoxeter Road
Derby
United Kingdom
DE22 3NE

Study participating centre**Royal Sussex County Hospital**

Eastern Road
Brighton
United Kingdom
BN2 5BE

Study participating centre**Royal Bournemouth General Hospital**

Castle Lane East
Bournemouth
United Kingdom
BH7 7DW

Study participating centre**Kettering General Hospital**

Rothwell Road
Kettering
United Kingdom
NN16 8UZ

Study participating centre
North Middlesex Hospital
Sterling Way
London
United Kingdom
N18 1QX

Study participating centre
Yeovil District Hospital NHS Foundation Trust
Higher Kingston
Yeovil
United Kingdom
BA21 4AT

Study participating centre
Barts Health NHS Trust
West Smithfield
London
United Kingdom
EC1A 7BE

Study participating centre
Northampton General Hospital NHS Trust
Cliftonville
Northampton
United Kingdom
NN1 5BD

Study participating centre
Calderdale and Huddersfield NHS Foundation Trust
Trust Headquarters
Acre Street
Lindley
Huddersfield
United Kingdom
HD3 3EA

Study participating centre

St John's Hospital

Howden West
Livingston
Lothian
United Kingdom
EH54 6PP

Study participating centre**North Devon District Hospital**

Raleigh Park
Barnstaple
United Kingdom
EX31 4JB

Study participating centre**The Royal Marsden Hospital**

Fulham Road
London
United Kingdom
SW3 6JJ

Study participating centre**Kingston Hospital**

Galsworthy Road
Kingston upon Thames
United Kingdom
KT2 7QB

Study participating centre**Worthing Hospital**

Lyndhurst Road
Worthing
United Kingdom
BN11 2DH

Study participating centre**Poole General Hospital**

Longfleet Road
Poole
United Kingdom
BH15 2JB

Study participating centre**Victoria Hospital**

Hayfield Road
Kirkcaldy
United Kingdom
KY2 5AH

Study participating centre**Addenbrooke's Hospital**

Addenbrookes Hospital
Hills Road
Cambridge
United Kingdom
CB2 0QQ

Sponsor information

Organisation

Imperial College London

ROR

<https://ror.org/041kmwe10>

Funder(s)

Funder type

Government

Funder Name

Health Technology Assessment Programme

Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location
United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request. Requests should be sent to REFINE-Lung@imperial.ac.uk and will be considered by the TMG and other relevant committees depending on the timing of the request. Consent has been obtained to share data with other researchers for future research. All data shared will be pseudonymised and only identifiable by the trial ID. The researchers requesting data will not be able to identify the participants. Data will not be available until the study has been published. Access criteria has not yet been defined by the TMG.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	version 2.0	21/02/2022	10/03/2022	No	Yes
Participant information sheet	version 2.2	12/07/2022	29/09/2022	No	Yes
Participant information sheet	version 3.0	05/10/2022	23/03/2023	No	Yes
Participant information sheet	version 3.1	10/03/2023	04/07/2023	No	Yes
Protocol file	version 2.0	21/02/2022	10/03/2022	No	No
Protocol file	version 4.0	05/07/2022	29/09/2022	No	No
Protocol file	version 5.0	19/10/2022	23/03/2023	No	No
Protocol file	version 6.0	10/03/2023	04/07/2023	No	No
Protocol file	version 8.0	12/11/2024	22/10/2025	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes