

# CORPUS 1 pilot study: evaluating the impact of physical capacity on the quality of single operator continuous-chest-compression-only cardio pulmonary resuscitation

<b>Submission date</b> 14/01/2010	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 09/02/2010	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
<b>Last Edited</b> 16/05/2012	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Dr Pascal Vranckx

### Contact details

Jessa Hospital  
Stadsomvaart 11  
Hasselt  
Belgium  
3500

## Additional identifiers

## Study information

### Scientific Title

An observational pilot study to evaluate the impact of physical capacity on the quality of single operator continuous-chest-compression-only cardio pulmonary resuscitation

### Acronym

CORPUS

### **Study objectives**

Different components of physical fitness (cardiopulmonary exercise capacity and muscle strength) affects the performance of sustained, single-operator, continuous-external-chest-compressions only resuscitation.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Medical Ethical Committee of Jessa Hospital approved on the 2nd October 2008 (ref: 08.53 /cardio08.11)

### **Study design**

Prospective observational pilot study

### **Primary study design**

Observational

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Critical cardiac care

### **Interventions**

Each subject was instructed to perform sustained, single operator, chest-compression-only cardiocerebral resuscitation (CCR) up to 15 minutes or until exhaustion when the 15-minute CCR session could not be completed. Resuscitation performance was assessed through a special designed laptop based technology (Laerdal PC Skill Reporting System), connected to an adult CCR training manikin (Laerdal, Resusci Anne & Skill Reporter), providing data on compression compliance, including numerical and graphical summaries. Compressions were recorded as correct if both depth and hand placements were in keeping with the standard 2005 International Consensus on Cardiopulmonary Resuscitation guidelines. Adequate compression depth is defined as appropriate between 38 and 51 mm. Adequate compression rate for adults was at least 100 compressions/minute; CCR quality is defined as the number of compressions with adequate depth within a certain timeframe (expressed as %).

### **Intervention Type**

Other

### **Phase**

Not Applicable

### **Primary outcome(s)**

1. Average values of compression rate (ECCs/min) and compression depth (mm), obtained after 1 minute, 3 minutes, and for two consecutive time periods of 5 minutes (5 - 10, 10 - 15)
2. Heart rate during CCR, measured continuously by using a commercial heart rate monitor

(Polar, Oy, Finland)

3. Blood lactate levels (mmol/L), measured with capillary blood samples taken at baseline and after 5, 10 and 15 minutes and using an automated lactate analyser

### **Key secondary outcome(s)**

1. Cardiopulmonary exercise capacity. All subjects performed a maximal cardiopulmonary bicycle exercise test, designed to reach peak oxygen uptake capacity ( $VO_{2peak}$ ) within 8 - 12 minutes, as executed in previous studies from our laboratory. During the cycling test, an electronically braked Ergo 1500 cycle (ErgoFit, Pirmasens, Germany) was used. The cycling frequency was set at 70 cycles/min and the test was ended when the subject failed to maintain a cycling frequency of at least 60 cycles/min. Both the starting and incremental cycling resistance was set between 30 and 45 Watts, depending on age, height, weight, and gender. The cycling resistance increased every single minute. Before every test, a gas and volume calibration was done automatically. During the tests, environmental temperature was kept stable ( $20^{\circ}C$ ). During the exercise tests, pulmonary gas exchange analysis was performed by a cardiopulmonary ergospirometry device (Schiller CS200, Schiller AG, Switzerland). Oxygen uptake capacity ( $VO_2$ ), respiratory exchange ratio (RER), and carbon dioxide output ( $VCO_2$ ) were collected breath-by-breath and averaged every 10 seconds. By plotting exercise  $VO_2$  against exercise  $VCO_2$ , ventilatory threshold was calculated by V-slope method, as executed in previous studies from our laboratory. A software programme of the ergospirometry device determined the break point of the two slopes. Maximal cycling power output ( $W_{peak}$ ) was reported.

2. Muscle strength. Muscle strength measurements of the upper body were performed using a standardised strength testing protocol on an isokinetic dynamometer (Biodex). Following a 5-minute standardised warming-up (Technogym Lat Pull, Pectoral, ArmCurl), subjects were positioned and fixated in a semi supine ( $15^{\circ}$  backward inclination) sitting position (shoulder and thorax fixation). The rotational axis of the dynamometer was aligned with the transverse shoulder/elbow-joint axis and strapped to the distal end of the humerus/radius-ulna. The alignment of the dynamometer was systematically controlled using anatomical reference points. To test upper body strength each subject performed two maximal isometric shoulder flexions, extensions, adductions, abductions, internal and external rotations as well as elbow flexion and extensions (3 seconds) at respective shoulder/elbow angles of  $45^{\circ}$  and  $90^{\circ}$  interspersed by 45-second rest intervals. Maximal isometric torque at each angle was calculated as the average of the manually smoothed static torque curves. After a brief rest period upper body muscle fatigue was tested. Subjects performed one bout of 20 maximal isokinetic ( $180^{\circ}/S$ ) shoulder flexions, extensions, adductions, abductions, internal and external rotations and elbow flexions and extensions from  $90^{\circ}$  to  $20^{\circ}$  (ROM of  $70^{\circ}$ ). After each contraction the upper/lower arm was returned passively to the  $90^{\circ}$  point. Muscle fatigue was calculated as the percentage total work decrease from the first three contractions to the last three. All tests were performed unilaterally. Immediately after the muscle strength measurements, the same examiner would ask subjects to perform dominant hand dynamometry (Jamar handgrip dynamometer; Sammons Preston Rolyan, Bolingbrook, IL) two times. Subjects were standing upright with their elbows at  $90^{\circ}$ .

### **Completion date**

01/02/2010

## **Eligibility**

### **Key inclusion criteria**

1. Active healthcare professionals (12 intensive cardiac care nurses, 3 physicians)
2. Between the ages 18 and 65 years, either sex

3. Working at the Jessa Hospital, Hasselt, Belgium

4. Subjects were required to be able to achieve a maximal voluntary cardiopulmonary exercise test

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

All

**Key exclusion criteria**

Any chronic disease

**Date of first enrolment**

01/07/2009

**Date of final enrolment**

01/02/2010

## **Locations**

**Countries of recruitment**

Belgium

**Study participating centre**

Jessa Hospital

Hasselt

Belgium

3500

## **Sponsor information**

**Organisation**

Heart Centre Hasselt vzw (Belgium)

ROR

https://ror.org/03tw90478

## Funder(s)

### Funder type

Hospital/treatment centre

### Funder Name

Virga Jesse Hospital - Heart Centre Hasselt (Virga Jesse Ziekenhuis - Hart Centrum Hasselt) (Belgium) - provided research grant

### Funder Name

Laerdal BeNeLux (Netherlands) - provided logistic support

## Results and Publications

### Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/02/2012		Yes	No