

Effect of music therapy in mitigating anxiety and enhancing satisfaction in elderly patients undergoing lower limbs orthopedic surgery under intravertebral anesthesia

Submission date 26/03/2024	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 02/04/2024	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 30/06/2025	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

This research examines how music therapy can help reduce feelings of worry and improve overall happiness for older patients who are having surgery on their lower limbs while under anesthesia injected into the spine.

Who can participate?

Older patients having surgery on their legs or hips with anesthesia injected into the spine at the Chinese People's Liberation Army General Hospital

What does the study involve?

Patients will be divided into two groups: the control group (with 60 patients) and the music group (also with 60 patients), using a random number table. Patients in the music group will wear headphones and listen to relaxing music via an MP3 player during the surgery. Patients in the control group will wear the same headphones but will not listen to music during the procedure.

What are the possible benefits and risks of participating?

The benefits of taking part are: (1) Getting access to new treatment options. (2) Lessening the financial strain. (3) Learning more about the most recent information concerning your illness. (4) Receiving additional attention from doctors. (5) Reaching high-quality medical care.

The risks of taking part are: Needing to use more energy and the chance that the treatment may not work.

Where is the study run from?

Department of Orthopedics, PLA General Hospital of China

When is the study starting and how long is it expected to run for?

February 2022 to October 2023

Who is funding the study?
Department of Orthopedics, PLA General Hospital of China

Who is the main contact?
Xuanxuan Zhang, gzgk2021@126.com

Contact information

Type(s)

Public, Scientific, Principal investigator

Contact name

Mr xuanxuan Zhang

ORCID ID

<https://orcid.org/0009-0003-1324-9745>

Contact details

The General Hospital of the People's Liberation Army of the Southern Theater, No. 111, Lihua Road, Yuexiu District
Guangzhou
China
510010
+86 199 6625 1009
gzgk2021@126.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Protocol serial number

20220201

Study information

Scientific Title

Analysis of therapeutic effect of music therapy on patients with lower extremity orthopedic operation under intravertebral anesthesia

Study objectives

Music therapy can reduce anxiety and improve satisfaction in patients undergoing lower extremity orthopedic surgery under intravertebral anesthesia

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 23/02/2024, Ethics Committee of the Chinese People's Liberation Army Southern Theater Command General Hospital (Southern War Zone General Hospital, No. 110, Liuhua Road, Yuexiu District, GuangZhou, 510010, China; +86 20-88686180; gzgcp2012@163.com), ref: None

Study design

Interventional randomized parallel trial

Primary study design

Interventional

Study type(s)

Efficacy

Health condition(s) or problem(s) studied

Patients undergoing lower extremity orthopaedic surgery under intravertebral anesthesia

Interventions

A total of 120 elderly patients will undergo lower extremity orthopaedic surgery under vertebral anaesthesia in the General Hospital of the Chinese People's Liberation Army. Patients will be divided into a control group (n=60) and a music group (n=60) according to a random number table. Patients in the music group will wear headphones and use MP3 to play relaxing music during the operation. Patients in the control group wore the same headphones but did not play music during the procedure.

Intervention Type

Mixed

Primary outcome(s)

1. Levels of perioperative anxiety measured using The Visual Analog Scale for Anxiety (VAS-A), at pre-operation and post-operation.
2. Hemodynamic Parameters: Heart rate, systolic blood pressure (SBP), diastolic blood pressure (DBP), and mean arterial pressure (MAP) measured using continuous monitoring during surgery to provide real-time information on autonomic response using standard medical monitoring equipment, at entry into the operating room (baseline T1), 5 minutes after entering the operating room (T2), before induction of anesthesia (T3), 5 minutes after anesthesia (T4), at the beginning of the operation (T5), 5 minutes after the operation (T6), 30 minutes (T7), 60 minutes of surgery (T8), end of surgery (T9), and one hour after surgery (T10).
3. Salivary Cortisol Levels which is an indicator of stress. Measured using saliva samples collected by the patients themselves using test tubes containing 0.2% trifluoroacetic acid (TFA), followed by centrifugation and measurement using an ELISA (AB154996 Cortisol ELISA kit, Abcam, Cambridge, UK) according to the kit instructions at 15 minutes after waking up on the day of surgery (pre-operation) and the next day (post-operation).
4. Patient Postoperative Satisfaction measured using a standardized homemade questionnaire that includes a question about the satisfaction of the surgery procedure with five answers: A. very satisfied; B. Satisfaction; C. undecided; D. Dissatisfied; E. Very dissatisfied at Post-operation

Key secondary outcome(s)

There are no secondary outcome measures

Completion date

31/10/2023

Eligibility

Key inclusion criteria

1. Age \geq 60 years
2. Undergoing lower limbs orthopedic surgery under intravertebral anesthesia
3. Volunteer to participate in this study and sign the informed consent form

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

60 years

Upper age limit

83 years

Sex

All

Total final enrolment

120

Key exclusion criteria

1. Patients with a history of mental illness, hearing impairment, or contraindications to listening to music;
2. Passive position, completely unable to swing anaesthesia position;
3. Coagulation disorders, bleeding tendency and patients taking anticoagulant drugs
4. Patients with shock, circulatory function is unstable
5. Patients with infection of the skin at the puncture site of the corresponding segment
6. Patients with injuries of the spinal column, which cause changes in the position or patients with a history of trauma to the spinal column

Date of first enrolment

01/02/2022

Date of final enrolment

30/09/2023

Locations

Countries of recruitment

China

Study participating centre

Department of Orthopedics, PLA General Hospital

The General Hospital of the People's Liberation Army of the Southern Theater, No. 111, Liuhua Road, Yuexiu District

Guangzhou

China

510010

Sponsor information

Organisation

PLA General Hospital, Southern Theater Command

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Department of Orthopedics, PLA General Hospital

Results and Publications

Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		30/09/2024	30/06/2025	Yes	No