

# People suffering from Chronic Kidney Disease-associated Pruritus (CKD aP) are treated with a new cream, MC2 25 Cream, or placebo (same Cream without active ingredient) for 12 weeks to reduce their burden and to prove the new cream is safe when used

<b>Submission date</b> 08/02/2022	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 18/05/2022	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 24/10/2025	<b>Condition category</b> Skin and Connective Tissue Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Patients with chronic kidney disease are often suffering from pruritus (severe itching of the skin). This is mostly caused by the lack of correct kidney function and dialysis. The pruritus often presents with fluctuating intensity, often worse during night-time than daytime, resulting in sleep disturbances. Most patients report itching over large, alternating, non-dermatomal regions of skin. The significant impact of itch on these patients' quality of life is generally accepted. In 23,264 haemodialysis patients from 21 countries, the proportion of CKD patients on dialysis who are at least moderately bothered by self-reported pruritus is 37%, and 7% are extremely bothered. It has also been found that this itch was associated with higher risk for hospitalizations and death, higher rates of withdrawal from dialysis and missing scheduled dialysis treatments, and lower rates of employment.

The sponsor of this study has developed a new cream formulation which is deemed to reduce this the itch and therewith the associated complaints and disadvantages in dialysis patients.

### Who can participate?

Adults over 18 years, with chronic kidney disease

### What does the study involve?

Patients will either receive MC2-25-Cream or vehicle (placebo).

### What are the possible benefits and risks of participating?

#### Benefits:

It is anticipated that the disease burden is reduced in participating patients.

#### Risks:

### Study Drug:

This is the first clinical trial with MC2-25 cream so no clinical data are available. According to an SmPC from an IV product containing similar active ingredient, there are no known undesirable effects when administered correctly. In the present trial a maximum of 50 g MC2-25 cream 3% (w/w) will be used per day, which corresponds to ~0.02 g of the active ingredient/kg/day (1.5 g per day in a 70-kg adult) and a resulting safety margin of 25 compared to the IV administration assuming that all topically applied ingredient is absorbed into the systemic circulation.

As for every unknown drug, there may occur adverse effects not yet known for patients receiving MC2-25 cream or even vehicle group, if a patient is allergic to one of the ingredients. This should be minimised by Exclusion Criterion # 9. Furthermore, no application site reactions were observed after dermal application of MC2-25 cream and MC2-25 vehicle to minipigs (twice daily).

### Study procedures:

It is not expected that patients will have a risk when filling study specific Questionnaires. Also, no risk is expected during physical examination, vital sign measurement or ECG. Skin biopsies are scientifically approved for examination of the skin. Patients may recognise the biopsy as unpleasant or painful. Pain induced by skin biopsies is usually not long lasting and disappears within a short time period. Blood samples will be taken from participants at several timepoints to assess the biochemistry and haematology. In some cases nausea could be observed when blood is taken. Furthermore, it might come to discomfort or pain and haematoma at the injection site with local infections. In very rare cases, skin nerves at the injection site can be damaged. Participants receiving MC2-creme could experience an improvement in their disease, which could also occur when patients receive the vehicle. For further information please refer to Table 1, "Summary of Risks and Mitigation Strategies" of the clinical trial protocol.

Where is the study run from?  
SRE GmbH (Germany)

When is the study starting and how long is it expected to run for?  
February 2022 to January 2024

Who is funding the study?  
MC2 Therapeutics Ltd (UK)

Who is the main contact?  
Dr Kieran McCafferty, kieran.mccafferty4@nhs.net  
Jon Bondebjerg, jbo@mc2therapeutics.com

## Contact information

### Type(s)

Principal investigator

### Contact name

Dr Kieran McCafferty

### Contact details

Whitechapel Road  
London  
United Kingdom  
E1 1FR

(+44) 7980 620627  
kieran.mccafferty4@nhs.net

**Type(s)**  
Scientific

**Contact name**  
Dr Jon Bondebjerg

**Contact details**  
MC2 Therapeutics A/S  
Agern Allee 24-26  
Horsholm  
Denmark  
2970  
+49 160 4186243  
jbo@mc2therapeutics.com

## **Additional identifiers**

**Clinical Trials Information System (CTIS)**  
2021-006971-40

**Integrated Research Application System (IRAS)**  
1004785

**Protocol serial number**  
MC-25-C1

## **Study information**

**Scientific Title**  
A parallel-group (2-arm), randomized, double-blind, 12 week trial to evaluate the efficacy and safety of MC2 25 cream and MC2-25 vehicle in subjects with chronic kidney disease-associated pruritus (CKD aP)

**Acronym**  
ITCHINESS

**Study objectives**

1. Exploration of the clinical efficacy of MC2-25 cream compared to MC2-25 vehicle in adults with chronic kidney disease-associated pruritus (CKD-aP)
2. Exploration of the safety of MC2-25 cream compared to MC2-25 vehicle in adults with CKD-aP and other objectives are to explore the subclinical effects of MC2-25 cream in adults with CKD aP

**Ethics approval required**  
Old ethics approval format

**Ethics approval(s)**

Approved 22/03/2022, London City and East REC (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 207 104 8144; cityandeast.rec@hra.nhs.uk), ref: 22/SW/0020

## Study design

Interventional double blind randomized parallel group placebo controlled trial

## Primary study design

Interventional

## Study type(s)

Treatment

## Health condition(s) or problem(s) studied

Patients who are on dialysis and experience itch due to this.

## Interventions

Patients will either receive MC2-25-Cream or vehicle (placebo).

Randomization is done via an electronic online system.

There will be a treatment period of 12 weeks for all arms plus a two week follow-up period (only in case of ongoing AEs).

## Intervention Type

Drug

## Phase

Phase II

## Drug/device/biological/vaccine name(s)

MC2 25 cream

## Primary outcome(s)

Weekly mean WI-NRS recorded in the subject's diary once daily from Baseline to Week 12.

Calculated as the average of minimum 4 WI-NRS values recorded in the subject's diary from 7 days prior to and including the visit days

## Key secondary outcome(s)

There are no secondary outcome measures

## Completion date

02/02/2024

## Eligibility

### Key inclusion criteria

1. Adult males or non-pregnant females of any race or ethnicity who are  $\geq 18$  years of age at the time of screening
2. Able to understand the trial and willing to comply with trial requirements
3. Has provided written informed consent
4. Chronic ( $>3$  months) kidney disease (CKD) stages G3-G5 (i.e., estimated glomerular filtration

rate [eGFR] by CKD-EPI creatinine 2021 equation  $<60 \text{ mL/min/1.73 m}^2$ )

5. Specifically for CKD subjects on haemodialysis (HD) or haemodiafiltration (HDF):

5.1. Subjects must be established on HD or HDF 3 times per week continuously for at least 3 months prior to the start of screening and must not have plans to change from HD to HDF or vice versa during the trial.

5.2. Subjects who require an occasional additional HD or HDF treatment to manage fluid overload may be enrolled as long as it is anticipated that no more than 4 such treatments will be required in any given month.

6. At least moderate CKD-aP defined as WI-NRS  $\geq 4$

7. Female subjects must be of either:

7.1. Non-childbearing potential, i.e., postmenopausal for a least 1 year or have a confirmed clinical history of sterility (e.g., hysterectomy or tubal ligation) or,

7.2. Childbearing potential with a negative urine pregnancy test at the Baseline visit or (in the case of anuria) a negative serum pregnancy test at the Baseline visit that is no more than 3 days old.

8. Female subjects of childbearing potential must agree to use a highly effective method of contraception (i.e., a method with a failure rate of less than 1% per year when used consistently and correctly) while receiving double-blind treatment. Highly effective contraception is defined as follows:

8.1. Combined (estrogen and progestogen-containing) hormonal contraception associated with inhibition of ovulation: oral, intravaginal, transdermal

8.2. Progestogen-only hormonal contraception associated with inhibition of ovulation: oral, injectable, implantable

8.3. Intrauterine device (IUD)

8.4. Intrauterine hormone-releasing system (IUS)

8.5. Bilateral tubal occlusion

8.6. Vasectomised partner (provided that is the sole sexual partner of the subject and that the vasectomised partner has received medical assessment of the surgical success)

8.7. Sexual abstinence if in line with the preferred and usual lifestyle of the subject and defined as refraining from heterosexual intercourse during the entire period of the trial. Periodic methods of abstinence (e.g., calendar, ovulation, symptothermal, post-ovulation methods) are not accepted methods of contraception.

### **Participant type(s)**

Patient

### **Healthy volunteers allowed**

No

### **Age group**

Adult

### **Lower age limit**

18 years

### **Sex**

All

### **Key exclusion criteria**

1. In the opinion of the investigator, the subject is unlikely to comply with the clinical trial protocol.

2. Has a functioning kidney transplant or is scheduled to receive a kidney transplant during the trial
3. Subjects who receive peritoneal dialysis
4. In the opinion of the investigator has pruritus attributed to a cause other than CKD or its complications, including but not limited to dermatological disease (e.g., atopic dermatitis, psoriasis) or liver disease (cholestatic pruritus)
5. Has localized itch restricted to the palms of the hands
6. Only has pruritus during haemodialysis sessions
7. Has concurrent skin conditions that may limit or prevent application of MC2-25 cream or MC2-25 vehicle or that may interfere with evaluation of the effects of MC2-25 cream or MC2-25 vehicle on the skin at the Screening or Baseline visits
8. Subjects who will have skin biopsies performed must not have any known hypersensitivity to the local anaesthetic or diagnosed bleeding disorders. Note: subjects with suspected uremic platelet dysfunction, without other bleeding diatheses, can be enrolled if the investigator agrees.
9. Known history of allergic reaction to any ingredients in MC2-25 cream or MC2-25 vehicle
10. Has a concurrent or recent medical condition that, in the opinion of the investigator, could pose undue risk to the subject, impede completion of the trial procedures, or would compromise the validity of the trial measurements, including, but not limited to:
  - 10.1. known or suspected abuser of alcohol, drugs, or narcotic substances
  - 10.2. severe physical, mental or cognitive disorder other than CKD
  - 10.3. malignancy
  - 10.4. failure to comply with local COVID-19 regulations on vaccination or testing (due to risk of transmitting the disease to other trial participants or trial staff).
11. Has a known current generalized infection
12. Is pregnant, breast feeding, or planning a pregnancy
13. Start of a new or change to existing systemic treatment for CKD-aP, including but not limited to antihistamines, corticosteroids, opioids, GABA analogues, or kappa opioid receptor agonists within 21 days prior to the Baseline visit
14. Use of emollients on CKD-aP areas within 10 days prior to the Baseline visit
15. Use of any topical treatment on CKD-aP areas, including but not limited to antihistamines, or corticosteroids within 21 days prior to the Baseline visit
16. Use of any light therapy for CKD-aP, including but not limited to UV-B within 35 days prior to the Baseline visit
17. Use of non-biologic systemic immunosuppressive treatment, including but not limited to corticosteroids, cyclosporin, and tacrolimus within 5 weeks prior to the Baseline visit
18. Use of biologic systemic treatment, including but not limited to etanercept, adalimumab, alefacept, infliximab, and ustekinumab within 3 months or 5 half-lives (whichever is longer) prior to the Baseline visit
19. Subjects who consent to having skin biopsies performed who are using anticoagulation treatment and are judged by the investigator to have an unacceptable risk of excessive bleeding in association with the skin biopsy
20. Subjects not currently on dialysis but who are likely to initiate routine dialysis during participation in the trial
21. Received another investigational drug within 30 days or 5 half-lives (whichever is longer) prior to screening or is planning to participate in another clinical trial while enrolled in this trial
22. Previously randomized in this trial

**Date of first enrolment**

19/04/2022

**Date of final enrolment**

09/11/2023

**Locations****Countries of recruitment**

United Kingdom

Germany

Hungary

Poland

**Study participating centre****The Royal London Hospital**

Department of Nephrology

Whitechapel Road

Whitechapel

London

United Kingdom

E1 1BB

**Study participating centre****Leicester General Hospital**

Gwendolen Road

Leicester

United Kingdom

LE5 4PW

**Study participating centre****University Hospitals Birmingham NHS Foundation Trust**

Queen Elizabeth Hospital

Mindelsohn Way

Edgbaston

Birmingham

United Kingdom

B15 2GW

**Study participating centre****Kings College Hospital**

Mapother House

De Crespigny Park

Denmark Hill  
London  
United Kingdom  
SE5 8AB

**Study participating centre**  
**West Suffolk NHS Foundation Trust**  
West Suffolk Hospital  
Hardwick Lane  
Bury St. Edmunds  
United Kingdom  
IP33 2QZ

**Study participating centre**  
**University Hospitals Sussex NHS Foundation Trust**  
Worthing Hospital  
Lyndhurst Road  
Worthing  
United Kingdom  
BN11 2DH

**Study participating centre**  
**The Newcastle upon Tyne Hospitals NHS Foundation Trust**  
Freeman Hospital  
Freeman Road  
High Heaton  
Newcastle upon Tyne  
United Kingdom  
NE7 7DN

**Study participating centre**  
**Oxford University Hospitals**  
John Radcliffe Hospital  
Headley Way  
Headington  
Oxford  
United Kingdom  
OX3 9DU

**Study participating centre**

**York and Scarborough Teaching Hospitals NHS Foundation Trust**

York Hospital  
Wigginton Road  
York  
United Kingdom  
YO31 8HE

**Study participating centre**

**Hull University Teaching Hospitals NHS Trust**

Hull Royal Infirmary  
Anlaby Road  
Hull  
United Kingdom  
HU3 2JZ

**Study participating centre**

**Ipswich Hospital NHS Trust**

Heath Road  
Ipswich  
United Kingdom  
IP4 5PD

**Study participating centre**

**South Tyneside and Sunderland NHS Foundation Trust**

Sunderland Royal Hospital  
Kayll Road  
Sunderland  
United Kingdom  
SR4 7TP

**Study participating centre**

**Central Manchester University Hospitals NHS Foundation Trust**

Trust Headquarters, Cobbett House  
Manchester Royal Infirmary  
Oxford Road  
Manchester  
United Kingdom  
M13 9WL

**Study participating centre**

**St Lukes Hospital**  
Little Horton Lane  
Bradford  
United Kingdom  
BD5 0NA

**Study participating centre**  
**Royal Free London NHS Foundation Trust**  
Royal Free Hospital  
Pond Street  
London  
United Kingdom  
NW3 2QG

**Study participating centre**  
**St Pancras Clinical Research**  
285-287 Grays Inn Rd  
Kings Cross  
London  
United Kingdom  
WC1X 8QD

## **Sponsor information**

**Organisation**  
MC2 Therapeutics Ltd

## **Funder(s)**

**Funder type**  
Industry

**Funder Name**  
MC2 Therapeutics Ltd

## **Results and Publications**

**Individual participant data (IPD) sharing plan**

Due to uncertainty about the protection of participants' privacy, the sponsor does not plan to share the dataset(s).

### IPD sharing plan summary

Not expected to be made available

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">HRA research summary</a>			28/06/2023	No	No
<a href="#">Other unpublished results</a>	version 1.0	11/12/2024	24/10/2025	No	No
<a href="#">Study website</a>	Study website	11/11/2025	11/11/2025	No	Yes