

Effect of omacor on heart rate variability (HRV) parameters in patients with recent uncomplicated myocardial infarction

Submission date 13/10/2003	Recruitment status No longer recruiting	☒ Prospectively registered ☒ Protocol
Registration date 13/10/2003	Overall study status Completed	☐ Statistical analysis plan ☐ Results
Last Edited 08/08/2008	Condition category Circulatory System	☐ Individual participant data ☐ Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Cornel Pater

Contact details
An der Trift 18
Hannover
Germany
30559
+49 (0)511 857 2074
cornel.pater@solvay.com

Additional identifiers

Protocol serial number
S1853001

Study information

Scientific Title

Study objectives

To evaluate the effect of Omacor on time-domain heart rate variability (HRV) parameters in comparison to placebo in patients with recent uncomplicated transmural myocardial infarction.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from the local medical ethics committee

Study design

A randomised, parallel group, double-blind, placebo-controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Myocardial infarction

Interventions

Based on the inclusion criteria, a first 24-hour Holter recording will be performed. Two to five days later, screened patients still eligible for the study will undergo a second 24-hour Holter recording. After the second Holter recording, all patients will be randomly allocated to treatment with Omacor 1 g once daily (o.d.) or placebo o.d.

Patients will be followed-up in double-blind fashion for a six-month period after randomisation. Visits, including 24-hour Holter recording and assessment of adverse events, will take place at one-month intervals $\pm$ five days after randomisation, i.e., six times in all.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Omacor

Primary outcome(s)

Superiority of Omacor to improve HRV from baseline to endpoint.

Key secondary outcome(s))

1. Improvement in time domain HRV indices
2. Safety

Completion date

01/01/2005

Eligibility

Key inclusion criteria

1. Males and females aged 40 years or older
2. Recent sustained acute myocardial infarction (AMI)
3. Women of childbearing age are subject to pregnancy testing and must agree to maintain adequate hormonal contraception
4. Signed informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Clinically or hemodynamically unstable condition
2. Further invasive investigation, a percutaneous transluminal coronary angioplasty (PTCA) or a coronary artery bypass graft (CABG) are required
3. Sustained antiarrhythmic therapy (other than a beta-blocking agent administered in the context of secondary prevention of MI)
4. Severe concomitant illness (related to any body organ or system) that is likely to affect outcome assessment
5. Compliance problems
6. Participating in another trial within the past 30 days
7. Pregnant or lactating
8. Known hypersensitivity to the ingredients in Omacor or intolerance to olive oil
9. Diabetes mellitus type I and II

Date of first enrolment

01/01/2004

Date of final enrolment

01/01/2005

Locations

Countries of recruitment

Germany

Lithuania

Poland

Study participating centre
An der Trift 18
Hannover
Germany
30559

Sponsor information

Organisation
Solvay Pharmaceuticals GmbH (Germany)

ROR
<https://ror.org/01xscrc43>

Funder(s)

Funder type
Industry

Funder Name
Solvay Pharmaceuticals GmbH (Germany)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article	Protocol	15/10/2003		Yes	No