

The effectiveness of motorised lumbar traction in the management of lumbosacral nerve root pain

Submission date 25/01/2007	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 21/03/2007	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 16/04/2008	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Jacqueline Gracey

Contact details
School of Health Sciences
University of Ulster
Shore Road
Newtownabbey
United Kingdom
BT37 0QB
+44 (0)28 9036 8284
jh.gracey@ulster.ac.uk

Additional identifiers

Study information

Scientific Title

Study objectives

To compare the effectiveness of two treatment protocols (manual therapy, exercise and advice with or without traction) in the management of acute/subacute Low Back Pain (LBP) with nerve root involvement.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approval received from the local ethics committee at the University Ulster in March 2004.

Study design

Pragmatic single-blind randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Low back pain

Interventions

Group 1: Manual therapy, exercise and advice

Group 2: Manual therapy, exercise, advice and continuous lumbar traction

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

1. Roland Morris Disability Questionnaire
2. McGill pain questionnaire
3. Acute low back pain screening questionnaire
4. Short Form health survey 36

Key secondary outcome(s)

1. Visual analogue scale - pain
2. Medication diary
3. Percentage improved - subjective patient score

Completion date

26/02/2005

Eligibility

Key inclusion criteria

1. Aged 18 to 65 years of age (male and female), presenting with acute/sub-acute LBP with accompanying radiculopathy
2. Nerve root was identified by the presence of:
 - a. dermatomal pain distribution radiating below the knee (one or both limbs)
 - b. sharp/severe quality, often worse in the leg than back (leg pain threshold of 3/10 on Visual Analogue Scale [VAS])With at least one of the following signs and symptoms:
 - c. pins and needles in the distal dermatome (where this was present patients with leg pain were accepted even if not extending below the knee)
 - d. increased pain in the leg on coughing, sneezing or straining
 - e. neurological deficit, i.e., decreased muscle strength/sensory loss/reflex loss
 - f. positive straight leg raise test stretch, i.e., limb pain reproduced on test
3. Acute/sub acute LBP, defined as LBP of less than 12 weeks duration, or a recurrent episode with a pain free period of at least three months prior to the onset of this episode. Only one study has considered recovery rates with sciatica and reported that both back and leg pain decreased, on average, by 69%, and disability decreased by 57% within one month from onset. Current physiotherapy practice would suggest that treatment begins as soon as possible; therefore patients were accepted after four weeks of onset of leg pain
4. Able to attend for physiotherapy two to three times a week for four to six weeks
5. Patients were literate with English as their first language

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Previous spinal surgery
2. Formal therapeutic or medical intervention within the last three months (e.g., epidural injection, facet joint block, physiotherapy etc..)
3. Co-existing conditions (ankylosing spondylitis, rheumatoid arthritis, spinal stenosis [diagnosed], spondylolysis, recent spinal fracture, spinal tumor or a patient where secondary metastases was suspected)
4. Concomitant severe medical problem preventing participation in the trial (cardiac condition, respiratory conditions, neurological disorder or organ disease)
5. Long term oral steroid intake (due to the risk of osteoporosis)
6. Current anti-coagulant therapy or blood clotting disorders
7. Pregnancy

8. History of major psychiatric illness

9. Roland Morris disability questionnaire score of below four, and/or a VAS score of less than three on a ten point scale for leg pain (to avoid floor effects)

Date of first enrolment

01/03/2004

Date of final enrolment

26/02/2005

Locations

Countries of recruitment

United Kingdom

Northern Ireland

Study participating centre

School of Health Sciences

Newtownabbey

United Kingdom

BT37 0QB

Sponsor information

Organisation

University of Ulster (UK)

ROR

<https://ror.org/01yp9g959>

Funder(s)

Funder type

University/education

Funder Name

University of Ulster (UK)

Alternative Name(s)

University of Ulster, Ulster, Ulster Uni, Ollscoil Uladh, Ulstèr Universitè, Ulstèr Varsitè, UU

Funding Body Type

Government organisation

Funding Body Subtype

Universities (academic only)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	29/11/2007		Yes	No