

Matrix metalloproteinases and their tissue inhibitors after selective laser trabeculoplasty in pseudoexfoliative secondary glaucoma

Submission date 13/03/2008	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 20/03/2008	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 30/12/2020	Condition category Eye Diseases	☐ Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Mauro Cellini

Contact details
Via Massarenti, 9
Bologna
Italy
40138

Additional identifiers

Study information

Scientific Title
Matrix metalloproteinases and their tissue inhibitors after selective laser trabeculoplasty in pseudoexfoliative secondary glaucoma

Study objectives
Pseudoexfoliative glaucoma (PEXG) is a degenerative fibrilloglycopathopathy characterised by the production and accumulation of extracellular fibrillar material not just in the anterior segment of the eye, but also in numerous extraocular tissues. At an ocular level, there is atrophy of the

iris dilator muscle fibre cells, degeneration of the irideal pigment with dispersion of melanin, peripupillar atrophy and an increase in the trabecular meshwork pigmentation. A typical feature is the excessive accumulation of the extracellular matrix material in the juxtacanalicular tissue. This accumulation explains the increased resistance to aqueous outflow found in eyes with PEXG.

Study aim:

To assess the variations in the metalloproteinases (MMP-2) and tissue inhibitor of metalloproteinases (TIMP-2) values following selective laser trabeculoplasty (SLT) in patients with pseudoexfoliative glaucoma (PEXG).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from the Institutional Ethics Committee of the St Orsola Malpighi University Hospital, University of Bologna on the 24th October 2007 (ref: 2007-007744-98).

Primary study design

Observational

Study design

Observational, open, single centre, case-control study

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Pseudoexfoliative secondary glaucoma

Interventions

Arm 1: Patients with pseudoexfoliative glaucoma with good intraocular eye pressure (IOP) control and cataract (15 patients). These patients received cataract surgery only.

Arm 2: Patients with pseudoexfoliative glaucoma only that underwent selective laser trabeculoplasty (SLT) without good control of IOP (15 patients). These patients received SLT, and after one month surgical trabeculectomy.

Arm 3 (control group): Patients with cataract only (15 patients). These patients received cataract surgery only.

Selective laser trabeculoplasty (SLT) was performed with Selecta 7000 (Q switched, frequency doubled, 532 Nd:YAG laser) using 50 non-overlapping applications in the inferior 180° of the trabecular meshwork, with a spot size of 400 µm and pulse duration of 3 ns. Aqueous humour was aspirated during surgery from patients.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

MMP-2 and TIMP-2 concentration, assessed once during the cataract surgery in Arm 1 and 3, and during trabeculectomy in Arm 2.

Key secondary outcome(s)

Intraocular pressure, measured before surgery and 7, 15 days and one month after surgery in all three arms of participants.

Completion date

15/06/2006

Eligibility

Key inclusion criteria

Fifteen patients with pseudoexfoliative glaucoma and cataract, and 15 patients with pseudoexfoliative glaucoma (30 patients in total):

1. Patients of either gender
2. Aged from 58 - 74 (in all three arms)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

30

Key exclusion criteria

1. Pseudophachia
2. Non-pseudoexfoliative glaucoma
3. History of laser treatment

Date of first enrolment

13/02/2006

Date of final enrolment

15/06/2006

Locations

Countries of recruitment

Italy

Study participating centre
Via Massarenti, 9
Bologna
Italy
40138

Sponsor information

Organisation
University of Bologna (Italy)

ROR
<https://ror.org/01111rn36>

Funder(s)

Funder type
Hospital/treatment centre

Funder Name
St Orsola Malpighi University Hospital Bologna (Italy)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	21/10/2008	30/12/2020	Yes	No