

Comparative evaluation of immunogenicity of Monovalent Type 1 Oral Poliovirus Vaccine (mOPV1) versus trivalent OPV (tOPV): a randomised double-blind trial set in India (Panacea Biotec Ltd mOPV1 study)

Submission date

01/02/2006

Recruitment status

No longer recruiting

☐ Prospectively registered

☐ Protocol

Registration date

01/02/2006

Overall study status

Completed

☐ Statistical analysis plan

☒ Results

Last Edited

17/01/2012

Condition category

Infections and Infestations

☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Anna-Lea Jenny Kahn

Contact details

World Health Organization

20, Avenue Appia

Geneva-27

Switzerland

CH 1211

+41 (0)22 791 3135

kahna@who.int

Additional identifiers

Protocol serial number

RPC133/Panbio/CR/0982004/CT

Study information

Scientific Title

Study objectives

One dose of mOPV1 induces significantly higher levels of seroconversion against poliovirus type 1 among Indian children in Indore and Hyderabad compared to tOPV.

Please note that as of 18/10/2007 the anticipated end date of this trial was updated from the 31st December 2006 to 28th March 2006.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received on the 20th October 2005.

Study design

Clinical trial, evaluation based, randomised double blind trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Polio

Interventions

Monovalent Type 1 Oral Poliovirus Vaccine (mOPV1) versus Trivalent OPV (tOPV).

Measurements:

1. Cord blood will be collected immediately after birth
2. Blood collection at 30 days of age and stool collection taken
3. During one week follow-up, stool sample collected at day 7

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Monovalent Type 1 Oral Poliovirus Vaccine (mOPV1), Trivalent OPV (tOPV).

Primary outcome(s)

Seroconversion after 30 days of a single dose of tOPV or mOPV1 produced for Biofarma bulk.

Key secondary outcome(s)

1. Prevalence of excretion of Poliovirus type 1 in stool specimens 7 days post-challenge with mOPV1 (age: 30 days + 7 days)
2. Comparison of mOPV1 produced by Biofarma bulk with the mOPV1 produced by Sanofi Pasteur bulk

Completion date

28/03/2006

Eligibility**Key inclusion criteria**

1. Healthy newborns (greater than or equal to 2.75 kg birth weight, apgar score at 5 minutes, greater than or equal to 9) at the study site(s) (large maternity hospitals)
2. Residing within a relatively short and easily accessible distance (less than 30 km)
3. Not planning to travel away during entire the study period (birth - 2 months)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Neonate

Sex

All

Key exclusion criteria

1. Newborns requiring hospitalisation
2. Birth weight below 2.75 kg
3. Apgar score at 5 minutes less than 9
4. Residence greater than 30 km from study site
5. Families expecting to be absent during the 60-day study period
6. A diagnosis or suspicion of immunodeficiency disorder (either in the participant or in a member of the immediate family) will render the newborn ineligible for the study

Date of first enrolment

25/10/2005

Date of final enrolment

28/03/2006

Locations**Countries of recruitment**

India

Switzerland

Study participating centre
World Health Organization
Geneva-27
Switzerland
CH 1211

Sponsor information

Organisation
Panacea Biotech Ltd (India)

ROR
<https://ror.org/01ew11x49>

Funder(s)

Funder type
Research organisation

Funder Name
Gates Foundation (USA) - grant received for a World Health Organization (WHO) Polio Eradication Initiative

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	05/08/2011		Yes	No