

Comparison of the response observed in the dominant and non-dominant hands when assessing neuromuscular block using the TOF-Watch® SX

| | | |
|--|---|--|
| Submission date 30/09/2009 | Recruitment status No longer recruiting | <input type="checkbox"/> Prospectively registered |
| | | <input type="checkbox"/> Protocol |
| Registration date 14/10/2009 | Overall study status Completed | <input type="checkbox"/> Statistical analysis plan |
| | | <input type="checkbox"/> Results |
| Last Edited 07/03/2017 | Condition category Surgery | <input type="checkbox"/> Individual participant data |
| | | <input type="checkbox"/> Record updated in last year |

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Ms Rosemary Hogg

Contact details
Department of Anaesthetics & Intensive Care Medicine
Queens' University Belfast
2nd Floor, Mulhouse Building
Grosvenor Road
Belfast
United Kingdom
BT12 6BJ

Additional identifiers

Protocol serial number
RGHT000353

Study information

Scientific Title

Comparison of neuromuscular monitoring in the dominant and non-dominant hand using the TOF-Watch® SX: a single-centre randomised controlled trial

Study objectives

Is there a difference in the response observed between the dominant and non-dominant hand when using acceleromyography to monitor neuromuscular blockade?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Health and Social Care Research Ethics Committee (HSC REC 3) (Northern Ireland), 23/07/2009, ref: 07/NIR03/1

Study design

Randomised controlled single-centre trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Routine surgery/neuromuscular block

Interventions

After pre-oxygenation, all patients will receive a standard anaesthetic regimen consisting of intravenous infusions of propofol (0.1 - 8.0 µg/ml) and remifentanyl (0.1 - 1.0 µg/kg/min) for induction and maintenance of anaesthesia. After induction, monitoring will be commenced on both arms using the TOF-Watch® SX. The monitors will be applied to the skin over the ulnar nerve at the wrist and calibrated according to Good Clinical Research Practice guidelines. After stabilisation of the TOF-Watch® SX trace in both arms, rocuronium (0.6 mg/kg) will be administered, following which tracheal intubation will be performed.

Surgery will be continued as normal with monitoring continuing until extubation. Reversal with neostigmine will be performed as required.

Intervention Type

Procedure/Surgery

Phase

Not Applicable

Primary outcome(s)

Time to return of TOF (train-of-four) ratio to 0.9

Key secondary outcome(s)

1. Time to onset of block
2. Time to return of T1 to 25%

Completion date

01/04/2009

Eligibility

Key inclusion criteria

1. American Society of Anaesthesiologists (ASA) grade I - III
2. Aged 18 - 65 years, either sex
3. Undergoing planned, elective surgery

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

65 years

Sex

All

Key exclusion criteria

1. Patients with known neuromuscular disease
2. Patients on medication known to interact with neuromuscular blocking agents
3. Pregnancy
4. Patients with known allergy to neuromuscular blocking agents
5. Patients with a known or suspected risk of difficult intubation

Date of first enrolment

01/04/2008

Date of final enrolment

01/04/2009

Locations

Countries of recruitment

United Kingdom

Northern Ireland

Study participating centre
Queens' University Belfast
Belfast
United Kingdom
BT12 6BJ

Sponsor information

Organisation
Belfast Health and Social Care Trust (UK)

ROR
<https://ror.org/02tdmfk69>

Funder(s)

Funder type
Government

Funder Name
Belfast Health and Social Care Trust (UK) (ref: RGHT000353)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration