

# The effect of acupoint application therapy in children with chronic coughs of varying severity

<b>Submission date</b> 22/07/2025	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 24/07/2025	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 23/07/2025	<b>Condition category</b> Signs and Symptoms	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

This study was designed to explore the effects of acupoint application on chronic cough of differing pain severity, based on validation of its efficacy in chronic cough, thereby guiding the clinical application of this therapy.

### Who can participate?

Children aged 4 to 12 with chronic cough

### What does the study involve?

Children with chronic cough were randomly divided into a control group and an observation group. Before receiving treatment, all participants underwent evaluation of cough symptoms and pain levels, which were used as baseline data to assess efficacy.

**Control Group:** After 2 weeks of diagnostic bronchodilator therapy and empirical inhaled corticosteroids (ICS), a reassessment was performed. Treatment involved a single medication or a combination of two to three medications administered via nebulisation, with dosages and methods adjusted according to clinical needs, along with routine symptomatic treatments for allergies, spasms and asthma.

**Observation Group:** In addition to the treatment given to the control group, the observation group underwent acupoint application therapy. Referencing the external treatment methods for cough (the syndrome of qi deficiency of the lung and spleen) outlined in the national textbook *Pediatrics in Chinese Medicine*, edited by Wang Shouchuan, the acupoint application was performed as follows: 12 g of cinnamon, 16 g of clove, 10 g of cassia twig, 15 g of frankincense, 10 g of *Codonopsis pilosula*, 10 g of *Astragalus membranaceus* and 30 g each of *Angelica sinensis*, safflower, red peony, chuanxiong and tuberculate *Speranskia* herb were ground into fine powder (100 mesh) and sealed in a bottle for later use. When used, the powder was mixed with honey to form a paste or shaped into pills the size of peanuts, followed by placing the prepared medicine in the centre of a sterile dressing.

**Acupoint Selection:** Days 1–2: Pills were used at the acupoints of Tiantu and Danzhong; pastes were used bilaterally at both Dingchuan and Feiyu. Days 3–5: Pastes were used at the acupoints of Zhongwan, Shenque, Qihai and Guanyuan.

**Selection of Application Method:** The skin was cleaned with alcohol, and the patient lay supine

to ensure stable application of the medicine. The sterile dressing was then placed on the designated acupoints and gently pressed to ensure full contact between the medicine and the acupoints, followed by securing it with adhesive tape. The application was performed once daily and removed after approximately 4 hours, with a treatment course of 5–7 days. The total intervention lasted 2 weeks.

What are the possible benefits and risks of participating?

**Benefits:** Traditional Chinese medicine can be used as a complementary treatment when conventional treatments are ineffective. Acupoint application is effective in treating chronic cough in children, significantly improving symptoms of chronic cough and increasing patient satisfaction.

**Risks:** Skin scraping and acupoint application may cause adverse reactions such as skin allergies, nausea, vomiting, and diarrhea.

Where is the study run from?

1. Guangdong Provincial Hospital of Traditional Chinese Medicine, China
2. Guangzhou Twelfth People's Hospital, China

When is the study starting and how long is it expected to run for?

April 2023 to December 2024

Who is funding the study?

1. Guangdong Provincial Hospital of Traditional Chinese Medicine, China
2. Guangzhou Twelfth People's Hospital

Who is the main contact?

Dr Lifang Lei, leifangli\_ll70@163.com

## Contact information

### Type(s)

Public, Scientific, Principal investigator

### Contact name

Dr Lifang Lei

### Contact details

No. 111, Dade Road, Yuexiu District

Guangzhou, Guangdong

China

511400

+86 13543478921

leifangli\_ll70@163.com

## Additional identifiers

## Study information

Scientific Title

The effect of acupoint application therapy

### **Study objectives**

To explore the efficacy of acupoint application therapy in children with chronic coughs of varying severity

### **Ethics approval required**

Ethics approval required

### **Ethics approval(s)**

approved 30/05/2023, Medical Ethics Committee of Guangzhou No. 12th People's Hospital (No. 111, Dade Road, Yuexiu District, Guangzhou, Guangdong, 511400, China; +86 020-62134001; gz12yy@126.com), ref: 2023082

### **Study design**

Randomized controlled study

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Children with chronic coughs of varying severity

### **Interventions**

One hundred children with chronic cough were randomly divided into a control group (n=50) and an observation group (n=50).

Participants will be randomly assigned (1:1 ratio) to either group according to a computer-generated sequence, with odd-numbered participants assigned to the control group receiving bronchodilator therapy and empirical inhaled corticosteroids conventional treatment, and even-numbered participants assigned to the observation group receiving acupoint application therapy in addition to conventional treatment.

To avoid the influence of the subjective consciousness of evaluators on the results, two full-time personnel were arranged to conduct efficacy evaluations. The personnel conducting the efficacy evaluation did not participate in other aspects of this study, such as enrolment and intervention.

Before receiving treatment, all participants underwent evaluation of cough symptoms and pain levels, which were used as baseline data to assess efficacy.

Control Group: After 2 weeks of diagnostic bronchodilator therapy and empirical inhaled corticosteroids (ICS), a reassessment was performed.

The following bronchodilators were used:

Ipratropium bromide solution for inhalation (Joincare Pharmaceutical Group Industry Co. Ltd., Approval No.: GYZZ H20203454, product specification: 0.5 mg/vial)

Terbutaline sulphate solution for nebulisation (Joincare Pharmaceutical Group Industry Co. Ltd., Approval No.: GYZZ H20223371, product specification: 5 mg/vial)

Salbutamol sulphate solution for inhalation (Suzhou Homesun Pharmaceutical Co. Ltd., Approval No.: GYZZ H20203292, product specification: 5 mg/vial) Empirical ICS included:  
Budesonide suspension for inhalation (Sichuan Purity Pharmaceutical Co. Ltd., Approval No.: GYZZ H20213286, product specification: 1 mg/vial)  
Fluticasone propionate (GlaxoSmithKline Pty Ltd., Imported Drug Registration No.: H20170361, product specification: 0.5 mg/vial)

Treatment involved a single medication or a combination of two to three medications administered via nebulisation, with dosages and methods adjusted according to clinical needs, along with routine symptomatic treatments for allergies, spasms and asthma.

**Observation Group:** In addition to the treatment given to the control group, the observation group underwent acupoint application therapy. Referencing the external treatment methods for cough (the syndrome of qi deficiency of the lung and spleen) outlined in the national textbook *Pediatrics in Chinese Medicine*, edited by Wang Shouchuan, the acupoint application was performed as follows: 12 g of cinnamon, 16 g of clove, 10 g of cassia twig, 15 g of frankincense, 10 g of *Codonopsis pilosula*, 10 g of *Astragalus membranaceus* and 30 g each of *Angelica sinensis*, safflower, red peony, chuanxiong and tuberculate *Speranskia* herb were ground into fine powder (100 mesh) and sealed in a bottle for later use. When used, the powder was mixed with honey to form a paste or shaped into pills the size of peanuts, followed by placing the prepared medicine in the centre of a sterile dressing.

**Acupoint Selection:** Days 1–2: Pills were used at the acupoints of Tiantu and Danzhong; pastes were used bilaterally at both Dingchuan and Feiyu. Days 3–5: Pastes were used at the acupoints of Zhongwan, Shenque, Qihai and Guanyuan.

**Selection of Application Method:** The skin was cleaned with alcohol, and the patient lay supine to ensure stable application of the medicine. The sterile dressing was then placed on the designated acupoints and gently pressed to ensure full contact between the medicine and the acupoints, followed by securing it with adhesive tape. The application was performed once daily and removed after approximately 4 hours, with a treatment course of 5–7 days. The total intervention lasted 2 weeks.

## **Intervention Type**

Mixed

## **Primary outcome(s)**

Cough symptoms will be measured using the cough symptom diurnal scoring system and cough symptom questionnaire, referencing the Diagnostic Standards for Efficacy of Syndrome in Traditional Chinese Medicine, before treatment and 2 weeks after treatment

## **Key secondary outcome(s)**

The pain level was measured using a Visual Analog Scale (VAS) before treatment and 2 weeks after treatment

## **Completion date**

31/12/2024

## **Eligibility**

### **Key inclusion criteria**

1. Aged 4–12 years
2. Meeting the relevant diagnostic criteria of both TCM and Western medicine as stated above
3. Obvious organic diseases or other severe complications
4. Patients and their parents or guardians willing to participate in the study and able to provide informed consent.

**Participant type(s)**

Patient, Carer

**Healthy volunteers allowed**

No

**Age group**

Child

**Lower age limit**

4 years

**Upper age limit**

12 years

**Sex**

All

**Total final enrolment**

100

**Key exclusion criteria**

1. Severe respiratory complications, e.g. pneumonia or bronchial asthma
2. Received other specific treatments for cough, e.g. antibiotic therapy or immunomodulatory therapy, within the past 3 months
3. Unable to cooperate with acupoint application procedures due to age, cognitive ability or other reasons

**Date of first enrolment**

01/06/2023

**Date of final enrolment**

30/10/2024

**Locations****Countries of recruitment**

China

**Study participating centre**

**Guangzhou no.12 People's Hospital**

No. 111, Dade Road, Yuexiu District

Guangzhou, Guangdong

China  
511400

## Sponsor information

### Organisation

Guangdong Provincial Hospital of Traditional Chinese Medicine

### ROR

<https://ror.org/01gb3y148>

## Funder(s)

### Funder type

Hospital/treatment centre

### Funder Name

Guangdong Provincial Hospital of Traditional Chinese Medicine

### Alternative Name(s)

### Funding Body Type

Government organisation

### Funding Body Subtype

Local government

### Location

China

### Funder Name

Guangzhou Twelfth People's Hospital

## Results and Publications

### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available.

### IPD sharing plan summary

Not expected to be made available