

Non-surgical treatment of benign paroxysmal positional vertigo (BPPV) - a randomised controlled trial of procedures and practices

Submission date 12/09/2003	Recruitment status Stopped	☐ Prospectively registered ☐ Protocol
Registration date 12/09/2003	Overall study status Stopped	☐ Statistical analysis plan ☐ Results
Last Edited 19/10/2011	Condition category Signs and Symptoms	☐ Individual participant data ☐ Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr R Palaniappan

Contact details
Audiological Medicine Department, RNTNE
Royal Free Hampstead NHS Trust
330 Grays Inn Road
Kings Cross
London
United Kingdom
WC1X 8DA
+44 020 7915 1300 ext 4259
rudipal@hotmail.com

Additional identifiers

Protocol serial number
N0256111276

Study information

Scientific Title

Study objectives

How efficient are the different non-surgical treatments that are in current clinical practice? Is any one method better than the other?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Not Specified

Health condition(s) or problem(s) studied

Signs and Symptoms: Benign paroxysmal positional vertigo (BPPV)

Interventions

1. Particle repositioning manoeuvre alone
2. Particle repositioning manoeuvre plus soft collar & sleeping instructions
3. Particle repositioning manoeuvre with bone vibrator
4. Particle repositioning manoeuvre with bone vibrator plus soft collar & sleeping instructions
5. Control group treated with rehabilitation exercises

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

All the above described methods are recognized standard treatment procedures for BPPV in current clinical practice. Patients will be reviewed in the clinic 8 weeks post-treatment, when they will be required to answer the Dizziness Handicap Inventory (DHI). Questionnaire. A negative Hallpike's test during this visit would be considered as a positive outcome.

Key secondary outcome(s)

Not provided at time of registration

Completion date

31/07/2003

Reason abandoned (if study stopped)

Poor recruitment

Eligibility

Key inclusion criteria

Outpatients who have a positive Dix-Hallpike's positional test diagnostic of BPPV
Five groups, approx 20 patients per group. 5 Patients recruited 2001- 2002.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Not Specified

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/02/2002

Date of final enrolment

31/07/2003

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Audiological Medicine Department, RNTNE

London

United Kingdom

WC1X 8DA

Sponsor information

Organisation

Department of Health (UK)

Funder(s)**Funder type**

Government

Funder Name

The Royal Free Hampstead NHS Trust (UK)

Results and Publications**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration