

A study to compare intubation conditions using the CTrach versus the Bonfils rigid fibrescope and CTrach intubating Laryngeal mask airway

Submission date 28/09/2007	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 28/09/2007	Overall study status Completed	☐ Statistical analysis plan ☐ Results
Last Edited 02/09/2015	Condition category Surgery	☐ Individual participant data ☐ Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr PS Venkatesan

Contact details

Anaesthetics Dept
Hull Royal Infirmary
Hull
United Kingdom
HU3 2JZ
+44 627047
abc@email.com

Additional identifiers

Protocol serial number

N0084186834

Study information

Scientific Title

A study to compare intubation conditions using the CTrach versus the Bonfils rigid fibrescope and CTrach intubating Laryngeal mask airway

Study objectives

Compare the airway management devices 'Bonfils' and 'CTrach' with respect to:

1. Time it takes to be placed it successfully in the throat
2. How quickly it helps to successfully place tube in windpipe for patients undergoing general anaesthesia for their operation

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Surgery: Intubation

Interventions

Patients are randomly selected to belong to either Bonfils group or CTrach group.

Procedure

In order to ensure that the conditions under which the study is conducted remains the same in all subjects, the following procedure will be followed at induction by the team managing the patient:

1. Pre-oxygenation for 3 minutes
2. Intravenous administration of Fentanyl 1-2 milligrams per kilogram patient weight to all patients
3. Either of two methods of induction of anaesthesia with target controlled infusion (TCI) of Propofol to target 3-7 milligrams per ml, then maintenance target (TCI) 2.5-4 micrograms per ml with 50% oxygen and 50% air, or intravenous induction of bolus dose of propofol 2-3mg/ml and then anaesthesia maintained with 50% oxygen, 50% air and sevoflurane
4. Muscle relaxation with intravenous administration of Atracurium 0.5 mg/kg to all patients to aim for T0F with 1/4 for adequate muscle relaxation

Following oxygenation, adequate manual ventilation and assessment for adequate level of anaesthesia and relaxation, the following procedure will then follow:

1. Direct laryngoscopy and airway grading by experienced anaesthetist using Macintosh blade (in absence of anaesthetist to use trial device), and using the modified Cormack and Lehane laryngoscopy grading
2. Call back anaesthetist to insert trial device

3. Pick envelope to identify device
4. Start stop clock at beginning of insertion of device and stop it at time capnograph trace is seen
5. Device insertion after appropriate positioning of subjects head

Parameters to be monitored and recorded:

1. Ctrach group
2. Bonfils fibrescope group

Intervention Type

Procedure/Surgery

Phase

Not Specified

Primary outcome(s)

Which of the two devices will be successful and quicker in placing the device in throat and placing the tube in windpipe.

Key secondary outcome(s)

To compare the easiness and quality of the windpipe view obtained, to compare performance of consultants and registrars, to find out any relation between conventional scope grading and successful placement of tube in windpipe with both devices.

Completion date

01/10/2007

Eligibility

Key inclusion criteria

1. Age over 16 years
2. Elective surgery requiring endotracheal intubation
3. ASA status 1-3
4. Airway Mallampati grade 1-3
5. Competency to give informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Morbid obesity (BMI > 35)
2. Pregnancy

3. Emergency surgery or inadequate starvation period
4. Gastro-oesophageal starvation period
5. Gastro-oesophageal reflux or hiatus hernia
6. Severe respiratory disease
7. Mental incapacity
8. Coagulation abnormalities
9. Oral surgery

Date of first enrolment

29/09/2006

Date of final enrolment

01/10/2007

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Hull Royal Infirmary

Hull

United Kingdom

HU3 2JZ

Sponsor information

Organisation

Record Provided by the NHSTCT Register - 2007 Update - Department of Health

Funder(s)

Funder type

Government

Funder Name

The North and South Bank Research and Development Consortium

Funder Name

Hull and East Yorkshire Hospital Trust

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration