

Chronic obstructive pulmonary disease (COPD) as syndrome of accelerated aging

Submission date 15/03/2010	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 18/05/2010	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 18/05/2010	Condition category Respiratory	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Erica Rutten

Contact details
Centre for Integrated Rehabilitation of Organ Failure (CIRO)
P.O. Box 4080
Horn
Netherlands
6080 AB
ericarutten@ciro-horn.nl

Additional identifiers

Protocol serial number
3.2.09.049

Study information

Scientific Title
Systemic manifestation and co-morbidity in chronic obstructive pulmonary disease (COPD) are associated with circulating markers of aging: a cross-sectional observational study with a longitudinal follow-up for two years

Acronym

AGOPD

Study objectives

We hypothesise that accelerated aging is a key pathophysiological mechanism of chronic obstructive pulmonary disease (COPD), and that aging markers are related to important domains of the disease, particularly to the systemic phenotype of COPD and the clinically manifested co-morbidity.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Maastricht Medical Ethical Commission, pending as of 16/03/2010

Study design

Cross-sectional observational study with a longitudinal follow-up

Primary study design

Observational

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Chronic obstructive pulmonary disease (COPD)

Interventions

At baseline and 2 years later, the participants will be invited for two test days; one day at the Center of Expertise for Chronic Organ Failure (CIRO), Horn, and one day at the Maastricht University Medical Center (MUMC).

For COPD patients, the test days will be planned before the start of the rehabilitation. The first day and after overnight fast, venous blood of about 30 ml venous blood in total will be collected, an amount which is not of clinical relevance, but the venepuncture can cause a blue spot. The electrocardiography and the pulse wave velocity will also be performed in the fasted state. During this procedure, the arm will be occluded for 5 minutes. This may give a tingling feeling, but this feeling disappears when the occlusion is removed. Dual x-ray absorptiometry scan will be performed after emptying the bladder and a lung function measurement will take place after consuming breakfast. On the second day at the MUMC, all subjects will be invited for a high resolution computed tomography (HRCT) scan of the thorax.

During the follow-up of 2 years, medical status of the participants will be followed by a telephone contact every three months. For the COPD patients, lung function measurement and dual energy x-ray absorptiometry (DEXA) scan will be performed during the assessment of the rehabilitation at baseline. These tests do not have to be repeated. In a subgroup of 25 patients with the emphysema like phenotype, 25 patients with the non-emphysema like phenotype and 50 smoking healthy controls, circulating concentration of hepatokines and deoxyribonucleic acid (DNA) repair mechanism will be detected in a second venous blood sample during the second test day.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

All analysed at baseline:

1. Markers of aging
2. Objective diagnosed co-morbidity
3. Circulating hepatokines

Key secondary outcome(s)

All analysed at baseline:

1. Markers of systemic inflammation and oxidative stress
2. Classic characterisation of COPD

Completion date

01/10/2014

Eligibility

Key inclusion criteria

COPD patients:

1. Diagnosis of COPD according to the American Thoracic Society (ATS) Global Initiative for Chronic Obstructive Lung Disease (GOLD) guidelines (forced expiratory volume in one second [FEV1] less than 80% predicted and FEV1/forced vital capacity [FVC] less than 70% and less than 10% predicted improvement in FEV1 after O_2 -agonist inhalation
2. Both male and female, aged from 50 to 75 years
3. No respiratory tract infection or exacerbation of the disease for less than 4 weeks before the study
4. Capable of providing informed consent

Healthy subjects:

1. Healthy subjects as judged by a physician
2. Without diagnosed COPD or any other described co-morbidity/chronic disease
3. Both male and female, aged from 50 to 75 years

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

COPD patients:

1. Any kind of carcinogenic pathology less than 5 years before study participation
2. Participation in any other studies involving investigational or marketed products concomitantly or less than 4 weeks prior to entry into the study

Healthy subjects:

1. Investigator's uncertainty about the willingness or ability of the subject to comply with the protocol requirements
2. Participation in any other study involving investigational or marketed products concomitantly or within two weeks prior to entry into the study

Date of first enrolment

01/10/2010

Date of final enrolment

01/10/2014

Locations

Countries of recruitment

Netherlands

Study participating centre

Centre for Integrated Rehabilitation of Organ Failure (CIRO)

Horn

Netherlands

6080 AB

Sponsor information

Organisation

Dutch Asthma Foundation (Netherlands)

ROR

<https://ror.org/00ddgbf74>

Funder(s)

Funder type

Research organisation

Funder Name

Dutch Asthma Foundation (Netherlands)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration