

40 Steps to Safety: Can comparing blood oxygen levels before and after a person has walked 40 steps help to decide whether they can be safely discharged from hospital?

Submission date 27/09/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 31/08/2023	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 06/09/2023	Condition category Signs and Symptoms	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Patients with suspected infection with COVID-19 might appear well but their oxygen levels might dangerously drop on exertion. The National Health Service In England has suggested to ask patients to walk for 40 steps on the flat to assure oxygen levels stay stable before allowing patients with suspected COVID to return home. The 40-steps test has not been used for this purpose before. The aim of this study is to investigate the ability of this test to detect low blood oxygen levels in patients who appear well, to find out whether developing low oxygen levels with exercise can be used to identify patients who are at higher risk of becoming unwell in the future.

Who can participate?

Acutely unwell patients who attend hospitals or are assessed by paramedic crews

What does the study involve?

Participants are asked to undertake the 40 steps test. This will involve taking 40 steps on the spot at their normal walking speed. Researchers will check whether oxygen levels or heart rate change after the test. After 30 days researchers will follow up the participants.

What are the possible benefits and risks of participating?

Participants will be contributing to important research which could help develop a better way to identify patients who can be safely discharged from hospital. They will also be helping us to better understand the normal response to exercise, and how this is different in a range of medical conditions, including in COVID-19.

The researchers anticipate that taking part in this study is generally very safe. However, there is a small possibility of a fall, which could result in injury, whilst doing the 40-step test. Some participants may experience symptoms such as breathlessness, light-headedness, or chest pain whilst taking part in the study.

Where is the study run from?
Betsi Cadwaladr University Health Board (UK)

When is the study starting and how long is it expected to run for?
September 2020 to May 2022

Who is funding the study?
Betsi Cadwaladr University Health Board (UK)

Who is the main contact?
Dr Christian P Subbe, christian.subbe@wales.nhs.uk

Contact information

Type(s)

Public

Contact name

Dr Christian P Subbe

ORCID ID

<https://orcid.org/0000-0002-3110-8888>

Contact details

Ysbyty Gwynedd
Penrhosgarnedd
Bangor
United Kingdom
LL57 2PW
+44(0)1248 384384
christian.subbe@wales.nhs.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

283998

Protocol serial number

IRAS 283998

Study information

Scientific Title

Exertional desaturation as a marker of risk – Validation study for the 40-steps-test: A multi-centre prospective observational cohort study

Study objectives

Absence of desaturation on performing the 40-steps-test is a predictor of safe discharge from hospital.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 16/10/2020, Wales Research Ethics Committee 5 (Health and Care Research Wales, Castlebridge 4, 15-19 Cowbridge Road East, Cardiff, CF11 9AB, UK; +44 (0)7970 422139; Wales. REC5@wales.nhs.uk), ref: 20/WA/0286

Study design

Observational cohort study

Primary study design

Observational

Study type(s)

Screening

Health condition(s) or problem(s) studied

All alert and mobile acutely ill patients considered for discharge to independent care including patients with suspected COVID-19

Interventions

Performance of the 40-steps-test and measurement of oxygen saturations and pulse rate prior to the test and after completion.

Intervention Type

Other

Primary outcome(s)

Validation of the 40 steps on the spot test as a marker for safe discharge from hospital, assessed by collecting the following information from the participants' medical records:

1. Change in decision to discharge following the 40 steps test
2. Outcome at 30 days following the test (30-day hospital admission and 30-day mortality)

Key secondary outcome(s)

Normal values for the 40 steps test challenge in a range of age groups:

1. Oxygen saturation measured by oximeter
2. Heart rate measured by oximeter
3. Breathlessness measured by number of breaths per minute and by using the rating scale for dyspnoea

The above will be measured immediately after the 40 steps test and at 2 minutes after ending the test. Baseline values will be collected from medical records

Completion date

01/05/2022

Eligibility

Key inclusion criteria

1. Patients who are being considered for discharge to independent care
2. Willing and able to give informed consent for participation in the study
3. Independent, stable gait
4. Alert, attentive, coherent and calm

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Minor injuries
2. Elective surgery patients
3. Post-operative patients at discharge
4. Requires supplemental oxygen
5. Shortness of breath at rest (i.e. Borg or Numerical Rating Scale ≥ 2)
6. Unstable angina
7. On long-term-oxygen therapy
8. Pregnancy as stated by patient
9. Oxygen saturation $< 95\%$ on room air
10. Resting heart rate > 100 bpm
11. Resting respiratory rate > 25 bpm
12. ECG with signs of acute ischemia in patients where an ECG has been requested by the treating clinician
13. National Early Warning Score of 5 or more
14. Nursing home residents, or those being transferred to a nursing home or similar care facility

Date of first enrolment

30/11/2020

Date of final enrolment

26/10/2021

Locations**Countries of recruitment**

United Kingdom

Wales

Denmark

Netherlands

Study participating centre

Ysbyty Gwynedd

Penrhosgarnedd

Bangor

United Kingdom

LL57 2PW

Sponsor information

Organisation

Betsi Cadwaladr University Health Board

ROR

<https://ror.org/03awsb125>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Betsi Cadwaladr University Health Board

Alternative Name(s)

Betsi Cadwaladr University Local Health Board, Health Board, Betsi Cadwaladr UHB, Betsi Cadwaladr Local Health Board, Bwrdd Iechyd Prifysgol Betsi Cadwaladr, BCUHB

Funding Body Type

Government organisation

Funding Body Subtype

Local government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown and will be made available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		20/04/2022	06/09/2023	Yes	No
Participant information sheet	version 1.2	30/12/2020	27/07/2023	No	Yes