

Investigation to see if cervical spine mobilisation can influence Cervicobrachial Syndrome

Submission date 28/09/2007	Recruitment status No longer recruiting	☐ Prospectively registered
Registration date 28/09/2007	Overall study status Completed	☐ Protocol
Last Edited 17/10/2016	Condition category Musculoskeletal Diseases	☐ Statistical analysis plan
		☐ Results
		☐ Individual participant data
		☐ Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Ms Emma Salt

Contact details
Musculoskeletal Physiotherapy Lead
Queen's Hospital
Belvedere Road
Burton upon Trent
United Kingdom
DE13 0RB

Additional identifiers

Protocol serial number
N0191187010

Study information

Scientific Title
Investigation to see if cervical spine mobilisation can influence Cervicobrachial Syndrome

Study objectives

1. To investigate whether a self-management protocol with lateral glide mobilisation is more advantageous than a self-management protocol in isolation for subjects with cervicobrachial syndrome.
2. To establish whether subjects with symptoms of parasthesia, numbness and or weakness in their affected arm in addition to their pain have a different outcome from subjects with pain alone.
3. To determine whether subjects preference is linked with outcome.
4. To determine feasibility and enable thorough planning for a phase III trial.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Added 21/09/09: South Staffordshire Local Research Ethics Committee

Study design

Pilot study using a randomised controlled design, with the assessor masked to allocation

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Musculoskeletal Diseases: Cervicobrachial syndrome

Interventions

Subjects will receive either lateral glide mobilisations and a self-management programme (Intervention group) or self-management alone (Control group) for the management of cervicobrachial syndrome. Due to the nature of the intervention, it will not be possible to mask subjects and treating physiotherapists to allocation. Assessment will occur at consent, and 6 weeks after initiation of treatment.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Added 21/09/09:

Visual analogue scale for pain

Key secondary outcome(s))

Added 21/09/09:

1. Neck and Upper Limb Index (NULI)
2. SF36 (RAND)
3. Global Rating of Change Score (GROC)

Completion date

01/06/2008

Eligibility

Key inclusion criteria

This study includes patients with any symptoms radiating from the neck to upper limb.

Added 21/09/09:

1. Patients with pain that had started in the neck and spread to one arm (below level of shoulder joint) in a dermatological distribution and been present for more than two months
2. There were no known systemic rheumatological (e.g. Rheumatoid Arthritis RA or Ankylosing Spondylitis AS) or neurological disorders (e.g. Multiple Sclerosis MS, Parkinson's Disease PD)
3. Not involved with any litigation issues regarding neck and arm pain and no symptoms of myelopathy or sinister pathology
4. No previous fractures to the neck or thoracic spine

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Not Specified

Key exclusion criteria

Does not meet inclusion criteria

Date of first enrolment

01/02/2007

Date of final enrolment

01/06/2008

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Musculoskeletal Physiotherapy Lead

Burton upon Trent

United Kingdom
DE13 ORB

Sponsor information

Organisation

Record Provided by the NHSTCT Register - 2007 Update - Department of Health (UK)

Funder(s)

Funder type

Government

Funder Name

Burton Hospitals NHS Trust (UK), Own Account NHS R&D Support Funding

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration