

# A UK trial for the investigation of stable chest pain: can we improve patient experience, outcomes and NHS cost-efficiency compared to the current NICE guidelines?

<b>Submission date</b> 19/01/2022	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 24/01/2022	<b>Overall study status</b> Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 28/07/2025	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Coronary artery disease (narrowing of the heart arteries) is the typical cause of angina (cardiac chest pain) and is becoming more common as people live longer with multiple heart disease risk factors (e.g. diabetes, smoking, high blood pressure, and high cholesterol). In the UK, about 2 million people have angina and about 200,000 are referred to hospital each year for further investigation. There are lots of different tests to investigate suspected angina, but doctors have different opinions as to the best approach, and international guidelines make very different recommendations. This leads to wide practice variation, inefficiency of healthcare resources and it may adversely impact on patient experience and outcomes. What most doctors agree on is that we do too many invasive X-ray angiograms. About 60% of angiograms performed do not find significant narrowing of the heart arteries, and as such, do not lead to a direct change in management for that patient.

Recent UK NICE guidelines recommend a non-invasive CT angiogram as the first-line test for all patients with suspected angina. This one-size-fits-all approach does not recognise individual patient characteristics, risk factors or likelihood of them having disease. Whilst cardiac CT is an excellent test to exclude narrowing of the heart arteries when they are normal, it can lead to over-estimation of severity when moderate disease is present. It has been suggested that this approach may lead to further increases in the rates of invasive angiography, more additional downstream investigations and increased NHS costs, with no difference in patient outcomes. In addition, the recent UK NICE guidelines if fully adopted would require major service reorganisation in every hospital, and large capital and revenue investment to accommodate the 700% predicted increase in cardiac CT capacity. Thus many doctors believe that this major NHS service reorganisation should first be robustly evaluated in the setting of a clinical trial to demonstrate that it is both appropriate and cost-efficient.

The aim of this study is to try to improve the investigation and management of patients with suspected cardiac chest pain (angina), both in terms of NHS resource use and also patient experience and outcomes; producing results that would be highly generalisable across the NHS.

### Who can participate?

Males aged 45 years and over and females aged 50 years and over with atypical or typical angina and at least one major cardiac risk factor (diabetes, peripheral arterial disease, cerebrovascular disease, current or past tobacco use, high blood pressure, dyslipidaemia or a family history of premature coronary artery disease) referred to NHS cardiology outpatient services, requiring further investigation according to NICE CG95 guidelines, and who are deemed suitable for coronary revascularisation if required.

### What does the study involve?

Patients will have either a computerised tomography cardioangiogram (a CT scan of the heart), a cardiac MRI scan, a stress echocardiogram (an ultrasound scan of the heart) or a nuclear perfusion scan of the heart with single-photon emission computerised tomography (MPS-SPECT), depending on which group they are randomly allocated to and their calculated risk of having heart disease. Patients will also complete three questionnaires at the start of the study and after 6 and 12 months.

### What are the possible benefits and risks of participating?

All of the study procedures are part of routine clinical care and no additional procedures will be undertaken. Participants may be less likely to have unnecessary tests. Patients will also have more contact with the clinical care team and have more opportunities to ask questions and be informed about their health, which some patients find helpful.

### Where is the study run from?

University of Leeds and Leeds Clinical Trials Research Unit (UK)

### When is the study starting and how long is it expected to run for?

May 2019 to April 2027

### Who is funding the study?

1. Heart Research UK
2. British Heart Foundation (UK) (quality of life sub-study)

### Who is the main contact?

Dr Laura Jones  
L.M.Jones@Leeds.ac.uk

## Contact information

### Type(s)

Principal investigator

### Contact name

Prof John Greenwood

### ORCID ID

<https://orcid.org/0000-0002-2861-0914>

### Contact details

Division of Biomedical Imaging  
LIGHT Building  
Clarendon Way

Leeds  
United Kingdom  
LS2 9JT  
+44 (0)113 343 8469  
J.Greenwood@Leeds.ac.uk

**Type(s)**

Principal investigator

**Contact name**

Dr Peter Swoboda

**ORCID ID**

<https://orcid.org/0000-0001-7162-7079>

**Contact details**

Division of Biomedical Imaging  
LIGHT Building  
Clarendon Way  
Leeds  
United Kingdom  
LS2 9JT  
+44 (0)113 39 28206  
P.Swoboda@Leeds.ac.uk

**Type(s)**

Public

**Contact name**

Dr Laura Jones

**Contact details**

Leeds Institute of Cardiovascular and Metabolic Medicine  
University of Leeds  
Leeds  
United Kingdom  
LS2 9JT  
+44 (0)7921121815  
L.M.jones@Leeds.ac.uk

## **Additional identifiers**

**Clinical Trials Information System (CTIS)**

Nil known

**Integrated Research Application System (IRAS)**

302218

**Protocol serial number**

CPMS 50598, IRAS 302218

# Study information

## Scientific Title

A pragmatic approach to the investigation of stable chest pain: a UK, multi-centre, randomised trial to improve patient experience, outcomes and NHS cost efficiency

## Acronym

CE-MARC 3

## Study objectives

The primary question is: 'can we improve the management of patients with new-onset chest pain, both from a patient perspective and an NHS (payer) perspective?'

The proposed trial will address this by answering the following: compared to the current UK NICE guidelines, can a more pragmatic diagnostic pathway to guide the management of patients with suspected cardiac chest pain, result in a) lower rates of invasive angiography without evidence of obstructive disease, b) less downstream resource utilisation and c) improved patient experience and outcomes.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 01/10/2021, London Bridge Research Ethics Committee (postal address: not available; +44 (0)207 1048202; +44 (0)207 1048124; londonbridge.rec@hra.nhs.uk), REC ref: 21/LO/0666

## Study design

Randomized; Both; Design type: Diagnosis, Imaging, Validation of investigation /therapeutic procedures

## Primary study design

Interventional

## Study type(s)

Diagnostic

## Health condition(s) or problem(s) studied

Stable chest pain

## Interventions

People who come to the hospital outpatients department with stable new-onset chest pain or those that have been admitted to cardiology wards will be approached to take part in the trial. If they take part they will be in one of the two arms of the trial and this will be randomly decided (by a computer). The two strategies for the first-line test are:

1. Usual care following the NICE guidelines (this essentially means having a non-invasive CT coronary angiography (CTCA), anatomical test)
2. Pragmatic care following the European Society of Cardiology (ESC) 2019 guidelines whereby a patient could get either a) an MRI scan of the heart (functional test), b) a stress echocardiogram (ultrasound scan) of the heart (functional test), c) a nuclear perfusion scan of the heart (functional test), d) a non-invasive CTCA with added computer calculations (fractional flow

reserve [FFR]-CT, anatomical test) which is no different for the patient than having a CTCA. The choice of the first test in this arm would be by shared decision making between the patient and the doctor (cardiologist) guided by contemporary risk stratification determined by the doctor. There will be no extra visits but patients may be asked to complete three short (optional) questionnaires that ask about their health and well-being at baseline, 6 and 12 months. The questionnaires are validated and are expected to take no longer than 20 minutes to complete in total (at each time point). They can be completed during hospital visits or submitted electronically via the Clinical Trials Research portal (the link will be sent out via email or SMS).

In either arm, if the first test is abnormal patients may be asked to go for an invasive angiogram (this would also be the case if a patient was not in the trial). If the first test is normal then NICE guideline-directed medical therapy would be recommended by the doctor. If the first test is inconclusive then the patient could have second-line non-invasive testing or invasive angiography, based upon shared decision-making and in line with current UK NICE guidelines.

The researchers will collect all the test results to see how many of the invasive angiograms were abnormal in each of the trial arms (and how many were normal - i.e. how many showed no evidence of significant disease). They will analyse questionnaire results completed at baseline, 6 and 12 months, and will review medical records at the same intervals.

Patients will also be followed up for up to 10 years to see if they have any further downstream tests, and if they have any health problems like a heart attack (thankfully only very few people have).

Setting: Secondary care cardiology departments that have well-established clinical NHS services for both anatomical and functional cardiac imaging. Initially nine high volume, experienced research centres that are geographically spread and with ethnically diverse populations have been selected (Leeds, Glasgow, Leicester, Bristol, Oxford, Barts (London), Liverpool, Wakefield and Southampton); other centres could join based on their clinical services and trial experience.

## **Intervention Type**

Other

## **Primary outcome(s)**

Time to composite endpoint of cardiovascular death, myocardial infarction, unobstructive coronary artery disease (CAD) at invasive angiography, defined as the invasive reference standard FFR measurement of  $>0.80$  (or instantaneous wave-free ratio [iFR]  $\geq 0.90$ ), i.e. no functional ischaemia, at the time of coronary angiography (or no coronary stenosis  $>70\%$  on quantitative coronary angiography should FFR/iFR be deemed clinically inappropriate/unsafe to perform)

## **Key secondary outcome(s)**

1. Patient-reported quality of life measured using the Seattle Angina Questionnaire (SAQ), Euroqol EQ-5D-5L, SF12v2 at 0, 6 and 12 months
2. Guideline-directed medical therapy (GDMT) usage measured using patient questionnaire results and/or GP records at 0, 6 and 12 months
3. Major Adverse Cardiovascular Events (all cause death, myocardial infarction [MI], unplanned percutaneous coronary intervention [PCI]/coronary artery bypass graft surgery [CABG]) measured using patient questionnaire results and/or GP records at 3 years
4. Resource use and cost efficiency measured using patient questionnaire results and/or GP records at 12 months

**Completion date**

26/04/2027

## Eligibility

**Key inclusion criteria**

1. Males  $\geq 45$  years or females  $\geq 50$  years
2. Atypical or typical angina
3. At least one major cardiac risk factor (diabetes, peripheral arterial disease, cerebrovascular disease, current or past tobacco use, hypertension, dyslipidaemia or family history of premature coronary artery disease)
4. Referred to NHS cardiology outpatient services
5. Requiring further investigation according to NICE CG95 guidelines
6. Deemed suitable for coronary revascularisation if required

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

45 years

**Sex**

All

**Key exclusion criteria**

In order to keep the trial generalisable to the UK population at large, only issues related to patient safety/appropriateness will form the exclusion criteria; these will include:

1. Prior normal CT coronary angiography (CTCA) within the last 2 years or prior CTCA with extensive calcification
2. Clinically unstable cardiac symptoms
3. Known coronary artery disease (including previous myocardial infarction [MI], acute coronary syndrome [ACS] or coronary revascularization)
4. Contraindication to CTCA or functional cardiac imaging
5. Pregnancy and/or breastfeeding
6. Known chronic renal failure (eGFR  $< 30$ ml/min/1.73m<sup>2</sup>)
7. Inability to give written informed consent

**Date of first enrolment**

27/04/2022

**Date of final enrolment**

08/08/2025

## Locations

**Countries of recruitment**

United Kingdom

England

Scotland

**Study participating centre****United Leeds Teaching Hospital NHS Trust**

Trust Hq

Leeds General Infirmary

Great George St

Leeds

United Kingdom

LS1 3EX

**Study participating centre****Oxford University Hospitals**

John Radcliffe Hospital

Headley Way

Headington

Oxford

United Kingdom

OX3 9DU

**Study participating centre****Pinderfields Hospitals NHS Trust**

Trust Hq, Rowan House

Pinderfields General Hospital

Aberford Road

Wakefield

United Kingdom

WF1 4EE

**Study participating centre****Northumbria Healthcare NHS Foundation Trust**

North Tyneside General Hospital

Rake Lane

North Shields

United Kingdom

NE29 8NH

**Study participating centre**  
**Harefield Hospital NHS Trust**  
Harefield  
United Kingdom  
UB9 6JH

**Study participating centre**  
**Royal Devon University Healthcare NHS Foundation Trust**  
Royal Devon University NHS Ft  
Barrack Road  
Exeter  
United Kingdom  
EX2 5DW

**Study participating centre**  
**South Tees Hospitals NHS Trust**  
Middlesbrough General Hospital  
Ayresome Green Lane  
Middlesbrough  
United Kingdom  
TS5 5AZ

**Study participating centre**  
**University Hospital Southampton NHS Foundation Trust**  
Southampton General Hospital  
Tremona Road  
Southampton  
United Kingdom  
SO16 6YD

**Study participating centre**  
**Central Manchester University Hospitals NHS Foundation Trust**  
Trust Headquarters, Cobbett House  
Manchester Royal Infirmary  
Oxford Road  
Manchester  
United Kingdom  
M13 9WL

**Study participating centre**

**Bradford Teaching Hospitals NHS Foundation Trust**

Bradford Royal Infirmary  
Duckworth Lane  
Bradford  
United Kingdom  
BD9 6RJ

**Study participating centre**

**Nenedoc (kettering)**

Kettering General Hospital  
Rothwell Road  
Kettering  
United Kingdom  
NN16 8UZ

**Study participating centre**

**The Royal Wolverhampton NHS Trust**

New Cross Hospital  
Wolverhampton Road  
Heath Town  
Wolverhampton  
United Kingdom  
WV10 0QP

**Study participating centre**

**Norfolk & Norwich University Hospital**

Colney Lane  
Colney  
Norwich  
United Kingdom  
NR4 7UY

**Study participating centre**

**Nottingham University Hospitals NHS Trust - City Campus**

Nottingham City Hospital  
Hucknall Road  
Nottingham  
United Kingdom  
NG5 1PB

**Study participating centre**

**NIHR Barts Clinical Research Facility**

11th Floor  
Royal London Hospital  
Stepney Way  
London  
United Kingdom  
E1 1BB

**Study participating centre**

**Kings College Hospital**

Mapother House  
De Crespigny Park  
Denmark Hill  
London  
United Kingdom  
SE5 8AB

**Study participating centre**

**Glenfield General Hospital**

Groby Road  
Leicester  
United Kingdom  
LE3 9QP

**Study participating centre**

**The James Cook University Hospital**

Marton Road  
Middlesbrough  
United Kingdom  
TS4 3BW

**Study participating centre**

**NIHR Barts Cardiovascular Biomedical Research Unit**

The Royal London Hospital  
Whitechapel  
London  
United Kingdom  
E1 1BB

**Study participating centre**

**North Bristol NHS Trust**

Southmead Hospital  
Southmead Road  
Westbury-on-trym  
Bristol  
United Kingdom  
BS10 5NB

**Study participating centre****Basildon University Hospital**

Nethermayne  
Basildon  
United Kingdom  
SS16 5NL

**Study participating centre****Queen Elizabeth University Hospital**

1345 Govan Road  
Glasgow  
United Kingdom  
G51 4TF

## Sponsor information

**Organisation**

University of Leeds

**ROR**

<https://ror.org/024mrx33>

## Funder(s)

**Funder type**

Charity

**Funder Name**

Heart Research UK; Grant Codes: TR2442/19/24

**Alternative Name(s)**

HUK

## Funding Body Type

Private sector organisation

## Funding Body Subtype

Trusts, charities, foundations (both public and private)

## Location

United Kingdom

## Funder Name

British Heart Foundation

## Alternative Name(s)

The British Heart Foundation, the\_bhf, BHF

## Funding Body Type

Private sector organisation

## Funding Body Subtype

Trusts, charities, foundations (both public and private)

## Location

United Kingdom

# Results and Publications

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request (Prof. John Greenwood; P.Greenwood@Leeds.ac.uk) following the publication of the pre-specified primary and secondary endpoints. Data will be fully anonymised.

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Protocol article</a>		13/06/2025	18/07/2025	Yes	No
<a href="#">HRA research summary</a>			26/07/2023	No	No
<a href="#">Participant information sheet</a>	version 1.0	16/08/2021	20/01/2022	No	Yes