

Effect of carni Q-gel (ubiquinol and carnitine) on cytokines in patients with heart failure in the TISHCON study

Submission date 14/03/2007	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 30/03/2007	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 08/08/2011	Condition category Circulatory System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Adarsh Kumar

Contact details
Professor and Head of the Department of Cardiology
Government Medical College Amritsar
Punjab
India
143001
adarshk27@jla.vsnl.net.in

Additional identifiers

Study information

Scientific Title

Acronym
The TISHCON Study

Study objectives

Chronic Heart Failure (CHF) represents a major public health burden, and its prognosis is comparable to that of different malignant diseases. CHF progresses because of the activation of neurohormones and pro-inflammatory cytokines, as well as coenzyme Q10 (CoQ) deficiency, following an initial cardiac injury such as Acute Myocardial Infarction (AMI), statin toxicity or a mutation of the genetic programme. It is becoming increasingly apparent that inflammatory mediators play a crucial role in the development of CHF and AMI and several strategies to counterbalance different aspects of the inflammatory response are considered. Possible targets involve pro-and anti-inflammatory cytokines and their receptors, endotoxin, adhesion molecules, nitric oxide and nitric oxide synthase, reactive oxygen species, CoQ and L-carnitine in mitochondria and different types of leucocytes.

The most important cytokines implicated in the progression of CHF are Tumour Necrosis Factor (TNF-alpha), Interleukin (IL)-1, and IL-6. These cytokines share some of their major characteristics (redundancy), and all act in a pro-inflammatory sense. Among those, TNF-alpha is the cytokine that has been studied in greatest detail. IL-10 is anti-inflammatory which is beneficial in heart failure. Cytokines form a vast array of relative low molecular weight, pharmacologically active proteins. These substances are secreted by different cell types for the purpose of altering either their own function (autocrine) or that of adjacent cells (paracrine). Several hypotheses have been suggested to describe the origin of immune activation in CHF. The production of pro-inflammatory cytokines has mostly been attributed to secretion by mononuclear cells, although the myocardium seems to be another important source. Some evidence suggests that catecholamines augment this myocardial cytokine production. There is evidence that CoQ and carnitine may be beneficial in patients with heart failure.

In the present study, we examine, that ubiquinol, in conjunction with L-carnitine, can decrease cytokines and may be useful in patients with CHF.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Reviewed and approved by the Institutional Ethics Committee at Government Medical College, Amritsar, India on the 28th August 2005.

Study design

Double blind, placebo controlled, randomised trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Chronic Heart Failure

Interventions

Nine carni Q-gel softgels (three with each meal thrice daily), which provide a total of 270 mg ubiquinol and 2250 mg L-carnitine daily, or nine matching placebos were administered to each patient, during the period of 12 weeks. All patients were kept on standard drug therapy, and

carni Q-gel and placebo were used as adjuncts to regular drug therapy not as a replacement or substitute.

Laboratory data:

Baseline measurements were taken and repeated after 12 weeks, and a complete blood profile was performed including:

1. Thiobarbituric Acid Reactive Substances (TBARS) and Malondialdehyde (MDA) were obtained by colorimetric methods
2. Resting electrocardiogram and prothrombin time (on anticoagulants) were performed in all the patients at baseline and during follow up whenever indicated
3. CoQ levels in the serum were measured by high pressure liquid chromatography based on an earlier method at Tishcon Corporation (USA)
4. Cytokines Interleukin-six (IL-6), Interleukin-ten (IL-10) and Tumour Necrotising Factors (TNF)-alpha were determined by using a solid phase, two sided chemiluminescence enzyme immunometric assay (immulite automated analyser) kit

The serum samples were kept at -24°C until assay. Results were expressed in pg/ml. The technicians measuring the laboratory data were blind to the groups.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Ubiquinol and carnitine

Primary outcome(s)

1. Increase in distance walked in six minutes
2. NYHA class of heart failure
3. Echocardiographic cardiac size

Key secondary outcome(s)

Reduction in TNF-alpha and IL-6

Completion date

30/09/2006

Eligibility

Key inclusion criteria

1. Both men and women affected by Chronic Heart Failure (CHF) of mixed etiology
2. Ischaemic, hypertensive, valvular heart disease and idiopathic dilated cardiomyopathy
3. Severity of disease was assessed by New York Heart Association (NYHA) functional Class II to IV

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

All

Key exclusion criteria

1. Hypertrophic cardiomyopathy
2. Amyloid cardiomyopathy
3. Not willing to give consent
4. Patients presenting with shock
5. Chronic renal failure
6. Cancer

Date of first enrolment

01/09/2005

Date of final enrolment

30/09/2006

Locations

Countries of recruitment

India

Study participating centre

Professor and Head of the Department of Cardiology

Punjab

India

143001

Sponsor information

Organisation

Tishcon Corporation (USA)

ROR

<https://ror.org/0133gy560>

Funder(s)

Funder type

Industry

Funder Name

Tishcon Corporation (USA)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/08/2007		Yes	No