

# The MAMA Study: Looking at the effects of stopping or continuing biologics for the treatment of inflammatory arthritis in pregnancy

<b>Submission date</b> 01/08/2024	<b>Recruitment status</b> Recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 23/10/2024	<b>Overall study status</b> Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 04/02/2026	<b>Condition category</b> Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

1-2% of women are affected by inflammatory arthritis. Many are treated with bDMARD medications known as 'biologics'. More women with inflammatory arthritis are considering starting a family, because biologics mean they are more able to manage their arthritis. They may face difficult decisions around treatments during pregnancy. Uncontrolled arthritis can lead to worse outcomes in pregnancy, so managing arthritis well is important. Biologics are often avoided during pregnancy because of limited understanding of how these drugs impact pregnancy or arthritis activity. There are concerns about the possible effects of these drugs on babies' immune systems, and some baby vaccinations are routinely delayed. Until recently, most women were advised to stop their biologics during pregnancy; however, due to mounting evidence of their safety, national guidance now states that women can stay on biologics throughout pregnancy. It is unknown whether there is any benefit to this in terms of arthritis control. Certain other medicines used to treat arthritis flares in pregnancy can pose potential harm. We are inviting pregnant women with rheumatoid arthritis (RA), juvenile idiopathic arthritis (JIA), psoriatic arthritis (PsA) or axial spondyloarthritis (axSpA) who are prescribed a regularly dosed biologic to join the MAMA. MAMA aims to find out the effects of stopping or continuing biologics during pregnancy. It will compare whether women who continue their biologics have better arthritis control than those who stop, and assess the impact on their pregnancy, their baby, and the costs associated with this decision.

### Who can participate?

Pregnant women at less than 28 completed weeks' gestation, aged 16 years old and over with a diagnosis of rheumatoid arthritis (RA), juvenile idiopathic arthritis (JIA), psoriatic arthritis (PsA) or axial spondyloarthritis (axSpA) and prescribed a regularly dosed biologic. You do not need to be receiving care at one of the MAMA Participant Sites in order to take part in MAMA. If you would like more information or are interested in taking part in the MAMA Study, please get in touch with the team at the MAMA Co-ordinating Centre.

What does the study involve?

Participants will have an equal (random) chance to continue their biologics throughout pregnancy or to stop by the 28 weeks of pregnancy. During pregnancy, women will be asked to complete a simple arthritis symptom severity questionnaire, monthly via an app (or in writing), and at 3, 6, and 12 months after the baby is born. The woman will be asked if they would be happy for their baby to have blood tests to check their immune response to vaccinations. Mums and babies will be followed up until 24 months after the end of pregnancy, their general health, and their baby's development.

This is a trial of two treatment strategies relating to continuing or stopping prescribed biologics. Biologics are a broad group of drugs which includes many classes and within each class, many marketed products are largely similar in safety profile. The classes of biologics included in this study are detailed in the inclusion criteria and the MAMA trial Reference Safety Information. All drugs are licensed for use in inflammatory arthritis and are already prescribed in pregnancy. Additionally, their use in pregnancy is outlined in the 2022 BSR guidelines on the use of medications in pregnancy.

What are the possible benefits and risks of participating?

Whilst there are no direct benefits to taking part, your contribution may help answer an important question, which could benefit women and their babies in the future.

Both continuing and stopping using biologics beyond 28 weeks of pregnancy are routine practices in the UK, so there are no additional risks to being involved in the trial. Stopping biologic medications can increase the risk of disease activity and flares during pregnancy. Continuing biologic medication in pregnancy may have possible effects on babies' immune systems, and some baby vaccinations may be delayed. This will be discussed with women as part of their clinical care they will also be encouraged to discuss any concerns with their rheumatologist or clinical team.

Optional Infant Immunology part of the Trial

Blood sampling can cause bruising or pain, minimised with local anaesthetic cream.

Where is the study run from?

National Perinatal Epidemiology Unit (NPEU), University of Oxford, UK

When is the study starting and how long is it expected to run for?

July 2024 to February 2030

Who is funding the study?

NIHR Health Technology Assessment Programme, HTA

Who is the main contact?

mama@npeu.ox.ac.uk

## Contact information

**Type(s)**

Public, Scientific

**Contact name**

Mrs Shan Gray

**ORCID ID**

<https://orcid.org/0000-0002-2551-2112>

**Contact details**

MAMA Trial Manager, National Perinatal Epidemiology Unit (NPEU) Clinical Trials Unit  
Nuffield Department of Population Health  
University of Oxford  
Old Road Campus  
Oxford  
United Kingdom  
OX3 7LF  
+44 (0)1865 743859  
mama@npeu.ox.ac.uk

**Type(s)**

Principal investigator

**Contact name**

Prof Marian Knight

**ORCID ID**

<https://orcid.org/0000-0002-1984-4575>

**Contact details**

National Perinatal Epidemiology Unit (NPEU)  
Nuffield Department of Population Health  
University of Oxford  
Old Road Campus  
Oxford  
United Kingdom  
OX3 7LF  
-  
marian.knight@npeu.ox.ac.uk

**Additional identifiers****Integrated Research Application System (IRAS)**

1009876

**Central Portfolio Management System (CPMS)**

65203

**Study information****Scientific Title**

The Monoclonal Antibody Medications in inflammatory Arthritis: stopping or continuing in pregnancy (MAMA) trial

**Acronym**

MAMA

## **Study objectives**

Primary objectives:

To compare the peak of disease activity up to 6 months after the end of pregnancy in pregnant women with AIA randomised to continue bDMARDs versus those randomised to stopping bDMARDs before the third trimester of pregnancy.

Secondary objectives:

In pregnant women with AIA randomised to continue bDMARDs versus those randomised to stopping bDMARDs before the third trimester of pregnancy:

To compare the peak of disease activity up to 12 months after the end of pregnancy

To compare other features of arthritis disease activity

To investigate pregnancy outcomes

In infants born to women with AIA randomised to continue bDMARDs versus those randomised to stop bDMARDs

To compare neonatal outcomes

To investigate infant and child outcomes including global development at 24 months of age, infection up to 24 months of age, and immune function at 2, 5 and 13 months (in a subset of infants where consent has been obtained with an address within geographical reach of Oxford)

Economic evaluation:

To examine using an economic evaluation, whether any additional benefits associated with continuing bDMARDs are justified by any additional health care resources needed

## **Ethics approval required**

Ethics approval required

## **Ethics approval(s)**

approved 30/09/2024, London - Central Research Ethics Committee (3rd Floor 3 Piccadilly Place, London Road, Manchester, M1 3BN, United Kingdom; +44 (0)207 104 8061; londoncentral.rec@hra.nhs.uk), ref: 24/LO/0678

## **Study design**

Interventional randomized controlled trial

## **Primary study design**

Interventional

## **Study type(s)**

Efficacy

## **Health condition(s) or problem(s) studied**

Pregnant women with rheumatoid arthritis (RA), juvenile idiopathic arthritis (JIA), psoriatic arthritis (PsA), or axial spondyloarthritis (axSpA)

## **Interventions**

This trial will compare two existing pathways of care for bDMARD use in pregnancy that are already being used in the UK, albeit with wide variation. MAMA is a pragmatic, comparative effectiveness trial of these two pathways of care.

The two pathways of care being assessed are:

1. Intervention: continuing bDMARDs throughout pregnancy. The woman's current bDMARD, dose and frequency of administration will continue.
2. Comparator: stopping bDMARDs before the third trimester (week 28) of pregnancy and restarting no earlier than 2 weeks after the end of pregnancy.

For both groups, all other aspects of clinical care are determined by the treating clinical team.

## **Intervention Type**

Drug

## **Phase**

Phase III

## **Drug/device/biological/vaccine name(s)**

Humira [adalimumab], Enbrel [etanercept], Remicade [infliximab], Simponi [golimumab], Cimzia [certolizumab pegol], ORENCIA [abatacept], RoActemra [tocilizumab], Kevzara [sarilumab], Kineret [anakinra], Ilaris [canakinumab], Cosentyx [secukinumab], Taltz [ixekizumab], Bimzelx [bimekizumab], Tremfya [guselkumab], Skyrizi [risankizumab], Stelara [ustekinumab]

## **Primary outcome(s)**

Peak disease activity measured by the highest RAPID3 total score [self-report] from randomisation up to 6 months after the end of pregnancy

## **Key secondary outcome(s)**

Measured at variable time points up to 24 months after the end of pregnancy:

1. Peak disease activity measured by the highest RAPID3 total score [self-report]
2. Peak pain level measured by the highest RAPID3 pain score [self-report]
3. Peak overall wellbeing measured by the highest RAPID3 patient global estimate score [self-report]
4. Any occurrence of arthritis flare [self-report]
5. Need for escalation of therapy due to inflammatory disease activity [self-report], defined as:
  - New or increased dose of any DMARD for arthritis;
  - New or increased dose of systemic glucocorticoid (GC) for arthritis (oral or intramuscular injection);
  - Received intra-articular GC joint injection
6. Any use of NSAIDs and frequency of use for treatment of joint pain [self-report] (described only using summary statistics)
7. Any occurrence of arthritis flare [maternal self-report] (described only using summary statistics)
8. Health-related quality of life measured using the EQ-5D-5L
9. Anxiety and depression measured using the EQ-5D-5L
10. Livebirth
11. Stillbirth (fetal loss greater than or equal to 24 weeks' gestation)(described using summary statistics)
12. Pregnancy loss less than 24 weeks' gestation (described using summary statistics)
13. Termination of pregnancy (with or without known congenital anomaly)
14. Mode of birth (described only using summary statistics)
15. Preterm prelabour rupture of membranes (described only using summary statistics)
16. New diagnosis of pre-eclampsia (described only using summary statistics)
17. New diagnosis of gestational diabetes (described only using summary statistics)

18. Venous thromboembolism (described only using summary statistics)
  19. Confirmed or suspected maternal infection (defined as positive culture from a usually sterile site and/or maternal treatment with antibiotics) (described only using summary statistics)
  20. Gestational age (continuous) and preterm birth at:
    - <28
    - 28 to <32 and
    - 32 to 37 gestational weeks
  21. Birth weight z-score
  22. Early-onset neonatal infection: (<72 hours after birth), microbiologically-confirmed or clinically suspected infection
  23. Late-onset infection (>72 hours after birth): microbiologically-confirmed or clinically suspected infection
  24. Neonatal death (up to 24 months)
  25. Neonatal unit admission
  26. Received intensive care (described only using summary statistics)
  27. Necrotising enterocolitis (NEC). [defined as babies born <32+0 gestational weeks where NEC is diagnosed at surgery, post-mortem or based on the following clinical and radiographic signs:
    - 27.1. At least one clinical feature from:
      - Bilious gastric aspirate or emesis
      - Abdominal distension
      - Occult or gross blood in the stool (no fissure)
    - 27.2. And at least one radiographic feature from:
      - Pneumatosis
      - Hepato-biliary gas
      - Pneumoperitoneum] (described only using summary statistics)
28. Brain injury [defined as any of neonatal seizures, intracranial haemorrhage (including intraventricular/periventricular haemorrhage grade 3 or 4), perinatal/neonatal stroke, hypoxic-ischaemic encephalopathy, central nervous system infection, bilirubin encephalopathy and among preterm infants only, cystic periventricular leukomalacia] (described only using summary statistics)
29. Chronic lung disease [defined as babies born <32+0 gestational weeks who are alive and receiving any respiratory support at 36+0 corrected gestational weeks] (described only using summary statistics)
30. Retinopathy of prematurity requiring treatment [defined as babies born <32+0 gestational weeks who received treatment with laser, cryotherapy or intravitreal injection for retinopathy of prematurity] (described only using summary statistics)
31. Any congenital anomaly [classified according to Eurocat] and to include congenital heart block as recommended by EULAR (described only using summary statistics)

Key long-term outcome:

32. Child development (measured using the parent-completed Ages and Stages Questionnaire-3 (ASQ-3) 24-month questionnaire total score)
33. ASQ-3 score below the cut-off for risk of developmental delay in the following domains:
  - Communication skills
  - Gross motor skills
  - Fine motor skills
  - Problem-solving skills
  - Personal social skills(described only using summary statistics)
34. Number of infections (defined by admission to hospital for infection or prescribed antibiotic treatment for infection)
35. Duration of breastfeeding

- 36. Infant death up to 24 months of age
  - 37. Healthcare utilisation and costs  
Including neonatal unit admission level of care: intensive care, high dependency, special care, transitional care, including length of stay at different care levels
  - 38. Quality-adjusted life years (QALYs)
  - 39. Cost-consequence analysis
  - 40. Cost-utility analysis
  - 41. Child Quality of Life (measured using the parent-completed PedsQL)
- Infant Immunology subset infants:
- 42. Child vaccine/immunological response:-
    - Total IgG/A/M immunoglobulin levels
    - Detailed lymphocyte phenotyping
    - Vaccine-specific antibodies (diphtheria, tetanus, pertussis antigens, Hib and measles antibody, and PCV13 at 13 months only)

### **Completion date**

28/02/2030

## **Eligibility**

### **Key inclusion criteria**

Current inclusion criteria as of 20/12/2024:

Pregnant women with Autoimmune Inflammatory Arthritis (AIA), satisfying the following criteria:

1. Have a diagnosis of rheumatoid arthritis (RA), juvenile idiopathic arthritis (JIA), psoriatic arthritis (PsA) or axial spondyloarthritis (axSpA)
2. Pregnant at less than 28 completed weeks' gestation
3. Prescribed a regularly dosed bDMARD (including biologic originators and biosimilars) for RA, JIA, PsA or axSpA.
4. Aged 16 years or over
5. Has provided informed consent

For Infant Immunology (optional element):

6. Address "within geographical reach" from the Oxford Vaccine Group
7. Ongoing consent from parents for infant immunological follow-up

---

Previous inclusion criteria:

Pregnant women with Autoimmune Inflammatory Arthritis (AIA), satisfying the following criteria:

1. Have a diagnosis of rheumatoid arthritis (RA), juvenile idiopathic arthritis (JIA), psoriatic arthritis (PsA) or axial spondyloarthritis (axSpA)
2. Pregnant at less than 28 completed weeks' gestation
3. Prescribed a regularly dosed bDMARD (including biologic originators and biosimilars) for RA, JIA, PsA or axSpA. This includes all current bDMARDs in the following classes and any future bDMARD licensed under these same classes:
  - 3.1. Biologic drugs which block Tumour necrosis factor (TNF)
  - 3.2. Biologic drugs which block CD80/86
  - 3.3. Biologic drugs which block Interleukin 6
  - 3.4. Biologics which block interleukin 1

- 3.5. Biologics which block interleukin 17
- 3.6. Biologics which block interleukin 23
- 3.7. Biologics which block interleukin 12/23
4. Aged 16 years or over
5. Has provided informed consent

For Infant Immunology (optional element):

6. Address "within geographical reach" from the Oxford Vaccine Group
7. Ongoing consent from parents for infant immunological follow-up

### **Participant type(s)**

Patient

### **Healthy volunteers allowed**

No

### **Age group**

Mixed

### **Lower age limit**

16 years

### **Upper age limit**

99 years

### **Sex**

Female

### **Total final enrolment**

0

### **Key exclusion criteria**

1. Prescribed rituximab either during pregnancy or in the 6 months prior to conception
2. Prescribed JAK inhibitors
3. Contraindication to cessation of bDMARDs (e.g. active, sight-threatening uveitis)
4. Current, active tuberculosis in the immediate or close family or household members
5. Plans to move in the first 6 months after birth with their infant to live in a country with a high rate of tuberculosis (incidence >40 per 100,000 population)

For Infant Immunology (optional element):

6. Temporary exclusion criteria for taking immunology samples from the babies - fever in previous 72 hours (or felt to be systemically unwell)

### **Date of first enrolment**

10/02/2025

### **Date of final enrolment**

31/08/2028

## **Locations**

**Countries of recruitment**

United Kingdom

England

Northern Ireland

Scotland

Wales

**Study participating centre**

**Doncaster and Bassetlaw Teaching Hospitals NHS Foundation Trust**

Doncaster Royal Infirmary

Armthorpe Road

Doncaster

England

DN2 5LT

**Study participating centre**

**Oxford University Hospitals**

John Radcliffe Hospital

Headley Way

Headington

Oxford

England

OX3 9DU

**Study participating centre**

**Liverpool Women's NHS Foundation Trust**

Liverpool Womens Hospital

Crown Street

Liverpool

England

L8 7SS

**Study participating centre**

**University Hospitals of Leicester NHS Trust**

Leicester Royal Infirmary

Infirmary Square

Leicester

England

LE1 5WW

**Study participating centre**

**Manchester University NHS Foundation Trust**

Cobbett House  
Oxford Road  
Manchester  
England  
M13 9WL

**Study participating centre**

**Kettering General Hospital NHS Foundation Trust**

Rothwell Road  
Kettering  
England  
NN16 8UZ

**Study participating centre**

**Great Western Hospitals NHS Foundation Trust**

Great Western Hospital  
Marlborough Road  
Swindon  
England  
SN3 6BB

**Study participating centre**

**The Royal Wolverhampton NHS Trust**

New Cross Hospital  
Wolverhampton Road  
Heath Town  
Wolverhampton  
England  
WV10 0QP

**Study participating centre**

**Chelsea and Westminster Hospital NHS Foundation Trust**

Chelsea & Westminster Hospital  
369 Fulham Road  
London  
England  
SW10 9NH

**Study participating centre**

**St George's University Hospitals NHS Foundation Trust**

St George's Hospital

Blackshaw Road

Tooting

London

England

SW17 0QT

**Study participating centre**

**University Hospital Southampton NHS Foundation Trust**

Southampton General Hospital

Tremona Road

Southampton

England

SO16 6YD

**Study participating centre**

**Imperial College Healthcare NHS Trust**

The Bays

St Marys Hospital

South Wharf Road

London

England

W2 1BL

**Study participating centre**

**University Hospitals Birmingham NHS Foundation Trust**

Queen Elizabeth Hospital

Mindelsohn Way

Edgbaston

Birmingham

England

B15 2GW

**Study participating centre**

**Guys and St Thomas' NHS Foundation Trust**

249 Westminster Bridge Road

London

England

SE1 7EH

**Study participating centre**  
**Tunbridge Wells Hospital**  
The Tunbridge Wells Hospital  
Tonbridge Road  
Pembury  
Tunbridge Wells  
England  
TN2 4QJ

## Sponsor information

**Organisation**  
University of Oxford

**ROR**  
<https://ror.org/052gg0110>

## Funder(s)

**Funder type**  
Government

**Funder Name**  
Health Technology Assessment Programme

**Alternative Name(s)**  
NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

**Funding Body Type**  
Government organisation

**Funding Body Subtype**  
National government

**Location**  
United Kingdom

## Results and Publications

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Data sharing statement to be made available at a later date

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Participant information sheet</a>		11/11/2025	11/11/2025	No	Yes
<a href="#">Participant information sheet</a>	version 1.0	18/09/2025	27/01/2026	No	Yes
<a href="#">Study website</a>		11/11/2025	11/11/2025	No	Yes