

Effects of walking with blood flow restriction on intraocular pressure and ocular perfusion pressure

Submission date 08/11/2022	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 16/11/2022	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 14/11/2022	Condition category Other	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Increased intraocular pressure (IOP) and decreased ocular perfusion pressure (OPP) are the main risk factors for the occurrence and development of glaucoma, a common eye condition in which the optic nerve, which connects the eye to the brain, becomes damaged. This study aimed to compare the effects of low-intensity walking with blood flow restriction and moderate-intensity continuous running on IOP and OPP.

Who can participate?

Healthy adult volunteers

What does the study involve?

Participants will be randomly assigned to perform 3 exercise tests:

1. 40% heart rate reserve (HRR) intensity walking (LOW group)
2. 40% HRR intensity walking with blood flow restriction belt of 60% maximum limb occlusive pressure (LOP) (L-BFR group)
3. 60% HRR intensity running (MCE group).

Blood pressure and IOP will be measured before exercise, immediately after exercise, and 5 and 10 minutes after exercise.

What are the possible benefits and risks of participating?

The study could provide potential exercise benefits for healthy people and has limited risks.

Where is the study run from?

Qingdao Haici Hospital (China)

When is the study starting and how long is it expected to run for?

June 2020 to December 2020

Who is funding the study?

1. Natural Science Foundation of Shandong Province (China)
2. Shandong Provincial Natural Science Foundation (China)

Who is the main contact?

Yinghao Li, zhengzhouliyinghao@163.com (China)

Contact information

Type(s)

Principal investigator

Contact name

Dr Yinghao Li

Contact details

Zhengzhou Shengda University

Zhengzhou

China

451191

+86 18203218267

zhengzhouliyinghao@163.com

Additional identifiers

Study information

Scientific Title

Effects of low-intensity walking with blood flow restriction and moderate-intensity running on intraocular pressure and ocular perfusion pressure among healthy adults

Acronym

WALKING+BFR

Study objectives

Low-intensity walking with blood flow restriction was more conducive to improving the ocular perfusion pressure (OPP) level than moderate-intensity continuous running. This may be due to an increase in mean arterial pressure.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 07/06/2020, Qingdao Haici Hospital Ethics Committee (No. 4, Renmin Road, Qingdao, China; +86 0532-83777800; qdhaiciwenhua@163.com), ref: 2022-HYJ-475

Study design

Single-centre cross-sectional cohort study

Primary study design

Observational

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Healthy adults

Interventions

First, we measured baseline height, weight, resting heart rate, resting blood pressure, and maximum limb occlusive pressure (LOP), and then performed an incremental load test on the same day. The treadmill speed was set at 40% and 60% of the heart rate reserve (HRR) using the recorded incremental load test data. Then, subjects were randomly assigned to perform three exercises through a randomization lottery: 1) 40% HRR intensity walking (LOW group); 2) 40% HRR intensity walking with a 60% LOP blood flow restriction belt (L-BFR group); and, 3) moderate-intensity continuous running with 60% HRR intensity (MCE group). Exercises were carried out from 13:00-16:00 h after lunch. Stimulating drinks containing caffeine or alcohol were prohibited. There must be 72 h between the exercise tests. HR and Rating of Perceived Exertion (RPE) were recorded every 5 minutes from before exercise to 10 minutes after exercise; blood pressure and intraocular pressure (IOP) were tested before exercise, immediately, and 5 min and 10 min after exercise. All exercises and tests were performed at the Xinzheng Second People's Hospital. This study adopts a double-blind design, and neither the experimenter nor the participants know the purpose of the experiment in advance.

Each test required 5 min of warm-up activity. The LOW group performed 30 min of walking exercise on a treadmill with 40% HRR intensity. In the L-BFR group, based on the LOW group, a blood flow restriction cuff with 60% LOP pressure should be worn on the proximal end of the thigh, inflated before the exercise, and deflated immediately after the end of the 30-minute exercise. The MCE group performed 30 min of running exercise on a treadmill with 60% HRR intensity.

Intervention Type

Behavioural

Primary outcome(s)

Intraocular pressure and ocular perfusion pressure measured using a non-contact jet tonometer at baseline (T0), immediately after exercise (30 min), and 5 and 10 minutes after exercise

Key secondary outcome(s)

1. Heart rate measured using polar band at baseline (T0) and every 5 minutes until 40 minutes
2. Blood pressure measured using a non-contact jet tonometer at baseline (T0), immediately after exercise (30 min), and 5 and 10 minutes after exercise

Completion date

01/12/2020

Eligibility

Key inclusion criteria

1. Aged \geq 18 years old
2. BMI $<$ 24kg/m²
3. No systematic training experience except for daily physical education
4. No bone, cardiovascular disease or other sports-related discomforts
5. A degree of myopia $<$ 200 degrees

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

16

Key exclusion criteria

Bone, cardiovascular disease or other sports-related discomforts

Date of first enrolment

01/08/2020

Date of final enrolment

20/08/2020

Locations**Countries of recruitment**

China

Study participating centre

Qingdao Haici Hospital

No. 4 Renmin Road

Qingdao

China

266033

Sponsor information

Organisation

Ocean University of China

ROR

<https://ror.org/04rdtx186>

Funder(s)**Funder type**

Government

Funder Name

Natural Science Foundation of Shandong Province

Alternative Name(s)

Shandong Provincial Natural Science Foundation, Shandong Natural Science Foundation, Natural Science Foundation of Shandong, Shandong Province Natural Science Foundation,

Funding Body Type

Government organisation

Funding Body Subtype

Local government

Location

China

Funder Name

Natural Science Foundation of Shandong Province

Alternative Name(s)

Shandong Provincial Natural Science Foundation, Shandong Natural Science Foundation, Natural Science Foundation of Shandong, Shandong Province Natural Science Foundation,

Funding Body Type

Government organisation

Funding Body Subtype

Local government

Location

China

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Yinghao Li, zhengzhouliyinghao@163.com. Anonymized intraocular pressure and ocular perfusion pressure data will be shared. Consent was required and obtained from participants.

IPD sharing plan summary

Available on request