

# Comparing inhaled nitrous oxide gas with cognitive behavioral therapy to reduce anxiety in children undergoing dental treatment

<b>Submission date</b> 30/11/2018	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 07/01/2019	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 11/09/2020	<b>Condition category</b> Mental and Behavioural Disorders	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Many people become anxious when they have dental treatment. This can put them off going to the dentist regularly, which might result in teeth being in a worse condition and needing more invasive treatment. Often the anxiety stems from a stressful experience at the dentist in childhood. This study aims to compare two methods of calming children during dental treatment. Nitrous oxide can be mixed with air and inhaled through a nose mask. It reduces anxiety, pain and the gag reflex. It is safe and the effects start and wear off quickly, so the amount breathed in can be easily adjusted if the patient is in pain or feeling too woozy. Cognitive behavioural therapy (CBT) is a talking therapy that help people understand and deal with distressing feelings.

### Who can participate?

Children aged 5 to 12 years who have dental anxiety and need pulp treatment in at least one molar tooth in the lower jaw.

### What does the study involve?

Participants will be randomly allocated to receive nitrous oxide or CBT during the dental appointment. They will describe their anxiety levels by selecting from a set of faces before and after the dental treatment. They will also provide saliva samples before and after treatment so that levels of the stress hormone cortisol can be measured.

### What are the possible benefits and risks of participating?

Patients will receive a complete dental treatment and oral health promotion as a benefit of participating in the study. All participants may leave the study any time and are informed about all the procedures that will be conducted. Nitrous oxide inhalation may provoke side effects, such as nausea or vomiting, but these effects are reversed quickly if the inhalation mask is removed.

### Where is the study run from?

Universidad Austral de Chile (Chile)

When is the study starting and how long is it expected to run for?  
January 2018 to December 2018

Who is funding the study?  
Universidad Austral de Chile (Chile)

Who is the main contact?  
1. Mrs Bruna Benso (scientific contact), bruna.benso@uc.cl  
2. Mrs Claudia Mautz (public contact), cpazmautz@gmail.com

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Mrs Bruna Benso

**ORCID ID**  
<https://orcid.org/0000-0002-4425-5174>

**Contact details**  
Marcoleta, 391  
Santiago  
Chile  
8320000

**Type(s)**  
Public

**Contact name**  
Mrs Claudia Mautz

**Contact details**  
Rudloff 1640  
Valdivia  
Chile  
5090000

## Additional identifiers

**Protocol serial number**  
2017-30

## Study information

**Scientific Title**  
Comparison of nitrous oxide inhalation and cognitive behavioral therapy for anxiety in pediatric dental patients in Chile

## **Study objectives**

The current study aimed to verify the safety and effectiveness of inhalation sedation with nitrous oxide when compared to cognitive behavioral therapy.

## **Ethics approval required**

Old ethics approval format

## **Ethics approval(s)**

Comité de Bioética en Investigación en Humanos, Universidad Austral de Chile, 19/06/2018, ref: 007/2018

## **Study design**

Randomised parallel-arm trial

## **Primary study design**

Interventional

## **Study type(s)**

Treatment

## **Health condition(s) or problem(s) studied**

Anxiety associated with dental treatment

## **Interventions**

Patients who accepted the invitation were randomly allocated to the intervention A or control group B using a closed envelope technique and a block assignment of two patients. Group A received dental treatment under pharmacological sedation with a mixture of nitrous oxide in oxygen delivered by mask inhalation. Gas concentrations ranged from 30% to 70%, accordingly to patients needs. Group B received dental treatment under cognitive behavioral therapy. The cognitive behavioral therapy was a verbal 'tell-show-do' explanation of dental procedures provided before and during treatment. This is a non-pharmacological technique that allows a link to be established between the child and the dentist. All communication was constructed in phrases appropriate to the development level of the patient. Each treatment session was performed in 30-40 minutes. The total number was defined depending on their needs and according to the DMFT (decayed missing and filled teeth) caries index. All efforts were made to obtain a 1- to 2-week follow-up.

## **Intervention Type**

Mixed

## **Primary outcome(s)**

Completion of planned treatment

## **Key secondary outcome(s)**

1. Anxiety levels assessed using a facial image scale (FIS) immediately before the start of dental treatment and after the end of procedure
2. Stress assessed by measuring saliva cortisol levels immediately before the start of dental treatment and after the end of procedure

## **Completion date**

12/12/2018

## Eligibility

### Key inclusion criteria

1. Aged 5 to 12 years
2. Moderate to severe dental anxiety
3. Parents' consent to participate in sampling process and two sessions of dental treatment
4. At least one mandibular primary molar needing pulp treatment
5. Previous history of dental treatment

### Participant type(s)

Healthy volunteer

### Healthy volunteers allowed

No

### Age group

Child

### Lower age limit

5 years

### Upper age limit

12 years

### Sex

All

### Total final enrolment

49

### Key exclusion criteria

1. Chronic obstructive pulmonary diseases
2. Acute otitis media
3. Current medications
4. Chronologically immaturity that may interfere with the ability to understand verbal communication
5. Systemic or congenital disorders
6. Mental retardation

### Date of first enrolment

01/07/2018

### Date of final enrolment

01/12/2018

## Locations

### Countries of recruitment

Chile

**Study participating centre**  
**Universidad Austral de Chile**  
Rudloff 1640, Valdivia, Región de los Ríos  
Valdivia  
Chile  
5090000

## Sponsor information

**Organisation**  
Universidad Austral de Chile

**ROR**  
<https://ror.org/029ycp228>

## Funder(s)

**Funder type**  
Government

**Funder Name**  
Dirección de Investigación, Universidad Austral de Chile

**Alternative Name(s)**  
DID, UACH

**Funding Body Type**  
Government organisation

**Funding Body Subtype**  
Local government

**Location**  
Chile

## Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available upon request from Claudia Mautz, [claudiamautz@uach.cl](mailto:claudiamautz@uach.cl)

**IPD sharing plan summary**

Available on request