

# The effectiveness of new bioresorbable coronary stents in the treatment of patients with acute coronary syndromes

<b>Submission date</b> 09/05/2021	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 12/05/2021	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 13/09/2021	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Acute coronary syndromes are a range of conditions associated with suddenly reduced blood flow to the heart. They can be treated with percutaneous coronary intervention (PCI), where a tube-shaped device called a coronary stent is placed in the coronary arteries that supply blood to the heart to keep them open. There is little evidence about the effectiveness and safety of magnesium-based bioresorbable (naturally-dissolving) coronary stents in patients with acute coronary syndromes. The aim of this study is to compare the bioresorbable Magmaris stent with the drug-eluting (drug-releasing) Xience stent in patients with acute coronary syndromes undergoing PCI.

### Who can participate?

Patients with acute coronary syndromes

### What does the study involve?

Participants are randomly allocated to undergo PCI with either a bioresorbable stent or a drug-eluting stent. Participants in both groups are followed for 12 months with heart scans at 12 months.

### What are the possible benefits and risks of participating?

The study will provide information about the effects of the coronary intervention at 12 months. There is a small risk (mainly minimal bleeding) with respect to repeat invasive imaging at 12 months.

### Where is the study run from?

University Hospital Kralovske Vinohrady (Czech Republic)

### When is the study starting and how long is it expected to run for?

March 2017 to January 2021

Who is funding the study?

1. University Hospital Kralovske Vinohrady (Czech Republic)
2. Charles University in Prague (Czech Republic)

Who is the main contact?

Petr Tousek  
petr.tousek@fnkv.cz

## Contact information

### Type(s)

Public

### Contact name

Mr Petr Tousek

### ORCID ID

<https://orcid.org/0000-0002-2598-3635>

### Contact details

Srobarova 50  
Praha 10  
Czech Republic  
10034  
+420 (0)267162621  
petr.tousek@fnkv.cz

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

### Protocol serial number

EK-VP-29-0-2017

## Study information

### Scientific Title

Bioresorbable magnesium-based sirolimus-eluting stent versus permanent metallic everolimus-eluting stent in patients with acute coronary syndromes

### Acronym

PRAGUE-22

### Study objectives

The bioresorbable magnesium-based sirolimus-eluting stent has similar 12-months efficacy compared to the everolimus-eluting metallic stent in patients with acute coronary syndromes.

### Ethics approval required

Old ethics approval format

### **Ethics approval(s)**

Approved 28/06/2017, University Hospital Kralovske Vinohrady Ethics Committee (Srobarova 1150/50, 100 34 Prague, Czech Republic; +420 (0)267 16 2272; eticka.komise@fnkv.cz), ref: EK-VP-29-0-2017

### **Study design**

Two-centre investigator-initiated academic randomized study

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Acute coronary syndromes

### **Interventions**

Participants are randomised using the envelope method into two arms. Both groups undergo percutaneous coronary intervention (PCI): one arm undergoes bioresorbable stent implantation (Magmaris stent) and the other group second-generation drug-eluting stent (XIENCE) implantation. Participants in both groups are followed for 12 months with control angiography and optical coherence tomography (OCT) imaging at 12 months.

### **Intervention Type**

Device

### **Phase**

Phase IV

### **Primary outcome(s)**

Late lumen loss assessed by quantitative coronary angiography (QCA) and optical coherence tomography (OCT) at 12 months follow-up

### **Key secondary outcome(s)**

1. Device and procedural success (devices implanted by physician with optimal expansion, time) recorded at the time of the procedure
2. Clinical combined endpoints (death, stent thrombosis, target vessel myocardial infarction, clinically driven target lesion failure), measured at 12 months
3. Magmaris resorption assessed by OCT at 12 months
4. Healing state assessed by OCT at 12 months

### **Completion date**

31/01/2021

## **Eligibility**

### **Key inclusion criteria**

Patients were included if they presented with:

1. ST-elevation myocardial infarction (STEMI) <24 hours since the onset of symptoms or
2. Non-ST-elevation myocardial infarction (non-STEMI) or
3. Unstable angina caused by thrombotic acute coronary stenosis and coronary artery with stenosis diameter suitable for implantation of both investigated types of stents (vessel diameter between 2.7 and 3.7 mm)

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Total final enrolment**

50

**Key exclusion criteria**

1. Cardiogenic shock or pulmonary oedema
2. Expected survival less than 3 years due to severe comorbidities
3. Contraindication of 12 months dual antiplatelet treatment including an indication to treatment with peroral anticoagulants
4. Diffuse calcifications or extreme tortuosity of the target vessel
5. In-stent restenosis or stent thrombosis as the culprit lesion
6. Left main stenosis

**Date of first enrolment**

01/07/2017

**Date of final enrolment**

20/01/2020

**Locations**

**Countries of recruitment**

Czech Republic

**Study participating centre**

University Hospital Kralovske Vinohrady

Srobarova 50

Prague

Czech Republic

10034

**Study participating centre****AGEL**

Cardiology Department

Kyjevská 44

Pardubice

Czech Republic

53203

**Sponsor information****Organisation**

Fakultní nemocnice Královské Vinohrady

**ROR**<https://ror.org/04sg4ka71>**Funder(s)****Funder type**

University/education

**Funder Name**

Univerzita Karlova v Praze

**Alternative Name(s)**

Charles University, Charles University in Prague, Univerzita Karlova, Karls-Universität zu Prag, UK

**Funding Body Type**

Government organisation

**Funding Body Subtype**

Universities (academic only)

**Location**

Czech Republic

**Funder Name**

University Hospital Kralovske Vinohrady, Intercardis project EU Nr. CZ.02.1.01/0.0/0.0/16\_026/0008388

# Results and Publications

## Individual participant data (IPD) sharing plan

Participant level data will be available upon request to principal investigator Petr Tousek (petr.tousek@fnkv.cz). Data are already available, will be available for 5 years, and are anonymized. Participant consent was not obtained for sharing other data.

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		10/09/2021	13/09/2021	Yes	No