

# Group therapy for a healthier lifestyle in psychiatric patients

<b>Submission date</b> 18/10/2024	<b>Recruitment status</b> Recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 30/10/2024	<b>Overall study status</b> Ongoing	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
<b>Last Edited</b> 14/11/2024	<b>Condition category</b> Mental and Behavioural Disorders	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

People with severe mental illness (SMI) often have unhealthy eating habits, even though they usually understand what a healthy diet is and want to follow it. This study aims to compare the effects of a new group therapy approach with the usual treatment methods.

### Who can participate?

People between 18 and 65 years old with a diagnosis of depression, bipolar disease or psychosis with a Body-Mass-Index (BMI) over 25 and the desire to lose weight.

### What does the study involve?

The study involves two groups and participants will be assigned randomly to one of the groups. The control group involves the usual treatment (nutritional counseling and standard group therapy). The intervention group involves the usual treatment, but the standard group therapy is replaced with a new group therapy specifically designed to help patients change their dietary behavior. The new group therapy takes place once a week for a duration of 8 weeks. All patients will be tested before and after the treatment (blood sample, stool sample, dietary behavior, physical activity, psychological assessment, sociodemographic data, medical history, current medication and treatment).

### What are the possible benefits and risks of participating?

There is no additional risk for the control group. For the intervention group, there is a minimal risk associated with the psychotherapy elements of the new group therapy, which has not been tested yet.

Nutritional counseling is known to have positive effects on dietary behavior. In general, group therapy is an effective psychotherapeutic intervention. Participants will receive CHF 100 for their time spent.

### Where is the study run from?

The Psychiatric University Hospital Zurich, Switzerland.

### When is the study starting and how long is it expected to run for?

October 2024 to June 2027

Who is funding the study?

Stiftung zur Förderung von Psychiatrie und Psychotherapie (Zürich, Switzerland)  
Fonds für wissenschaftliche Zwecke im Interesse der Heilung von psychischen Krankheiten  
(Zürich, Switzerland)

Who is the main contact?

Dr med. Florian Hotzy, [florian.hotzy@pukzh.ch](mailto:florian.hotzy@pukzh.ch)  
Dr Eileen Neumann, [eileen.neumann@gmx.ch](mailto:eileen.neumann@gmx.ch)

## Contact information

### Type(s)

Public, Scientific, Principal investigator

### Contact name

Dr Eileen Neumann

### ORCID ID

<https://orcid.org/0000-0001-8543-6265>

### Contact details

Psychiatrische Universitätsklinik Zürich  
Militärstrasse 8  
Zürich  
Switzerland  
8021  
+41583847471  
[eileen.neumann@pukzh.ch](mailto:eileen.neumann@pukzh.ch)

## Additional identifiers

### Protocol serial number

181024

## Study information

### Scientific Title

Behavioral therapy to induce a healthier lifestyle and more self-efficacy in psychiatric patients (HEALTH study): A naturalistic randomized controlled trial

### Acronym

HEALTH

### Study objectives

Study participants receiving the HEALTH intervention will have better anthropometric measures (BMI and WHR) after the intervention (T1) and in the long term (T2) compared to individuals receiving TAU only.

### Ethics approval required

Ethics approval required

### **Ethics approval(s)**

submitted 13/11/2024, Kantonale Ethikkommission Zürich (Stampfenbachstrasse 121, Zürich, 8090, Switzerland; +41 43 259 79 70; info.kek@kek.zh.ch), ref: 2024-02177

### **Study design**

Monocenter open-label randomized head-to-head study with two study arms

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Psychosis, depression, bipolar disease

### **Interventions**

HEALTH Intervention:

The intervention is planned to consist of 8 group sessions in groups of 5–10 patients every 7 days.

The sessions cover the following main topics:

- Healthy diet
- Physical activity
- Circadian rhythm

Therapeutic focus is set on: behavioral therapy support to identify and change problem behavior, small goals (microsteps), and enhance self efficacy.

The control group will receive treatment as usual (TAU):

Nutritional counseling plus group sessions in a standard group therapy (e.g. art therapy or recreational contact group).

RedCap software will be used for assignment either to the intervention or control group (allocation ratio 1:1, block randomization).

### **Intervention Type**

Behavioural

### **Primary outcome(s)**

1. BMI is measured using weight and height at baseline (T0), week 9 (T1), and week 24 (T2)
2. WHR is measured using a measuring tape at baseline (T0), week 9 (T1), and week 24 (T2)

### **Key secondary outcome(s)**

1. Inflammatory markers are measured using a venous blood sample of 24 ml at baseline (T0), week 9 (T1), and week 24 (T2)
2. Gut microbiome composition is measured using a stool sample at baseline (T0), week 9 (T1), and week 24 (T2)
3. Dietary behavior is measured using the Rapid-Eating-Assessment of Participants-Short version (REAP-S), Food-Frequency-Questionnaire (FFQ), Practical Knowledge about Balanced Meals scale (PKB-7), and Salzburg Emotional Eating Scale (SEES) at baseline (T0), week 9 (T1), and week 24

(T2)

4. Physical activity is measured using the Xiaomi Smart Band 8 and a self-rating scale for physical activity at baseline (T0), week 9 (T1), and week 24 (T2)

5. Psychological assessment is measured using the Goal Attainment Scale (GAS), Beck Depression Inventory (BDI), Self-efficacy Scale (SWE), Symptom-Checklist (SCL-K-9), World Health Organization Quality of Life (WHOQOL-BREF), and Schlafqualitäts-Fragebogen (PSQI) at baseline (T0), week 9 (T1), and week 24 (T2)

**Completion date**

30/06/2027

## Eligibility

**Key inclusion criteria**

1. Age 18-65 years
2. SMI including ICD-10 F2 or F3 diagnosis
3. BMI >25 kg/m<sup>2</sup> and desire to lose weight

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Upper age limit**

65 years

**Sex**

All

**Key exclusion criteria**

1. Comorbid substance use disorder (F1) or eating disorder (F5)
2. Current medical conditions such as acute infectious disease
3. Taking medication to reduce weight (e.g. GLP-1 receptor agonists) and participating in another nutritional counselling

**Date of first enrolment**

01/01/2025

**Date of final enrolment**

31/12/2026

## Locations

## Countries of recruitment

Switzerland

## Study participating centre

Psychiatrische Universitätsklinik Zürich

Militärstrasse 8, Postfach 2019

Zürich

Switzerland

8021

## Sponsor information

### Organisation

University of Zurich

### ROR

<https://ror.org/02crff812>

## Funder(s)

### Funder type

University/education

### Funder Name

University of Zurich / Psychiatric University Clinic Zurich; Fonds für wissenschaftliche Zwecke im Interesse der Heilung von psychischen Krankheiten.

### Funder Name

Stiftung zur Förderung von Psychiatrie und Psychotherapie

## Results and Publications

### Individual participant data (IPD) sharing plan

The datasets generated during and / or analysed during the current study will be available upon request from Dr. med. Florian Hotzy ([florian.hotzy@pukzh.ch](mailto:florian.hotzy@pukzh.ch)).

### IPD sharing plan summary

Available on request

