

TURP and laser therapy for men presenting with lower urinary tract symptoms (LUTS) and chronic retention of urine

Submission date

23/01/2004

Recruitment status

No longer recruiting

Registration date

23/01/2004

Overall study status

Completed

Last Edited

14/01/2010

Condition category

Urological and Genital Diseases

☐ Prospectively registered

☐ Protocol

☐ Statistical analysis plan

☒ Results

☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Jenny Donovan

Contact details

University of Bristol

Department of Social Medicine

Canynge Hall

Whiteladies Road

Bristol

United Kingdom

BS8 2PR

+44 (0)117 928 7214

jenny.donovan@bristol.ac.uk

Additional identifiers

Study information

Scientific Title

Acronym

The CLasP Study

Study objectives

We assessed the effectiveness of laser therapy versus transurethral prostatic resection in men with symptomatic chronic urinary retention secondary to benign prostatic enlargement.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Urological and genital diseases: Lower urinary tract symptoms

Interventions

1. Laser therapy involved neodymium:YAG noncontact visual prostate ablation
2. Transurethral prostatic resection (TURP) was performed by standard electroresection

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

I-PSS quality of life score, maximum urinary flow and post-void residual urine volume.

Key secondary outcome(s)

Treatment failure, complications, hospital stay and catheterization time.

Completion date

30/05/1998

Eligibility**Key inclusion criteria**

Patients reporting moderate to severe lower urinary tract symptoms with an International Prostate Symptom Score (I-PSS) of 8 or more, benign prostatic enlargement and a persistent post-void residual urine volume of more than 300 ml.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Male

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/10/1994

Date of final enrolment

30/05/1998

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

University of Bristol

Bristol

United Kingdom

BS8 2PR

Sponsor information**Organisation**

NHS R&D Regional Programme Register - Department of Health (UK)

Funder(s)**Funder type**

Government

Funder Name

NHS Executive South West (UK)

Results and Publications

Individual participant data (IPD) sharing plan**IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/07/2000		Yes	No