

Screening and Intervention Programme for Sensible drinking - Accident and Emergency Department

Submission date 30/04/2007	Recruitment status No longer recruiting	☐ Prospectively registered
Registration date 08/06/2007	Overall study status Completed	☐ Protocol
Last Edited 26/06/2014	Condition category Mental and Behavioural Disorders	☐ Statistical analysis plan
		☒ Results
		☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Colin Drummond

Contact details

Section of Addictive Behaviour
Division of Mental Health
St George's University of London
Cranmer Terrace
London
United Kingdom
SW17 0RE

Additional identifiers

Protocol serial number

SIPS-AED

Study information

Scientific Title

A randomised controlled trial of different methods of alcohol screening and brief interventions in routine accident and emergency department care

Acronym

SIPS (AED)

Study objectives

Brief intervention by an alcohol health worker for hazardous and harmful drinkers identified by targeted screening is more effective and cost effective than brief advice conducted by Accident and Emergency Department (AED) staff in the typical AED setting.

As of 11/03/2009 this record was updated to include an extended anticipated end date; the initial end date at the time of registration was 31/12/2008. At this time, the sponsor field was also updated to include the current sponsor of Institute of Psychiatry, King's College London (UK). The initial sponsor at the time of registration was St George's University of London (UK).

Ethics approval required

Old ethics approval format

Ethics approval(s)

London Local Research Ethic Committee (LREC), 05/04/2007, ref: 07/MRE02/6

Primary study design

Interventional

Study design

Cluster prospective randomised controlled trial

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Alcohol use disorders/accident and emergency

Interventions

1. Control condition: AED staff in the control condition will be trained to apply the appropriate screening method and record the screening outcome, will feedback the result of screening to patients and offer them a Patient Information Leaflet (PIL). The PIL is from the Drink-Less Programme and has been extensively pre-tested with clinicians and patients in PHC and will be used in this trial. The PIL will also contain a number for Drinkline where the patient can access further information. The PIL to be used in this trial will be Alcohol and Sensible Drinking as in the PHC study (trial details in <http://www.controlled-trials.com/ISRCTN06145674/>).

2. Simple structured advice condition: AED staff will be trained to carry out screening and deliver up to 5 minutes of brief advice for hazardous and harmful drinkers presenting to the AED, using the Drink-Less brief intervention materials (level 1) as in the PHC study. Patients in this condition will also receive a PIL as above, including a number for Drinkline.

3. Alcohol Health Worker condition: This is based on the St Marys Hospital model. AED staff will be trained to carry out universal alcohol screening and to refer hazardous and harmful drinkers

identified by screening to an Alcohol Health Worker (AHW), by making an appointment usually the following day or as soon as possible after AED attendance. Before leaving the AED the patients in this condition will be given a PIL and Simple structured advice as above. The AHW will be experienced in carrying out alcohol assessment and brief interventions. The AHW will carry out a brief lifestyle counseling intervention lasting for 15-20 minutes as in the protocol for PHC lifestyle counseling intervention above (using the Drink-Less brief intervention materials [level 2]).

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Alcohol Use Disorders Identification Test (AUDIT) at baseline and 6 months.

Key secondary outcome(s)

The following will be assessed at baseline and 6 months:

1. Alcohol Problems Questionnaire
2. EQ-5D questionnaire (quality of life assessment)
3. Service utilisation
4. Staff attitudinal and organisational measures

Completion date

30/06/2010

Eligibility**Key inclusion criteria**

1. AED patients
2. Scoring positive on the Paddington Alcohol Test (PAT), FAST or SASQ
3. Patients who are alert and orientated
4. Aged 18 or over, either sex
5. Able to speak English sufficiently well to complete study questionnaires

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

1. Already in contact with alcohol services
2. Those already included in the study
3. Those requesting help with alcohol problems at a tertiary level

Date of first enrolment

01/04/2007

Date of final enrolment

30/06/2010

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre**Section of Addictive Behaviour**

London

United Kingdom

SW17 0RE

Sponsor information**Organisation**

Institute of Psychiatry, Kings College London (UK)

ROR

<https://ror.org/0220mzb33>

Funder(s)**Funder type**

Government

Funder Name

Department of Health (UK) - part of an action under the governments Alcohol Harm Reduction Strategy for England (2004)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	25/06/2014		Yes	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes