

The use of sling exercise therapy for weakness and functional disability on one side of the body after a stroke

Submission date 13/05/2022	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 27/05/2022	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 26/05/2022	Condition category Circulatory System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Stroke patients have problems with trunk control, which is linked with restricted balance, an increased risk of falls, poor walking performance, and dependence on people in daily activities. The goal of physical therapy is to improve the trunk stability of stroke patients. In addition, recent studies have found that trunk exercises performed on an unstable surface are more effective than those performed on a stable surface. Sling exercise therapy (SET), one of the common unstable surfaces, can help patients perform the training program more effectively by supporting their body weight with an elastic cord. The aim of this study is to determine the effects of core stability exercises using sling exercise training on balance, walking, and activities of daily living for patients with subacute hemiparesis (weakness of one side of the body).

Who can participate?

Patients diagnosed with their first stroke with hemiplegia within the last 6 months who can walk independently

What does the study involve?

Participants are randomly allocated to the sling exercise training or conventional physical therapy. Participants receive the training for 8 weeks and their trunk function, balance, walking and daily living activities are assessed before and after 8 weeks of training.

What are the possible benefits and risks of participating?

Balance, walking and daily living activities may be improved after the training, and trunk function may be improved after the sling exercise intervention.

Where is the study run from?

Yuan's General Hospital (Taiwan)

When is the study starting and how long is it expected to run for?

October 2017 to March 2018

Who is funding the study?

1. Ministry of Science and Technology (Taiwan)
2. National Sun Yat-sen University (Taiwan)
3. Kaohsiung Medical University (Taiwan)

Who is the main contact?

Miss Shih-Chi Tseng
origin0725@gmail.com

Contact information

Type(s)

Scientific

Contact name

Miss Shih Chi Tseng

Contact details

No.100
Shih-Chuan 1st Road
Sanmin Dist
Kaohsiung
Taiwan
807
+886 (0)7 312 1101#2646#624
origin0725@imst.nsysu.edu.tw

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

The effect of core stability exercises using sling exercise therapy on trunk function, balance ability, walking performance and daily living activities in subacute hemiparesis patients

Study objectives

The sling exercise training improved balance ability, walking performance and activities of daily living significantly compared to baseline measurement and the conventional therapy intervention.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 25/12/2017, Institutional Review Board of Yuan's General Hospital (No. 162, Chenggong 1st Road, Lingya Dist., Kaohsiung City, Taiwan (R.O.C.); +886 (0)7 3350205; irb@yuanhosp.com.tw), ref: 20171031B

Study design

Single-center interventional randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Subacute stroke

Interventions

Participants were randomly assigned to study groups by sealed envelopes to receive sling exercise training for the experiment group and conventional physical therapy for the control group. All participants receive a total of 20 intervention sessions, 3 intervention sessions a week and 1 hour per session, over 8 weeks.

Intervention Type

Other

Primary outcome(s)

1. Balance ability measured by Timed Up & Go, Berg Balance scale and postural assessment scale at baseline and post-intervention (8 weeks)
2. Walking performance measured by 5-meter walking test and 6-minute walking test at baseline and post-intervention (8 weeks)

Key secondary outcome(s)

1. Trunk function measured by the Trunk Impairment Scale and the pelvic backward tilt force by using stabilizer pressure biofeedback at baseline and post-intervention (8 weeks)
2. Daily living activities measured by the Barthel index and the activities-specific balance confidence scale at the baseline and post-intervention (8 weeks)

Completion date

12/03/2018

Eligibility

Key inclusion criteria

1. Less than 6 months since diagnosis of first stroke with hemiplegia
2. Modified Rankin Scale scores less than or equal to 4 (MRS \leq 4)
3. Independent walking

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

40

Key exclusion criteria

1. Trunk Impairment Scale (TIS) score of 17 or more
2. Other musculoskeletal or neurologic problems

Date of first enrolment

25/12/2017

Date of final enrolment

15/01/2018

Locations**Countries of recruitment**

Taiwan

Study participating centre

Yuan's General Hospital

No. 162, Chenggong 1st Rd., Lingya Dist.

Kaohsiung

Taiwan

802

Sponsor information**Organisation**

Kaohsiung Medical University

ROR

<https://ror.org/03gk81f96>

Funder(s)

Funder type

Government

Funder Name

Ministry of Science and Technology, Taiwan

Alternative Name(s)

Ministry of Science and Technology, R.O.C. (Taiwan), Ministry of Science and Technology of Taiwan, MOST

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Taiwan

Funder Name

National Sun Yat-sen University

Alternative Name(s)

NSYSU

Funding Body Type

Government organisation

Funding Body Subtype

Universities (academic only)

Location

Taiwan

Funder Name

Kaohsiung Medical University

Alternative Name(s)

-, Kaohsiung Medical University | Kaohsiung Medical University, Kaohsiung Medical University in Taiwan, Kaohsiung Medical University (KMU) - Taiwan, Kaohsiung Medical University, Taiwan, Kaohsiung Medical University | , Kaohsiung Medical College, KMU

Funding Body Type

Private sector organisation

Funding Body Subtype

Universities (academic only)

Location

Taiwan

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study will be included in the subsequent results publication

IPD sharing plan summary

Published as a supplement to the results publication