

Back pain and Rescuer fatigue following CardioPulmonary Resuscitation by kneeling, standing on taboret, and standing postures

Submission date 27/11/2007	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 18/12/2007	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 08/04/2021	Condition category Signs and Symptoms	☐ Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Foo Ning Ping

Contact details
201, Taikang Village
Liou Ying township
Tainan
Taiwan
736

Additional identifiers

Protocol serial number
CLFHR9611

Study information

Scientific Title
Back pain and Rescuer fatigue following CardioPulmonary Resuscitation by different postures

Acronym

BRCPR

Study objectives

Back pain as an occupational disorder in health care providers has been well documented in the literature. But the association between back pain and Cardiopulmonary Resuscitation (CPR) was seldom studied. The objectives of this study were to determine whether different CPR postures including kneeling, standing, and standing on taboret are able to induce back pain in rescuers, and to compare differences in rescuer fatigue in three CPR postures.

Hypothesis:

Back pain and rescuer fatigue are different among three different postures of CPR.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from the Institutional Review Board of Chi Mei Medical Center from 5th February 2007 to 16th February 2008 (ref: IRB09602-0011).

Primary study design

Interventional

Study design

Randomised, controlled, cross-over trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Rescuer fatigue for resuscitation

Interventions

Cardiopulmonary resuscitation for 10 minutes by using kneeling, standing, and standing on taboret postures.

Since this is a prospective and cross-over trial, each of the participants should complete three different CPR posture within 3 weeks in one week apart. Therefore, each of the sequences of CPR should follow up for at least 48 hours, but the total follow-up period for each of them will be 3 weeks and two days.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

1. Severity of back pain (by Brief Pain Inventory Short Form), measured 24 hours and 48 hours post CPR
2. Rescuer fatigue (effective compression of cardiac massage) - detected by Laerdal Resusci-Anne® Skillreporter™ mannequin, measured during 10 minutes of cardiac massage

Key secondary outcome(s)

Range of motion of elbow and back with different CPR postures (detected by flexible goniometer), measured during 10 minutes of cardiac massage.

Completion date

31/12/2007

Eligibility**Key inclusion criteria**

Health care providers working at the Emergency Department that have:

1. Clinical work experience longer than 2 years
2. Experience of performing CPR for more than 20 times

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

24

Key exclusion criteria

Candidates with:

1. A herniated intervertebral disc
2. Previous spine surgery
3. Underlying ankylosing spondylitis or other autoimmune diseases
4. Current pregnancy

Date of first enrolment

01/03/2007

Date of final enrolment

31/12/2007

Locations**Countries of recruitment**

Taiwan

Study participating centre
201, Taikang Village
Tainan
Taiwan
736

Sponsor information

Organisation

Chi Mei Foundation Medical Center (Taiwan) - Liou Ying Campus

ROR

<https://ror.org/02y2htg06>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Chi Mei Foundation Medical Center (Taiwan) - Liou Ying Campus

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/05/2010	08/04/2021	Yes	No