

# A study to evaluate the safety and efficacy of PTC518 in participants with Huntington's disease (HD)

<b>Submission date</b> 03/03/2022	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 08/08/2022	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 02/02/2026	<b>Condition category</b> Genetic Diseases	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

The safety and efficacy of three doses of PTC518 (investigational medicine), for participants with Huntington's Disease (HD), will be studied in this Clinical Trial.

PTC518 is being studied to determine if it can slow the progression of HD and its associated symptoms by reducing the production of an abnormal version of a protein called huntingtin protein (sometimes called a mutant version or mHTT).

### Who can participate?

Approximately 252 participants, aged 25 years and older will participate across approximately 20 sites in North America, Europe, and Australia.

### What does the study involve?

The study is divided into the below periods:

- Screening period (to check participant eligibility to join the study)
- Baseline visit (to further confirm participant eligibility and initial dosing)
- Treatment period
- Safety follow-up period (to check participants' overall health and safety)

The study medicine comes in tablet form and is to be taken orally, in the morning, at least 2 hours before the first meal of the day (fasting).

Participation in the study will last up to 18 months, from screening to the final follow-up visit. The study treatment period will last approximately 12 months. Participation is voluntary. Participants will be expected to attend clinic visit whereby study assessments will take place e. g., blood and urine collection, vital signs monitoring, assessments of motor function & gait, cognition & behavioral and mental health, and completion of questionnaires.

### What are the possible benefits and risks of participating?

Not provided at time of registration

Where is the study run from?  
PTC Therapeutics International Limited (Ireland)

When is the study starting and how long is it expected to run for?  
September 2021 to July 2025

Who is funding the study?  
PTC Therapeutics, Inc. (USA)

Who is the main contact?  
Dr Sarah Tabrizi, s.tabrizi@ucl.ac.uk

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Patient Advocacy

**Contact details**  
PTC Therapeutics, Inc.  
100 Corporate Court  
South Plainfield  
United States of America  
07080  
+1 908 2227000  
medinfo@ptcbio.com

**Type(s)**  
Principal investigator

**Contact name**  
Dr Sarah Tabrizi

**Contact details**  
University College London  
National Hospital for Neurology and Neurosurgery  
Queen Square  
London  
United Kingdom  
WC1N 3BG  
-  
s.tabrizi@ucl.ac.uk

## Additional identifiers

Clinical Trials Information System (CTIS)  
2021-003852-18

## **Integrated Research Application System (IRAS)**

1004960

## **ClinicalTrials.gov (NCT)**

NCT05358717

## **Protocol serial number**

PTC518-CNS-002-HD

## **Central Portfolio Management System (CPMS)**

51614

# **Study information**

## **Scientific Title**

A Phase 2a, randomized, placebo-controlled, dose-ranging study to evaluate the safety and efficacy of PTC518 in subjects with Huntington's disease

## **Acronym**

PIVOT HD

## **Study objectives**

Current study hypothesis as of 25/07/2023:

Primary objective:

- Evaluate the safety of PTC518 compared with placebo in participants with Huntington's Disease (HD)
- Evaluate the pharmacodynamic (PD) effects of PTC518 through the reduction in blood total huntingtin (tHTT) protein levels

Secondary Objectives:

- Assess the effects of PTC518 on change in caudate volume via volumetric magnetic resonance imaging (vMRI) (key secondary)
- Assess the effects of PTC518 on change in composite Unified Huntington's Disease Rating Scale (cUHDRS)
- Determine the effect of PTC518 on mutant huntingtin (mHTT) protein in cerebrospinal fluid (CSF) at Month 12
- Determine the effect of PTC518 on blood mHTT levels at Month 12

Previous study hypothesis:

Primary objective:

- Evaluate the safety and pharmacodynamic effects of 3 different doses of PTC518 and placebo in subjects with Huntington's Disease (HD)

Secondary Objectives:

- Determine the effect of PTC518 on huntingtin (HTT) mRNA in blood and mHTT protein in CSF
- Reduction in blood mHTT levels

Exploratory Objectives:

- Assess the effect of PTC518 on change in whole brain, caudate, and putamen volume via

volumetric magnetic resonance imaging (vMRI)

- Assess the effect of PTC518 on change in ventricular volume via vMRI
- Assess change after 12 weeks of treatment in clinical scales

Pharmacokinetic Objective:

- Evaluate the concentration of PTC518 in subjects with HD

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Approved 03/05/2022, London Research Ethics Committee (80 London Road, Skipton House, London, SE1 6LH, UK; +44 (0)2071048387; londonbridge.rec@hra.nhs.uk), ref: 22/LO/0229

### **Study design**

Interventional double-blind randomized parallel group placebo-controlled trial

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Huntington's disease is an illness caused by a faulty gene in the DNA. Huntington's affects the body's nervous system – which can cause changes in movement, learning, thinking and emotions.

### **Interventions**

Current interventions as of 25/07/2023:

Participants will first be randomized to Part A or Part B or Parts D or E in a 1:1 randomization ratio, depending on their Huntington's disease Integrated Staging System (HD-ISS) staging criteria and then to active treatment (PTC518 5 mg in Parts A and D and 10 mg in Parts B and E) or matching placebo within each part in a 2:1 ratio of active treatment to placebo. A Drug Safety Monitoring Board (DSMB) will undertake an unblinded review of safety data from the 5 and 10 mg dosing groups and provide a recommendation on when Parts C and F (with a 20 mg active treatment arm) can be initiated. At that time, participants will be randomized to any study Part that is currently open for enrollment, and then to either active treatment or placebo (in a 2:1 ratio) within that Part. The participant will receive the assigned treatment for approximately 12 months. After the treatment period ends, there is a safety follow-up period for all arms with follow-up safety visits at Month 13, Month 16, and Month 18.

Previous interventions:

There are three treatment arms: 1) Part A (5 mg or placebo), 2) Part B (10 mg or placebo), and 3) Part C (20 mg or placebo). Participants will be assigned to either Part A, Part B, or Part C based on which parts of the study are open at the time of consent. A computer program (an interactive response technology system) will do the part assignment and assign the participant to either the study drug or placebo. The participant will have a 2 in 3 chance of receiving the study drug versus placebo. The subject will receive the assigned treatment for approximately 85 days or 12 weeks. After the treatment period ends, there is a safety follow-up period for all arms. The safety follow-up period includes two visits on Day 113 (approximately one month after the treatment period ends) and Day 185 (approximately 4 months after the treatment period ends.)

## **Intervention Type**

Drug

## **Phase**

Phase II

## **Drug/device/biological/vaccine name(s)**

PTC518

## **Primary outcome(s)**

Current primary outcome measures as of 25/07/2023:

1. Number of participants with adverse events (AEs) [time frame: baseline up to month 18]
2. Change from baseline in blood total huntingtin protein (tHTT) at Month 3 [time frame: baseline, month 3]

Previous primary outcome measures:

Primary Safety Endpoints:

1. Safety profile as characterized by TEAEs, laboratory abnormalities, NFL levels in plasma and CSF, YKL-40 levels in CSF, ECG, vital signs, slit lamp eye examination, and physical examination

Primary Efficacy Endpoint:

2. Change from Baseline in blood tHTT protein at Day 85

## **Key secondary outcome(s)**

Current secondary outcome measures as of 25/07/2023:

1. Change from baseline in caudate volume as assessed via vMRI at month 12 [time frame: baseline, month 12]
2. Change from baseline in Composite Unified Huntington's Disease Rating Scale (cUHDRS) Scores at month 12 [time frame: baseline, month 12]
3. Change from baseline in blood tHTT protein at month 12 [time frame: baseline, month 12]
4. Change from baseline in CSF mHTT at month 12 [time frame: baseline, month 12]
5. Change from baseline in blood mHTT protein at month 12 [time frame: baseline, month 12]

Previous secondary outcome measures:

1. Change from Baseline in blood HTT mRNA at Days 29, 57, and 85
2. Change from Baseline in CSF mHTT at Day 85
3. Change from Baseline in blood mHTT protein at Day 85

Exploratory Endpoints

4. Change from Baseline in whole brain, caudate, putamen, and ventricular volume (as assessed by vMRI) at Day 85
5. Change from Baseline in UHDRS scores, with the exception of Behavioral Examination, at Day 85
6. Change from Baseline in the short form of the Problem Behaviors Assessment (PBA-s) at Day 85 (substituting for the UHDRS Behavioral Examination)
7. Change from Baseline in wearable accelerometer assessment of Timed Up and Go (TUG), 2-minute walk distance, and postural sway at Day 85
8. Change from Baseline in the FuRST 2.0 questionnaire at Day 85

## Pharmacokinetic Endpoints

9. Plasma trough concentration (C<sub>trough</sub>) and accumulation ratio of PTC518 in plasma at Visits 3, 4, and 5 and accumulation ratio of PTC518 in CSF at Visit 5

## Completion date

31/07/2025

## Eligibility

### Key inclusion criteria

Current inclusion criteria as of 25/07/2023:

Key Inclusion Criteria:

1. Genetically confirmed HD diagnosis with a cytosine-adenine-guanine (CAG) repeat length from 40 to 50, inclusive

Eligibility for HD-ISS Stage 2 Group (Parts A, B, and C):

1. A Unified Huntington's Disease Rating Scale (UHDRS)-Independence Scale (IS) score of 100
2. A UHDRS Total Functional Capacity (TFC) score of 13
3. A score between 0.18 and 4.93 inclusive on the normed version of the HD prognostic index (PINHD)

Eligibility for HD-ISS Mild Stage 3 Group (Parts D, E, and F):

1. A UHDRS Total Functional Capacity (TFC) score of 11 or 12, or a UHDRS TFC score of 13 with an UHDRS IS score of <100

Note: Other inclusion criteria may apply.

Previous inclusion criteria:

1. Ambulatory male or female patients aged 25 years and older, inclusive
2. Subject (or legally authorized representative) is willing and able to provide informed consent and comply with all protocol requirements
3. Genetically confirmed HD diagnosis with a CAG repeat length from 42 to 50, inclusive
4. A UHDRS-IS score of 100
5. A UHDRS TFC score of 13
6. A score between 0.18 and 4.93 inclusive on the normed version of the HD prognostic index (PINHD)
7. Women of childbearing potential (WOCBP) must agree to use highly effective methods of contraception during dosing and for 6 months after stopping the study medication.
8. Sexually active and fertile males must agree to use a condom during intercourse while taking the study drug and for 6 months after stopping the study drug and should neither father a child nor donate sperm in this period. A condom is required to be used also by vasectomized men in order to prevent the potential delivery of the drug via seminal fluid.

### Participant type(s)

Patient

### Healthy volunteers allowed

No

### Age group

Mixed

**Lower age limit**

25 years

**Upper age limit**

70 years

**Sex**

All

**Total final enrolment**

159

**Key exclusion criteria**

Current exclusion criteria as of 25/07/2023:

Key Exclusion Criteria:

1. Receipt of an experimental agent within 90 days or 5 half-lives prior to Screening or anytime over the duration of this study, including ribonucleic acid (RNA)- or deoxyribonucleic acid (DNA)-targeted HD-specific investigational agents such as antisense oligonucleotides, cell transplantation, or any other experimental brain surgery
2. Any history of gene therapy exposure for the treatment of HD
3. Participation in an investigational study or investigational paradigm (such as exercise/physical activity, cognitive therapy, brain stimulation, etc) within 90 days prior to Screening or anytime over the duration of this study
4. Any medical history of brain or spinal disease that would interfere with the lumbar puncture process safety assessments
5. Any medical history or condition that would interfere with the ability to complete the protocol-specified assessments (for example, implanted shunt, conditions precluding magnetic resonance imaging [MRI] scans)
6. Pregnancy, planning on becoming pregnant during the course of the study or within 6 months of the end of treatment, or currently breastfeeding

Note: Other exclusion criteria may apply.

Previous exclusion criteria:

1. Inability or unwillingness to swallow oral tablets
2. Receipt of an experimental agent within 90 days or 5 half-lives prior to Screening or anytime over the duration of this study, RNA- or DNA-targeted HD-specific investigational agents such as antisense oligonucleotides, cell transplantation, or any other experimental brain surgery
3. Any history of gene therapy exposure for the treatment of HD
4. Participation in an investigational study or investigational paradigm (such as exercise/physical activity, cognitive therapy, brain stimulation, etc) within 90 days prior to Screening or anytime over the duration of this study
5. Presence of an implanted deep brain stimulation device
6. Family history of early onset cataracts or presence of cataracts at Baseline
7. Brain and spinal pathology that may interfere with CSF homeostasis and circulation, increased intracranial pressure (including presence of a shunt for the drainage of CSF or an implanted central nervous system catheter), malformations, and/or tumors
8. Hospitalization for any major medical or surgical procedure involving general anesthesia within 12 weeks of Screening or planned during the study

9. At significant risk of suicide as measured by the C-SSRS with a moderate risk rating or higher score
10. Risk of a major depressive episode, psychosis, confusional state, or violent behavior as assessed by the investigator
11. Any medical history of brain or spinal disease that would interfere with the lumbar puncture processor safety assessments
12. History of malignancy of any organ system (other than localized basal cell carcinoma of the skin or in situ cervical cancer), treated or untreated, within the past 5 years, regardless of whether there is evidence of local recurrence or metastases
13. Any medical history or condition that would interfere with the ability to complete the protocol-specified assessments (eg, implanted shunt, conditions precluding MRI scans)
14. Antidepressant or benzodiazepine use, unless receiving a stable dose for at least 6 weeks prior to Screening and with a dose regimen that is not anticipated to change during the study
15. History of illicit/illegal drug use, or alcohol use in the high-risk category of risk drinking levels according to the World Health Organization for a duration of 1 month or longer that in the opinion of the investigator could compromise the interpretability of study results
16. Clinically significant medical condition, which in the opinion of the investigator could adversely affect the safety of the subject or impair the assessment of study results (e.g. inability to fast)
17. Current significant renal impairment defined as estimated glomerular filtration rate  $<60$  mL/min/1.73 m<sup>2</sup> at Screening
18. Current hepatic impairment resulting in elevated liver function tests (aspartate transaminase [AST], alanine transaminase [ALT], alanine phosphatase [ALP]) at 3 times the upper limit of normal at Screening
19. Pregnancy, planning on becoming pregnant during the course of the study or within 6 months of end of treatment, or currently breastfeeding
20. Use of medications that are moderate or strong inhibitors of cytochrome P450 (CYP) 3A4 within 1 week of Screening or medications that are moderate or strong inducers of CYP3A4 within 2 weeks of Screening or planned use of moderate or strong CYP3A4 inhibitor or inducer medications during the study period.

**Date of first enrolment**

03/06/2022

**Date of final enrolment**

31/12/2023

## **Locations**

**Countries of recruitment**

United Kingdom

England

Wales

Australia

France

Germany

Netherlands

United States of America

**Study participating centre**

**UCL Queen Square Institute of Neurology National Hospital for Neurology and Neurosurgery**  
Box 104  
Queen Square  
London  
England  
WC1N 3BG

**Study participating centre**

**University Hospital of Wales**  
Heath Park  
Cardiff  
England  
CF14 4XW

**Study participating centre**

**University Hospitals Birmingham NHS Foundation Trust NIHR/Wellcome Trust Clinical Research Facility**  
First Floor, Blue Zone  
Heritage Building  
Queen Elizabeth Hospital  
Edgbaston  
Birmingham  
England  
B15 2TH

**Study participating centre**

**Manchester University NHS Foundation Trust**  
6th Floor  
St Marys Hospital  
Oxford Road  
Manchester  
England  
M13 9WL

**Study participating centre**

**Leiden University Medical Center**

Leiden  
Netherlands  
2333 ZA

**Study participating centre**

**George-Huntington-Institut**

Münster  
Germany  
48149

**Study participating centre**

**Ruhr-Univ.Bochum St. Joseph-Hospital**

Bochum  
Germany  
44791

**Study participating centre**

**Ulm University, UKU, Dep. of Neurology**

Ulm  
Germany  
89081

**Study participating centre**

**Centre Hospitalier Universitaire d'Angers**

Angers  
France  
49100

**Study participating centre**

**Hôpital Universitaire de Marseille Hôpital de la Timone**

Marseille  
France  
13385

**Study participating centre**

**Brain and Spine Institute Paris**

Paris

France

75013

**Study participating centre****Westmead Hospital**

Sydney

Australia

2145

**Study participating centre****Monash Health**

Clayton

Australia

3168

**Sponsor information****Organisation**

PTC Therapeutics International Limited

**Funder(s)****Funder type**

Industry

**Funder Name**

PTC Therapeutics

**Alternative Name(s)**

PTC Therapeutics Inc., PTC Therapeutics, Inc., PTC Therapeutics Incorporated, PTC Therapeutics, Inc, PTC

**Funding Body Type**

Private sector organisation

**Funding Body Subtype**

For-profit companies (industry)

## Location

United States of America

# Results and Publications

## Individual participant data (IPD) sharing plan

Study participants will provide appropriate consent to enable the sharing of study data. Through the consent form, participants will be informed of the possibility of their data being shared with other research teams and research collaborators in other countries.

All third parties involved with the research will have appropriate safeguards and confidentiality agreements in place to assure that same level of confidentiality as within the UK will be maintained.

## IPD sharing plan summary

Other

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">HRA research summary</a>			28/06/2023	No	No