

Finding the Optimal Resistance Training Exercise for type 2 diabetes

Submission date
14/01/2010

Recruitment status
No longer recruiting

☐ Prospectively registered

☐ Protocol

Registration date
21/06/2010

Overall study status
Completed

☐ Statistical analysis plan

☐ Results

Last Edited
21/06/2010

Condition category
Nutritional, Metabolic, Endocrine

☐ Individual participant data

☐ Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Paul Oh

Contact details
Toronto Rehabilitation Institute
347 Rumsey Rd
Toronto
Canada
M4G 1R7

Additional identifiers

Protocol serial number
#19581

Study information

Scientific Title
Finding the optimal volume and intensity of resistance training exercise for type 2 diabetes: a randomised controlled trial

Acronym
The FORTE Study

Study objectives

A higher intensity, hypertrophic resistance training protocol added to aerobic training would lead to more marked improvements to glycaemic control and cardiovascular risk factors compared to lower intensity endurance training.

Ethics approval required

Old ethics approval format

Ethics approval(s)

University of Toronto Research Ethics Board (REB) approved in February 2006 (ref: 19581)

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Type 2 diabetes

Interventions

Six months of aerobic and resistance training:

1. Aerobic training and minimal resistance training (usual care)
2. Aerobic training and high intensity resistance training
3. Aerobic training and low intensity resistance training

The intervention for all groups was 6 months. Following completion of the program, patients will not be followed-up.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Measured at baseline and 6 months:

1. HbA1c
2. Maximal aerobic power (VO2 max)

Key secondary outcome(s)

Measured at baseline and 6 months:

1. Fasting glucose
2. Insulin
3. Proinsulin
4. Blood lipid profile
5. Body composition (dual energy x-ray absorptiometry [DEXA])

6. Cytokines (interleukin-6 [IL-6], interleukin-1B [IL-1B], c-reactive protein [CRP])
7. Adipokines (tumour necrotising factor-alpha [TNF-alpha], adiponectin, leptin, leptin receptor)
8. Depressive mood (Center for Epidemiologic Studies Depression Scale [CESD])
9. Quality of life (Diabetes Quality of Life Clinical Trial Questionnaire [DQLQ])
10. Nutrition intake

Completion date

31/03/2009

Eligibility

Key inclusion criteria

1. Adults aged greater than 18 years, either sex
2. Type 2 Diabetes
3. Participating in the Diabetes Exercise and Healthy Lifestyle Program of the Toronto Rehabilitation Institute

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

1. History of diagnosed cardiovascular disease
2. Contraindications to high intensity exercise participation:
 - 2.1. Glaucoma
 - 2.2. Retinopathy
 - 2.3. Musculoskeletal limitations
 - 2.4. Kidney disease
 - 2.5. Balance impairments

Date of first enrolment

01/05/2006

Date of final enrolment

31/03/2009

Locations

Countries of recruitment

Canada

Study participating centre
Toronto Rehabilitation Institute
Toronto
Canada
M4G 1R7

Sponsor information

Organisation
Toronto Rehabilitation Institute (Canada)

ROR
<https://ror.org/00mxe0976>

Funder(s)

Funder type
Research organisation

Funder Name
Toronto Rehabilitation Institute (Canada)

Alternative Name(s)
Toronto Rehab, UHN Toronto Rehabilitation Institute, TRI

Funding Body Type
Private sector organisation

Funding Body Subtype
Other non-profit organizations

Location
Canada

Funder Name
Human Physiology Performance Laboratory (Canada)

Funder Name

Canadian Diabetes Association (Canada)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration