

Development of a program to monitor patients who were started on HIV treatment on the same day of HIV diagnosis

Submission date 22/01/2025	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 03/04/2025	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 03/04/2025	Condition category Infections and Infestations	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

The South African government wants to improve HIV treatment and achieve global targets. One way to do this is by starting patients on antiretroviral therapy (ART) on the same day they're diagnosed with HIV. This approach is called same-day initiation (SDI). Research shows that starting ART early reduces illness, death, and the spread of HIV. It also boosts patients' hope and optimism. However, putting SDI into practice is challenging due to various obstacles. Some of the challenges include structural barriers such as healthcare system issues, like laboratory result delays; attitudes such as patients' and healthcare providers' attitudes toward SDI; and additional challenges in developing countries such as substance abuse, housing instability, and managing clinical symptoms. Although South Africa has adopted SDI, there are concerns about whether patients stay in treatment and how effective the treatment is. This study aims to create a comprehensive plan (standard operating procedure) to:

1. Monitor how well SDI is implemented.
2. Measure the outcomes of treatment.

By doing so, the study hopes to improve the effectiveness of SDI in South Africa.

Who can participate?

All newly diagnosed people aged 12 years and over living with HIV

What does the study involve?

Participants are randomly allocated to one of two groups, namely the intervention group and the control group. The intervention group would receive the interventions aimed at improving treatment outcomes, while the control group would use the normal protocol which does not involve new interventions. Lastly, this study involves the collection of blood, home visits, and social worker involvement among patients in the intervention group.

What are the possible benefits and risks of participating?

The participants may benefit directly from this study due to improved treatment outcomes

(adherence, viral load suppression and prevention of HIV-related deaths). Secondly, patients have an opportunity to see a social worker, assigned to a treatment advisor, and have their treatment delivered to their homes. There are no risks in participating in this study.

Where is the study run from?

This study was run in the O.R. Tambo district, Eastern Cape, South Africa. The study was conducted at three healthcare clinics, namely, Mthatha Gateway Clinic, Tsolo Clinic, and Flagstaff Clinic.

When is the study starting and how long is it expected to run for?

The study was conducted (begin data collection) from 09 January 2023 to 16 August 2023. This study was expected and ran for six months. However, the study plan was proposed on 06/08 /2021.

Who is funding the study?

This study was supported by the Chemical Industries Education and Training Authority (CHIETA) and the Strategic Health Innovation Partnerships (SHIP) of the Medical Research Council (MRC), funding attributed to Prof. Eugene Jamot Ndebia.

Who is the main contact?

For the study: Siyakudumisa Nontamo, siyakudumisanontamo86@gmail.com and 206607687@mywsu.ac.za

For funding: Professor Eugen Ndebia endebia@wsu.ac.za

Contact information

Type(s)

Public, Scientific, Principal investigator

Contact name

Dr Siyakudumisa Nontamo

ORCID ID

<https://orcid.org/0009-0001-6929-0275>

Contact details

03 Swift Street
Southernwood
Mthatha
South Africa
5099
+27 (0)712341138
206607687@mywsu.ac.za

Type(s)

Scientific

Contact name

Prof Eugene Ndebia

ORCID ID

<https://orcid.org/0000-0002-5840-0715>

Contact details

Private Bag X1
Walter Sisulu University
Mthatha
South Africa
5099
+27 (0)767550574
siyakudumisanontamo86@gmail.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

Developing a Standard Operating Procedure for monitoring same-day initiation of antiretroviral therapy among newly diagnosed persons living with HIV in South Africa

Study objectives

By and large, when carefully measured and monitored, same-day antiretroviral therapy (ART) initiation improves treatment adherence, retention to care, and viral load suppression. A procedure to measure and monitor same-day ART initiation will be associated with positive treatment outcomes. As a result, if developed and implemented, positive treatment outcomes will flourish even further in South Africa. Therefore, the HIV care continuum will be optimized, and the 959595 UNAIDS strategy will be achieved by 2024. Subsequently, the Millennium Development Goal (MDG) that aims to achieve an ambition of 'free HIV generation' by 2030 is to be achieved, and life expectancy will further improve.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 28/09/2022, Walter Sisulu University Faculty of Health Sciences Research Ethics and Biosafety Committee (Walter Sisulu University, NMD Campus, Private Bag X1, Mthatha, 5099, South Africa; +27 (0)475022100; fhs@wsu.ac.za), ref: 027/2022

Study design

Multicenter interventional unblinded randomized controlled trials

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Improving adherence to HIV treatment among newly HIV-diagnosed patients

Interventions

Following the patients' written informed permission, the researcher conducted a screening evaluation to assess the study exclusion criteria. Eligible participants were then enrolled and randomly assigned on the same day of HIV testing. A computer-generated random-number list was used to randomly assign participants to either standard ART or same-day ART initiation in a 1:1 ratio, with allocation concealment. Participants were enrolled and divided into study groups by the researcher. However, participants, site personnel, and the study supervisor were not blinded to the group assignments.

After randomization:

1. Standard group: Received ART initiation as per national guidelines, with follow-up on Day 28
2. Intervention group: Received additional support, including:
 - 2.1. Assignment to patient navigator after starting treatment
 - 2.2. Day 5: Nurse support (blood result interpretation, side effect management, counseling)
 - 2.3. Day 14: Social worker support (counseling, disclosure support, needs assessment, opportunistic infection evaluation)

The duration of the interventions was 6 months from the date the participant was enrolled in the study.

Intervention Type

Mixed

Primary outcome(s)

Retention in care is defined as maintaining an HIV-1 RNA level <50 copies/ml 6 months following HIV testing. This outcome was measured at baseline and month 6 using the whole blood that would measure HIV-1 RNA level (viral load)

Key secondary outcome(s)

1. Retention in care with HIV-1 RNA <1,000 copies/mL at 6 months post-testing, which was also measured from baseline and at month 6
2. Adherence is measured using pharmacy refill records at months 1, 2, 3, 4, 5, and 6
3. Loss to follow-up measured using patient charts and phone calls at 6 months into treatment
4. Mortality rate was measured using phone calls to verify from the family members and the clinic file at months 1, 2, 3, 4, 5, and 6

Completion date

16/08/2023

Eligibility

Key inclusion criteria

1. All patients above 18 years are eligible for ART on the same day as HIV testing services (HTS) (same-day initiation)
2. Patients whose status is known but who have never been initiated on ART
3. Adolescents from 12 years after the completion of the HIV disclosure process in

accompaniment of the parent or caregiver

4. Stable mental health patients deemed fit to be initiated on ART

All patients whose permanent and temporary addresses are known and reachable for the duration of the study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

12 years

Upper age limit

80 years

Sex

All

Total final enrolment

142

Key exclusion criteria

1. Symptomatic client who does not meet the criteria of same-day ART initiation
2. Active mental health condition
3. Clients below the age of 12 years
4. Antenatal clients enrolled in the prevention of mother-to-child transmission program
5. Known HIV-positive client who was once initiated on treatment and defaulted
6. TB co-infected client
7. A newly HIV-positive person who is likely to seek a transfer to unreachable locations during the study period.

Date of first enrolment

03/01/2023

Date of final enrolment

17/02/2023

Locations

Countries of recruitment

South Africa

Study participating centre

Tsolo Clinic

Dr Malizo Mpehle Hospital Road, main road, Tsolo, 5170
Mthatha
South Africa
5099

Study participating centre**Flagstaff Clinic**

76 Main Street
Flagstaff
South Africa
4810

Study participating centre**Mthatha Gateway Clinic**

Mthatha General Hospital
71 Nelson Mandela Drive
Mthatha
South Africa
5099

Sponsor information

Organisation

South African Medical Research Council

ROR

<https://ror.org/05q60vz69>

Funder(s)

Funder type

Other

Funder Name

Chemical Industries Education and Training Authority (CHIETA)

Funder Name

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Dr Siyakudumisa Nontamo (siyakudumisanontamo86@gmail.com) and Professor Eugene Ndebia (endebia@wsu.ac.za)

IPD sharing plan summary

Available on request